

102760" 0005660

					AC006080, AP001567, AP001567, and AP001567.
HTWBF58	2448	747706	1 - 1338	15 - 1352	BE177364, BF028245, AW798985, BF211148, BF208569, BF673866, BG259980, BE714535, AW798936, BF980362, AA758151, BE177404, BF812852, AI084648, AW069510, BF667616, BG223397, AW795505, AA601125, BF945606, BF874758, AA722039, AI889566, AI298061, BF965259, AL038498, BF726747, BF813686, AL133909, AI205010, BF857083, BF792316, AW939065, BE092200, BF931462, BE072376, AA584733, BF030591, BE092116, AL046549, AV702817, AW104793, BF854223, BE439876, BF678990, AL120083, BF986433, BF825540, AI524360, T41374, AW963750, AV703259, AI499301, BF916526, AW303196, BF725029, BE155414, AW238092, C18550, AI247199, BE077301, AA191659, AI002744, AI254980, AW854410, AV755654, AL537797, BF680286, AL044340, AV733830, AI436601, AW301350, AW961160, BF853276, AV654705, AW440215, AV759632, AW274772, AL048135, BF874773, AA912287, AA601202, AI873990, AL042853, R61498, AW971114, AV734120, BE902217, AV704541, AA663512, BG152927, BF904472, BF347791, BF974349, AW854498, BF342223, AF304447, AF275945, AC005254, AL035413, AC006512, AC004531, AL121890, AC037193, AL391260, AL357992, AL137784, AC008482, AC004491, AC009481, AC004841, AL035404, AC006012, AC005881, AP002898, AL031283, AC002350, AC020610, AC004019, AC008372, AC007242, AL049779, AL139109, AC005251, AL157938, AC066583, AC008041, Z98036, AC010748, AC005484, AL050348, D84394, AC005052, AC004967, AC010201, AF205588, AC007969, AC010605, AL137067, AC004895, AC006115, AC008403, AC005323, AL357752, AJ010770, AL136443, AL121871, AC011510, AC004158, AC007954, AP001717, AL355382, AC004813, AF001548, AC007619, AL035417, AC010150, AL031667, AL118520, AL117667, AC025765, AL035407, AL161666, AL117694, AC002089, AC008685, AL121658, AC002078, AC004887, AC002477, AL133282, AC009087, AL050311, AC006487, AC022027, AC005737, AL132773, AC002492, AL121747, AL353659, AC008042, AF055066, AL121578, AC022201, AL050335, AC000353, AC003957, AL136981, U47924, AP001870, AC012312, AC008265, AC007199, AC005962, AC004966, AL365505, AC005231, AC010083, AC020934, AC007782, AL354720, AL031664, AL157688, AC005043,

					AC005971, AC006013, AF053356, AC007963, AL359763, AC006077, AL031681, L81694, AC005837, AL096710, AC002563, AC008745, AP000356, AP000115, Z97056, U63630, AL139100, AJ277662, AL158823, AL035653, AL009181, AC008038, AF196972, AC000070, AL117336, Z95114, AF168787, AC005901, AC011445, AL022721, AC004789, AC005330, AL079304, AL136418, AC009244, AC005815, AL121774, AC005755, AC005731, AC006459, AL049709, AP001727, AC005011, AC058790, AC008999, AL109805, AL135978, AC012442, Z82203, AL049843, AC007386, AC083864, AL008719, AL022238, AL035683, AC007371, AC010553, AL356969, AB023048, AC005018, AC007541, AC004098, AC004996, AC002366, AC004125, AC005839, AL137230, AC004551, AC004149, AC022148, AL138706, AC004583, AL080243, AL109939, AC005015, AC007225, AC010462, AC004785, AP001718, AC007064, AF207550, U89387, U85195, AC011497, AC005480, AC003663, AP000313, AC021999, AC008271, AC005696, AL133247, AC004552, U96629, AC007546, AC007327, AC002565, AC004549, AC004778, AL009174, AL049795, AC020754, AL031427, AL138876, AF126403, AP001965, AC008969, AL163280, AC007688, AP000008, AC003104, AC005519, AC006318, AC073316, AL354864, AE000658, AL356379, Z98304, Z98752, AC020954, AP002792, AP002792, AP002792, AP002800, and AP002800.
HTWBO30	2449	655371	1 - 759	15 - 773	AA765129, AC013608, AC013608, AF235095, AF235095, AF235095, AP001574, and AP001574.
HTWBZ57	2450	828030	1 - 1653	15 - 1667	AL526540, AL532214, AL530769, AL526361, AL530770, AI038407, AL535007, BE737062, AI990446, BE736620, AI817294, BE736053, AI813752, BE674610, BF206639, AI887489, AW242426, BE904233, AW967040, AW188622, AV724936, BE169551, BE736681, BE787662, AA534945, BE539341, AI697490, AI970622, AW957112, BE393703, BE779956, AI826969, AI040380, AW512975, AI040037, AA779071, AA707247, BE787595, AW770455, BE695527, AI144222, AL535006, BE695442, AA749283, AL534456, AW019955, AI085178, AW022748, AI219163, AW192223, AI248630, BF110235, AA861436, AI206049, AI082771, AI038933, N24045, AA314458, AA452873, AI418455, AL534457, AV712871, H48028, BF724656, BF154707, AI095331, AI246185, AW572089, AA428132, AA428087, AA442266, W51755, AA281896, W06990, AW955681, N20861, AV712274, H80671, BE832400, H81664,

					AV757676, AA527742, W26241, AA084697, AA639542, BG006495, AA369695, AA862628, AA281739, BE243064, AA215459, AI745037, N27703, AA369694, AI299779, T77691, AI630540, H82924, AI219263, AW009819, AA369994, AA912335, AA610391, R35906, AW015184, BE764274, H60465, H88095, AA386027, AI630604, H53894, N80028, BF149156, N74576, BE243467, C02123, AI868490, AA995185, BE909040, BE831884, AI523294, H82925, H88096, BE764236, AI904772, AA317113, AA993010, BE674735, N35866, AV727421, BF369786, R89713, BF432789, H81665, AW383154, BE833623, AI383624, H53573, BE696239, AI904776, AA373117, N55673, BE699339, BG006486, BF830131, AA453043, H60420, AW389466, AA878674, BF992454, AI050888, N78248, N78305, AA489156, AR079021, AF132034, AF082569, AF113535, AF246144, U50734, AR083263, AC068724, AC068724, AC068724, AC009852, AC009852, and AC009852.
HTWCC10	2451	747687	1 - 1227	15 - 1241	R97267, R97030, AI061294, AA490845, AA632711, BF876324, AC002565, AC002470, AL122023, AC009516, AL359382, AC007688, AF131215, AC004231, AL117337, AF205588, AC018751, AP001753, AC009087, Z84484, AC009194, AC025470, AC005722, Z93017, AC018764, AC008733, AC004031, AL049569, AL031276, AL121906, AL118520, AL109797, AL033397, AL080314, AC007425, AC005088, AC003681, AC005049, AL031727, AL035494, AC021868, AL034420, AC018738, AC005577, AC008395, AC008372, AL022165, AP001670, AC019209, AC010618, AF134726, AC005519, AC004782, AC007284, AL008725, AC005815, AF224669, AL356748, AP001713, AC002302, Z92846, AC010358, AC006360, AC023511, AC008551, AC004382, AC004965, AF111168, AL161445, AC005249, AL035404, AC004847, AC007365, AC016397, AC083864, AP001760, AC011290, AC007225, AC006285, and U47924.
HTWCE14	2452	762839	1 - 1040	15 - 1054	BE621444, BG119409, BE621686, BF316579, BE267101, BE266139, AI369695, AI697690, AA557161, AW770378, BE208026, BG222463, AI206932, AI334278, AI830165, AW248509, AA027833, AI400954, AI469528, AI421560, AW194022, AI760695, AA760826, AW130878, AI249786, BG027713, AA171518, H05814, AA357204, AA027876, AI871059, BE746884, AA731891, AI208277, AA649957, AI694290, AW474641, AW474699, AI475699, AA350549, AW517662, AW957287, AA350548, AA676464, AI928706, AA352387, AA693347, AW005167, AW247418, BE728351,

				<p> BF526343, BE241425, BE242337, BE389611, AA665587, AI250627, AI364135, AI829495, BF814449, AI677797, AI860783, AA830821, AW058330, AI689614, AA514684, AI349012, AW189245, AI621171, AI824357, AI250646, BF830213, AL045859, AW163823, AA873151, AI471517, AW265004, AW152415, AI961393, AI050666, AI336575, AI811603, AW162071, AI203372, AI287476, BF915537, AL038445, AI439903, AI583578, AI699211, AW301344, AL040127, AW151034, AW087533, AI431962, AI702540, AW009066, BF725001, AW085786, AI679550, BE876766, AI097253, AW026633, BF726160, AI570966, AI826230, AI539071, AW262565, AI469516, AI089766, AI283996, AI689463, AI801032, AW089006, AW089258, AA769318, AI702902, AI371442, AW084915, AW963490, AI860694, AW161098, AI654258, AI886055, BG179295, BE875786, BE263582, AA916133, AW188438, AI285514, AW025464, AI872489, AI784028, AW162189, AI623596, AA858370, AI698437, AI699011, AI630947, AI370322, AI440444, AW131994, BE393551, AW900453, AI803740, AL046463, AV742931, AW858254, AI684164, AI815239, AI244380, AI744121, AI360560, BF793891, AI538850, AI373276, AW410878, AA832154, AI886124, AI249946, AW059713, AI744768, BF813177, AI434242, AI921266, AI468872, AI624962, AI270039, AW148303, AI589391, AI624604, AA648751, AI141288, AI474076, AI452993, AI624942, BE613727, BG106515, AI365256, AI679800, BG107420, AI559863, AI696621, AA279795, AI648502, AW071689, AB040950, X59414, AX046753, AK025209, AF067790, L10353, AF197929, AL122098, AF016628, AB049887, AB025103, AF061263, AL050172, AK026613, AK026583, AF218033, AF217977, AF065135, AK000559, AL137557, AF217998, AF089818, AF119857, AF114170, AF217993, AK026894, AK025092, AB044390, AF081571, A65336, AK000432, AF119909, AK025407, AR062106, I66342, AL137283, AB028451, AF155221, AL162062, E03349, AK027221, AL117635, AK026533, AL137538, AJ003118, I80062, AF266284, AF155148, AF113222, AL353938, M27260, AB047631, X62773, AL161802, AK026534, A32826, A30330, A32827, A30331, AL133636, AL137294, AF242525, AK025383, S54890, AK027116, AL137562, AK000074, AL049460, AL133015, AR029580, AB048910, AL122050, AK024601, AK027161, AB049629, AF159615, I17544, AF218023, AF208850, AL137268, AF019298, AF012536, E03348, AL359623, AK000652, AK024533, AK000636, AF217994, M77345, AF106934, </p>
--	--	--	--	---

0950003 01001

					AL137662, U95738, AK000568, and AF038191.
HTWCT76	2453	767758	1 - 1546	15 - 1560	AV764259, AV656063, AI345695, BF974756, BF801143, AA829044, AA644090, AI962030, AI863046, AA515048, AV709871, AI583466, AV762033, AI306232, BF854308, AA779783, AW958962, AA526326, AV755654, AW148507, AW471332, BF891585, BF832074, AI580707, AV764535, AW965058, AV764334, AV762633, AI434197, AA463590, AW189068, AW502237, AA225406, AA502991, AI251576, AV656851, AA833896, AA833875, AV711711, AL038606, AW272294, AA128511, AV713072, AA707849, T05834, BE044774, AV760014, BF852924, AW970940, AV711987, AW166611, AV738285, AW855730, AI933714, AA714011, AI254779, AA679937, BF915799, AW855736, AI254913, AI570943, AA714110, AV759559, AA581247, AI925579, AV695478, AW855803, AV762242, AA653139, AI688837, AA452887, N57681, AW975095, AA526625, AW130446, AA601356, AV763457, AV756491, AC005522, AL162424, AC007172, AC006449, Z85986, AC005512, Z97054, AL049795, AL021707, AC012384, AL033529, AC002395, AF241726, AC008119, AC004955, AC005305, AC009955, AL080243, AX033912, AD000813, AL138919, AX033911, Z83840, AC005235, AF155238, AC007649, AC010643, AC005247, AL133294, AC007036, AC010150, AC004097, AC009470, AF222856, AF042484, AC007056, AC005052, AF222854, Z93023, AF222855, AL356057, AF168787, AC023105, AC004686, AC018758, AC005479, AC078843, AC083866, AF117829, AC003967, AL133244, AC011514, AC005914, AL445248, AC011482, AL021154, U91326, AL121897, AC024085, AC005955, AL021807, AL138836, AC018719, AL122023, AL161422, AP000299, AL138741, AP000104, AC004099, AC011462, AL031295, AC004595, AL445483, AC007160, AL035683, AL049776, AF069291, AL121905, AC006468, AL022165, AL109627, AC009154, U50537, AC008636, AC011489, AL121891, AC024075, AC011811, AL079342, AC002067, AP000113, AP000045, Z97985, AC006345, AP000497, AL160175, AK024160, AP000270, AX039602, AL035400, AC005220, AC009600, AC006023, AC007551, AL139343, AC020906, AC005626, AC006441, AC005796, AC009194, AC010605, AC007385, Z81364, AC002116, AL050318, Z84466, AC004824, AL139188, AJ400877, AL136300, AL162584, AC004922, AC005409, AC003969, AL161670,

					AL365505, AL163280, AC004510, AC004947, AL035462, AL138755, AL117382, AL162377, AL035670, AC005562, AK001397, AP001716, AC005933, Z82215, AC006111, AC000073, AC010251, AC004156, AC003104, AF001549, AC007539, AF258547, AP000032, AL121949, AC004778, AC006080, AC004525, AL035405, AL355809, AL135839, AC002073, AL390028, AC008733, AC002544, Z75888, U51244, AC005258, AL121652, AL049839, U71210, AC005878, AP000691, AL121893, AC008115, AC008372, AL133448, AC004883, AC005589, AC007383, AL135912, AL049632, AC000070, AC002045, AC010677, AL138875, AL049764, AC006329, Z98304, AC009123, AC020913, AL117377, AP000159, AL121916, AC004382, AC003982, AL050313, AL137073, AF196970, AC004500, AP000289, AL121712, AL354866, AL022162, AC005837, AL031282, AC008033, U96629, AC016995, AC008474, AC002039, U80017, AP001725, AL139087, AL161731, AC008812, AC004841, AP000110, AP000043, AP000111, AP000350, AC005104, AL022145, AC011455, AP000258, AL121886, AL023284, AC008699, AL035458, AL163202, AL034376, AL020998, AL109976, AL049766, AC010326, AC004867, AC008379, AL157713, AL049653, AF020503, AC008736, AC000159, AF064858, Z68162, AC067721, AC067721, and AC067721.
HTWDJ17	2454	784037	1 - 1376	15 - 1390	AV686983, AV646121, AW961968, AV646138, AA424670, BF700169, AV659105, AA004250, AV660261, AV658479, H72674, AV658573, AV660130, AV660299, R00934, AV683433, AW501010, AW501128, AV646353, AV646230, T08315, T77004, AW796868, AW976358, AI538436, AV707357, F13040, F12949, AA699932, AA761228, AI807087, AA766036, BE549529, W27156, AA761066, T75185, AI866469, BG110517, BG029667, BF061529, AI433157, AI554821, BG252929, AW151136, AI539771, BE897632, AI537677, AI494201, AI500659, AI815232, AI801325, AI500523, AI582932, AI872423, AI923989, AI284517, AI500706, AI491776, AI445237, AW151138, AI521560, AI889189, AI500662, AW172723, AI284509, AI889168, AI440263, AI866573, AI633493, AI434256, AI805769, AI434242, AI888661, AI284513, AI888118, AI859991, AI436429, AI889147, AI371228, AI440252, AI866786, AI860003, AI610557, AI242736, AI887499, BF812963, AW858243, AI538850, BE885490, BF812936, BG257535, BF338002, AL045500, BF795712, BE886728, BG260144, AI890907, AI539800, BE965355, AI866465, AV743631, AI538885, BF910810, BF812438, AI371251,

095003-09101

				AI274759, AI866510, BF525578, AI923046, AI500714, BF921092, AI355779, AI499986, AL047422, BE963918, BE964512, BF811804, AI433976, BE895765, AI690835, BG108334, AI648567, AI539781, AI366900, BF970652, AI476086, BF968504, AI581033, AI559957, AI432666, BE963035, AW197139, AI539707, BF814409, BG113712, AI348880, AI521571, AI275175, AI568138, BF814541, BE964959, AI608805, BE964576, AI499463, AI920968, AI280670, BE964767, AI610362, AI620284, AI440239, AI521596, BF727352, BG120046, BE965724, AI537273, AI436456, AI371265, AI963846, AI744185, AI567940, AI610357, AI817244, AI491710, AI612913, BF904178, AL119319, AI285826, AI863014, AI521594, AI499512, AI889133, BF526393, AL042365, AL042551, AL042787, BG029053, AL042572, AL513743, AI610402, AI434223, AI697243, AI610429, AI628850, AI828574, AI433968, BF812961, AI539632, AL515375, BE895585, AI889148, AW118237, AL042377, AI539847, AI872074, AI828583, AI887775, BF987113, BE875407, AL042538, AI872300, BF968907, AW172745, AL042557, AI538878, AI354998, BG179586, BG111590, BG121959, BF968622, BG114012, AI567935, AI805762, BG165260, AI432656, AW268060, AI308035, BE895801, BF036448, BG164752, BG033403, BF347093, BG034554, BG029194, BF983875, AI349967, BG180046, BF038804, AW302973, BG181100, AI572717, BG024457, BG116926, BG261116, AI475377, AW151485, AI049851, BF525811, AI690946, BG178146, BE537531, BE963960, AW191003, AI225248, AI432644, BE784387, AI339388, AI554827, AI801619, AI561170, AV745008, AX040958, AX040974, I89947, A08916, A08913, I48978, AR038854, AK026086, AF032666, A08910, A08909, I46765, A93016, AL122049, AB048953, X70685, AF113222, A08912, E12747, I89931, AK026865, AL080086, AR087170, A08908, AK026627, U66274, AR079011, AR079012, AL162062, AL133665, AL389939, AK000718, AK024992, AF271350, I96214, AR034830, AL137271, AK025414, AK027204, M86826, AF185576, AJ012755, AF116682, AB049892, AL049466, AL162006, AL133072, AL137283, AK000212, AK025524, AF085809, AL137557, AF119875, AF143723, AB052191, AF130077, AF090896, Y07905, AL110159, AL133560, AF118094, AF130055, AK026642, AK026784, AF153205, AB048975, AL117460, AL359601, AK026927, AF090901, AL389935, AF081197, AK025383, AK026462, AL122110, AF067728, L13297, AX042059, AL049283, AK026480, AL133080, A77033, A77035, AF087943, AL133081, AK026744, AX045627, Y11587, AB050431, AF090903,
--	--	--	--	--

					AL137533, AL133077, AL050146, AL353940, AL133049, E15569, AL137292, AF081195, E07361, AL353625, AL137705, A18777, A08907, AK024538, AL353957, AF110640, AF130105, AL359941, AL133070, AF217987, I48979, AF130104, AK027114, AF116676, A65341, E04233, AB041801, AJ299431, AB048919, AX005848, AL050172, AK025092, AF116654, AX005804, E06743, AL117585, AK025632, AL080158, AF262032, AF130082, AF130110, AL122123, AB034701, X72889, AK026532, AK026408, AK027113, AB051158, AK027096, AK026528, AL137478, AF217966, AR000496, U39656, AK026626, Z82022, AB050533, AL117416, AK025435, AK027160, A93350, AF026124, AF116602, AL110221, AB052200, I09499, AF125948, AF158248, AX010492, AL137550, AL050393, AL110222, AL122121, X83508, AL137476, AB049848, AK025889, Y10655, AR013797, AK026550, X82434, AL137526, X87582, AL050024, AL049430, AF113699, AL133640, AK024601, E02349, AL359596, AL117583, AK026741, AK000083, AF111849, AF219137, Y09972, A90832, AL117457, AK026452, AF177401, AX027129, AL353956, U78525, AF106657, AL133113, AF008439, AF285167, AK000450, AL390167, AK026542, AL049938, AK025857, I17767, AR059958, AK026950, AF130087, AF090900, AL096744, AR081734, U58996, AF061943, AF113013, AK026591, AL121656, AF119337, L19437, AL389978, AK025391, AK025375, U30290, AF061795, E08631, AF151685, AL133568, AL137488, AK026506, AF000145, AK026647, AL133112, AL359618, S78214, AK000486, AK026600, U00763, AL137300, I89934, AK026593, Y10080, AF111112, AF078844, AL080060, AF090934, AK024588, Y16645, AK025967, AL080074, AL110196, AK026583, AL359583, and AL137648.
HTWDM89	2455	569787	1 - 1458	15 - 1472	AI358384, T06598, AW965008, AC016627, AB002353, Z97832, AL022302, AL117333, AC007225, AL121895, AC005082, Z98941, AC005332, D83253, AC018663, AL121903, AL049589, AL050335, AC011450, AF196779, AL158830, AL031681, AP000501, AC005821, AC009086, AC006345, AF196972, AC009032, AC004167, AL021407, AC011475, AL353804, Z94056, AC007637, AL163281, AP001628, AC008760, AC005049, AL109839, AP001754, AF228703, AL132826, AC009060, AC003982, AL139100, AC000353, AP001717, AL078477, AL136137, AL158040, AL035086, AF053356, AL135960, AJ131016, Z95152, AC000004, AP001747, Z98036, AC003007, AC018644, AC005052, AC008736, AL355476, AC005484, AJ400877, AL049697, AC005015, AP000510, AC007687,

0950083-091201

					AC006241, AC011465, AP000513, AL121585, AC005527, AL022721, Z94801, AL049780, AP001745, AC007722, AC002511, AC018731, AL391601, AL133229, AP001711, AC005529, AC007546, U91323, AF195658, U80460, AP001760, AL031733, AL138812, AC005288, AC010422, AL353807, AC005004, U91322, AC008041, and AL022336.
HTWEA05	2456	844386	1 - 879	15 - 893	AW878238, AW827205, AL045891, AL138455, BF337291, AV756220, AL514075, AW858522, AL042753, AV760019, BF923684, AI064787, AA521299, AV760466, AV682853, AV698043, AL042898, AL043089, AV733824, AL046356, AL135012, AL043321, AL047163, AV762645, BF347269, AW963250, AL042488, AL041862, AL514791, BF342223, BF816035, AW021195, BE906410, BF990285, AA577696, BE904906, BF996894, AL045327, AL513991, AL047611, AL513979, AL514627, BE965169, AL043052, AL514357, AI440133, AL042515, AL134524, AV681613, AI432644, AL513817, AL040207, AI431307, AL040459, AI636719, BE252421, AL047092, BE966577, BF970536, AL035847, BF968622, AA598742, AL045328, AL043091, AW080565, AL514691, AI431316, BE070818, AL047675, AL079977, BF868994, AL514219, BG106270, AL514449, AL513693, AL514415, AL042853, AA760655, AL042538, AL040038, AI432666, AL047037, AW967257, BF527274, BG180622, AL515293, AL514511, AL042745, AL514983, BE966547, BG256324, AL119748, AL042729, BE893169, AV662292, AL514487, BE047915, BG251851, AV733431, AW858243, AL037683, AL042787, AI285419, AL043196, AW959837, BG110517, BG113493, BG029667, AV729953, BG033220, BF828567, BG181066, AI433157, AI648567, AI690946, BG252929, AW151136, AI539771, BE897632, BF925370, AI537677, AI494201, BF797072, AL043239, AI500659, BE883591, AI866465, AI815232, AI801325, AI500523, AI538850, BE885490, AI887775, AI582932, AI284517, AI923989, AI872423, AI590043, AI500706, AI491776, AI445237, BF965904, AW151138, AI521560, AI889189, AI500662, AW172723, AI539800, AI582912, AC007172, AC002542, AL121601, AC004837, AC021068, AL033521, AC021019, AL358133, AC020923, AL355103, AL034422, AC002394, AC011484, AL353807, AL109753, AC004594, AC022402, AC007114, AC006313, AC002287, and AC002457.
HTWEG06	2457	762856	1 - 1052	15 - 1066	AA856945, BG110768, AI803741, AF075680, AA810477, AI923265, AA340226, AA102707, BF800649, BF678590, AI890324, AI559720, AC006111, AL391724, Z83844, AL050335,

					AL031662, Z69707, AP000553, AC005513, AC004166, AC007686, AC006946, AP000116, AP000557, AP000269, AP000563, AC002395, AC005225, AL354720, AC011453, AP000555, AP000103, AL034549, AC005098, AC005779, AF190464, AL035587, AL133286, AP000212, AP000134, AC009079, AL121983, AC004167, AC004129, AP001711, AC006312, AC018511, AL138733, AL031587, AC009060, Z95114, AL135905, AC005480, AC016898, AF001548, AC006965, AL109804, AP001717, AP001714, AC004854, AC000353, AC007404, AL078581, AC004858, AC020916, AC011455, AP000030, AC010328, AL158828, AC005670, AC010205, AC004971, AC007308, AC003041, AC008953, AC005911, D86995, AP001726, Z93241, AC006530, AC004491, AP000689, AP001719, AL096703, AF111168, AC008039, AC004765, AP001725, AP001694, AC010618, AB023049, AL133246, AC009247, AF254822, AC005522, AC007011, AL136105, AC006333, AL160231, AC008403, AC004447, AC010285, AC011500, AL354861, AC006388, AC011479, AC007021, AF109907, AP001724, AC006484, AC003684, AC010326, AL162430, Z93930, AC009086, AC005280, AL136358, AJ009616, AB003151, AC008848, AP000133, AP000211, AP000032, AL096840, AL132639, AC007956, AL138836, AC005722, AL096870, AC005300, AC000026, AP000240, AC017100, AL121928, AL139100, AL139396, AC002404, AL049840, AL031283, AP001728, AC006057, AC025575, AL132855, AC006449, AL031727, AF126403, AL035669, AC005529, Z98946, AL121655, AP001712, AC002551, AC007057, AL031433, AC020955, AL031846, AP000696, AL022318, AC007536, AL022316, AC005081, AC005211, AL020993, AC008556, AC004907, AC006285, AC003688, AL157372, AC005531, AC023880, AF030453, AL031284, AC023490, AP000692, AC020603, AC004895, AC027644, AL138919, AC010271, AC022073, AL138741, AC005356, AC006252, AC006430, AP001753, AC005666, AL163853, AC027279, AC006001, AC002306, AP000354, AL121905, AC007114, AC008784, AC005690, AL021918, AC005484, AJ277546, AC004883, Z98949, AL135927, AC007227, AL136228, AL034372, AL035079, AJ295844, AL139384, AC004973, U47924, AC011477, AC016025, AC004216, AP000359, AC005913, AC008040, AP002408, AC021325, AC021325, AC074288, AC074288, and AC074288.
HTWEQ36	2458	823369	1 - 1422	15 - 1436	BE740976, BF688976, BE388462, AL515860, BE618810, AI144183, BE619718, BE743638,

					AI753326, AI123548, AW148675, AI762763, AW665598, AI991105, BE246225, AW151784, AA632384, BF970549, AA687754, AI360286, AW731898, AI281497, BE206192, AA662315, AW173365, AI304830, AI581361, AA653146, BE409628, AI093405, AA837430, AA989346, AW129580, AW079976, AW589241, AA888131, F24031, AA765564, AA161018, AA197309, AA149756, AI554179, AI439376, BE963919, BE243366, AA826614, AA088666, AA552595, AI191165, AA993464, AW901000, AW241223, AW519257, W92934, AW473246, BG152577, AA132941, AA768562, AA160619, AA976014, AI346746, AI538906, AA132715, AI640891, H22143, AA135107, W92935, R48285, BE545547, AA677271, BF316389, R74055, T65148, Z28491, R43359, F02594, D53802, AA743537, BF805352, Z41473, AA831142, AA524214, F09495, AI872452, AA827021, F36480, BF213155, AA975550, AI671936, AA366296, AA992018, AA652934, BF837632, AA585480, BF342033, AI133151, BE255649, C00346, BE707838, AI908756, AC000394, AF072711, AK000194, AY007123, AK024510, AF133124, AL162417, AC000394, AC000394, and AC000394.
HTWFA21	2459	824551	1 - 670	15 - 684	AA905353, AI093811, D60081, D60080, AA035076, AA035493, D81399, AA771809, AL162385, and AL162385.
HTWFA88	2460	596942	1 - 1837	15 - 1851	BF035327, AL038785, AW972871, BF684828, AW088202, AV720371, AA522942, BG029216, AI570261, AI679782, AW977134, AA523815, AV763777, AI625244, AI368745, AW162049, AI929531, AI339850, AW969698, AW969694, AW029038, AI962050, AA496260, AV729947, AV728632, AA837084, AV719311, AW088616, AI251436, AV728309, AV719316, AV710306, AI918421, AI567674, AA482711, BE389903, AA713815, AI251002, AU148742, AU145462, AA582911, BE787738, AA768905, AW166815, AI049722, BF677892, AA634146, AI133636, AW440836, AI890923, AA504871, AW104748, AI917271, AV752696, BE390977, AA488573, BE277602, AW080134, AA826303, AI499938, AI469003, AW079659, AA111909, AA873560, C06458, AA729721, AA767376, AL120674, C06327, AI608626, C05755, AI805607, AW075511, AV734666, AV760777, BG254180, H78195, AA644207, U51696, BF819858, AA550758, AL046409, AI284640, AV763354, BE744242, BF675718, AI431303, AI904894, AI064864, AA312845, BF211188, N22018, AV760937, AV762098, AL119391, AV762139, AA132833, AA483068, AV761056,

0950083-091001

				<p> BF668217, AV735370, AL515495, AW193265, AV709064, AW303196, AW274349, F36273, AA486885, AW500125, AI064952, AI334443, AI270117, AI358571, AV710066, T08438, BG249643, AW301350, AV761362, AI568678, AA487858, AV764578, AI963720, AA491814, AA531372, BG036934, AW954829, AV740801, AI350211, AV764307, AV762050, AA652764, AW872676, AF330238, AI613280, AI281881, AV728425, AV762111, AV761489, BF241967, AI305766, BF940837, AI610159, BE387744, AA113186, AA719080, AV759274, AI281468, AV763255, AW473163, AV761786, BF679274, AA490183, R95704, AW438643, AI457397, AI821271, AA469451, AV735495, AA151980, AW872575, BF697673, AU147800, BE042649, AV763540, AA196796, AL138265, AV763971, AI254615, AI345654, AV725423, AL138455, AV749274, AV742057, AA491284, BE259894, AA115968, BF678329, AW513362, AW327868, AV764329, AW974109, BE047069, AA620464, BF130107, BE350475, BF820241, AA581903, AW503666, AA623002, AW517377, AV760774, AA715217, BG171096, BG177307, BF854876, AW020992, BG180305, AV762826, AW833862, AV764241, AA857486, AV741609, AV742309, AV742337, AA720702, AV652936, AV761925, AI053672, AA908687, AL119691, AI754955, AV759382, AA507547, AA595825, AA548689, AA480790, AW965008, AW021583, AV682003, AW419262, AW502975, AW276827, BF541116, R91159, BG109996, AI471481, AC005969, AL354766, AL353706, AL132987, AP000426, AC010386, AL022318, AC006050, AC024239, AL035467, AL158093, AC007611, AP001728, AC025472, AL162430, AL357774, AL109653, AL136134, U49973, AL023755, AP001689, AL158159, AC003081, AC015600, AL365193, AL136992, AL137787, AC016395, AL139331, AC007130, AC008155, AC004158, AL138756, AC022163, Z98745, AC002981, AC005953, AC002123, AC006006, AC006080, AC005678, AL136137, AL354942, AC005537, AC004019, AP001053, AC004941, AL359763, AL135841, AL121944, AC011299, AC004103, AC009225, AC004069, AL162872, AC000004, AL355178, AC005013, AP001752, AP000354, AL121983, AC006325, AC009406, AP000355, AL137918, AL050335, AC016623, AC002303, AP000043, AL022159, AC000092, AL121591, AC008102, AL157791, AC010485, AL138731, AC011456, AL353643, AP001716, </p>
--	--	--	--	---

09950083 091201

					<p>AP001676, AC004764, AL031585, AL031768, AL357041, AL163279, AC006975, Z72001, AB020869, AL022717, AL096712, AL022578, AL031602, AC008831, AL139182, AL356513, AL049835, AC020939, AC009481, AC006566, AL135905, AC004014, AC002369, AC073316, AP001818, AL353573, AL031671, AC026341, AL096861, AC006346, AL160237, AC007214, AC008598, AC004845, AC027663, AL139188, AP000885, AF130342, Z93016, Z84719, AL157915, AL158040, AC004828, AL359218, AL096821, AL109984, AL365444, AP000516, AC006500, AP001051, AC018719, AC005544, AC006334, AC005145, AP001708, AC000086, AL136131, AC005837, AC004067, AL031622, AL035634, AC004126, AC083871, AC003667, AL137878, AC009275, AC024167, AL132826, AL121753, AC012081, Z70049, AL117334, AC004413, AL096816, AL133283, AC006984, AC009511, AC006480, AC007380, AL031677, AL078460, AP000493, AC007026, AL049650, AC008124, AE000660, AP000470, AC016144, AC006039, AL133391, AC023134, AL139286, AL031287, AC004142, AC005828, AC009331, D83253, AL034410, AC005358, AL133517, AC016568, AL109830, AL138690, AC000094, AC004049, AL034551, AL139375, AL122020, AC004980, AL050321, AC007652, AC009405, AL031276, AC002377, AL117354, AC006965, AC020550, AP000009, AC018642, AL021408, AC068658, AC005592, AC007316, AL136179, AP000072, AC005411, AC006213, AP002028, AC019227, AF064864, AL139009, AC022392, AL359382, AC007282, AC011738, AL079342, AL136419, AC005513, AL034376, AC016620, AC009483, AC006060, AC019106, AC007384, AL161781, AL163853, AC018448, AL022336, AC007097, AL034343, AL133396, AL161666, AL138894, AC069275, AL162811, AC015971, AF181897, AL163283, AC005562, AC000370, AC002995, AL158830, AC007326, AC007671, AL121782, AC083862, AL136987, AL391869, AC073934, AC000359, AC025201, AC023008, AC053473, AC041018, AC021959, AC026240, AC011175, AL365218, AC022695, AC005664, AL163193, and AL163193.</p>
HTWFM85	2461	844677	1 - 1679	15 - 1693	<p>AL530147, C75513, AI796284, AA425431, Z42827, AA973025, C75646, H92159, AI017480, AA164842, BE004359, AA151143, AA047527, BF734980, AA114885, N50209, AI623294, AI341910, BE720812, N32573, N32594, AB046820, AC005881, AC005881,</p>

095005660

HTWFO43	2462	858741	1 - 1284	15 - 1298	and AL137025. BF792326, AF074667, AV700988, AV700498, BE796439, AV756726, BG164166, AV762783, BG260565, AV733710, AL040921, AV759172, AV762145, AV700545, AV764490, AL135377, AV710705, AW965008, BF339640, AV699709, AV762693, AW600804, AV734607, AV763135, AL138265, AI076616, AW820787, BE071877, AW406162, AI924251, AU118745, BE071876, AW088224, AU117456, AV714931, AV760723, AV760599, BE908796, AV704375, BG250302, AA604395, BF828714, AW820795, BE387734, BG032943, BE538259, AV759356, BF826830, BF915799, AV761286, BG177715, AW820127, AW406447, BF968610, BF906248, AV731286, AI963600, AA836811, AV729960, AU159301, AI567674, AA631507, AW968187, AV759362, AV762900, AV762902, BF906239, AL121235, BF913258, AV759711, AV760364, BE740358, AV763174, AV700663, AV762022, AI016704, AI038990, AV759683, AA515829, AW245747, AW969988, AV701844, AV704740, AV759686, AW961994, AV762779, AV762001, AV761309, AV759684, AL120343, AW243793, AA908687, AA837740, AV701613, AW953071, BE969647, BF805173, BE541237, AU146953, AV764388, AW079792, BF965406, BF965477, AA502104, BF337291, AU121297, AV764389, AV699393, AV762781, AV759682, AV763683, AV759685, AV762395, AV763154, AA722372, AU156861, AW474168, AL120483, AL038705, AV764387, AV763892, BF213459, AI204304, BF736198, AV763608, AC005280, AL357153, AC008372, AC004019, AC008569, AL034372, AL121897, AC004675, AC004638, AC000052, AL158830, AP001725, AL049759, AP001666, AF088219, AC002350, AC008519, AC018751, AC003007, AC005971, AJ009611, AC005696, AC005632, AL354720, AC009516, AC005722, AL117352, AC005324, AC004971, AL353807, AC000025, AP001731, AL035587, AP000046, AL159977, AP001760, AJ295844, AL109804, AP000114, AC008543, AC007227, AF001549, AC005527, AL135927, AL163249, AC005520, U80017, AC006449, AC004796, AL355385, AC027689, AC008155, AC007216, AC010739, AC005412, AC005180, AC006195, AL096840, AL023575, AP000553, Z82244, AC005839, AC009247, AC006285, AL035422, AL133215, AC027644, AC005288,
---------	------	--------	----------	-----------	--

					AC005771, AC011455, AL023284, AC002470, AL163282, AP002534, Z93017, AC002996, AC007043, AL022238, AP001688, AL080243, AC005859, AL049830, AL353194, AF134726, AC023274, AC005740, AC004686, AF123462, AC004884, AC005529, AC005081, AC009086, AL021154, AL137073, AC006501, AP001684, AC018644, AC007298, AP000402, AC006071, AJ400877, AF029308, Z93241, AJ277892, AP001689, AL031257, Z74739, AL118520, AP001691, AP001714, AP001672, AL137818, AD000092, AL356299, AL109965, AC011475, AL121658, AC005911, AC005516, AC004382, AJ009616, AL158040, U95742, AP001687, AC005231, AC037193, AC011477, AL035411, AC004690, AP001346, AC005154, AC007562, AP001685, AC004166, AL138759, AL021578, AC021752, AP001717, AP001732, AC005046, AC004972, AC008064, AL021368, AL133153, AC002041, AC006480, AC011489, AC009228, AL022328, AL163247, AC009060, AP001667, AP001695, AP001670, AC005246, AL135744, AC011464, AC006213, AC079610, AL391867, AL121653, AP000359, D84394, Z97054, AL136980, AC008770, AP001753, AC008537, AC020904, AC007097, AC008039, AL031650, AF196969, AL049766, AP001727, AC024082, AL008721, AP001671, AP001679, AC004673, AC010879, AP001681, AC002429, AC007308, AC006111, AC005821, AC004980, AL356244, AP001759, AC013737, AL078461, AL133332, AC006441, AC011482, AC018720, AC009194, AC004887, AC005255, AC022137, AL035450, AF314058, AP001698, AC006028, AC011491, AC018639, AC007537, AL133551, AC005531, AC021868, AC019209, AC005392, AP001708, AL354915, AL445687, AL008718, AF109907, AP001715, AP001699, AC068993, AL109759, U62317, AP000020, AL163207, AC016026, AF042090, AC002059, AL109865, AC004933, AL163278, AL161757, AC008745, AL008629, AP001172, AJ011930, AC025593, AL163300, AL117693, AC006998, AP000513, AC035150, AL031281, AC005522, AC010326, AP001694, AC005013, AC009953, AL117336, AC027319, AC009487, AP001680, AC034243, AC068755, AC026936, AC027264, AC011962, AC040904, AC040904, and AL353694.
HTWLG39	2463	853406	1 - 490	15 - 504	AA813011, AI091612, AI204546, AI560743, AI418276, AA858118, AI560812, AI992293, AA522719, AW130435, AA402002, AW051814, AI291876, AI362691, AA905625, AW020847, AI356940, AW008254,

					AI149942, AV729360, AI703226, N55455, AI221349, AA975506, AI240036, AA970179, AA885640, AI685154, N80142, AI283256, AW028863, AW261910, AA883234, AA865830, AA765503, AA906638, H95085, AA995348, AW512013, AI240974, N35116, AW128861, AI025840, H95663, AI346239, AI207741, H24210, AA738193, H54683, H94256, N20084, D59957, N23206, N29785, AW661747, BE644944, AI089271, AA063258, H38653, AI027068, AA443008, AI520954, T78586, AI359626, H96792, H96239, H18631, AV648478, N90414, AI362623, BF083306, D60276, BF083308, AI962075, D80608, R54942, BE044404, AI985444, AW026456, AA894582, AI800639, AA972097, AA609747, H53597, AI804799, BF083311, R20753, D59884, T56966, W96084, N22716, H48906, BF083237, W96085, H91182, D52425, T27025, AW852853, N79026, H67369, BF083234, AW797454, BF083282, AA761468, AA815412, H54445, BF083309, Z28499, AA846821, AA972438, BG058544, R20754, AA970691, W19987, R89812, H73921, AA811823, AI671824, AA235389, F10605, AI923477, AA236543, AI370536, H69003, AA427749, AA384679, AI300180, AA301182, AA492083, T79403, AA232733, AA768615, AA505078, BF980167, AA282083, H60026, AA701943, T86974, AA247535, R08289, H61736, H54573, R94479, H38643, BF790455, BF212832, AA854918, AI371632, BF057907, AI609846, AI349670, C01267, AI252251, AI345661, BG170214, T90553, R94069, BE043757, N56269, BF055861, AW302033, AB018288, AJ297360, and AC023288.
HTXAA20	2464	821342	1 - 747	15 - 761	AV730383, AV732734, AV731010, AW139425, BF807866, AI590899, AI199332, BF858888, AI343963, AI040225, AV730097, AA947175, AA640032, BG230657, BE247164, AK024449, AC016823, and AC016823.
HTXAD75	2465	745410	1 - 1910	15 - 1924	AA634116, BF570310, BF793193, BG257515, AU134921, BF980040, AW194709, AI983851, BG250091, AI064728, BE889585, AW071012, AI660958, AU123297, AI766131, BG249846, AW148839, AL040880, AV707299, BE671248, BG251065, BE893272, AW469227, AI982831, BE882137, AL537034, AW338696, BE878531, BE081362, AI042399, AI583975, BE327675, AI096908, AW305017, BE219767, AW612867, AA665917, AW363287, AA522903, AI696887, AI050865, AI587311, BF240570, AW363286, AW769190, AA523283, AA582781, AA494509, BF448151, AW363297, AI005382, BE698583, AW363298, AV709045, AW129288, AW237848, AW131711,

					<p>BF939460, BF351450, AW590749, AW190991, AA563896, AW363278, AW363290, AW374132, AW513093, BF572301, BF216042, AW078702, AU154337, AA633098, AI678882, AA983260, AI623366, AA872788, AI183629, AI129791, BF030649, AA836948, AW008817, AW517993, BF433652, BF667783, AW239528, AI192887, AA487970, AA970798, BE502093, AI587536, AA102454, N57489, AA043263, BF246636, AA977904, BF593235, AV725022, AA155612, AW316964, AW170327, AW950029, AI445245, AI189089, AA603286, BF433643, AW273766, AI168534, AI744547, AW339027, AI354742, AU150339, AA724163, AI692612, AA962012, BF594651, AI354743, AI955531, BE439411, AI707977, W52119, BF572649, BF440046, AA125892, N75252, AW250965, AA146599, AA847445, N58837, W37586, W39332, AW191016, AA487819, AI921182, N29228, W63799, AW089771, AA131636, AI635094, AI769687, AA968495, BE000945, AI160953, W42875, BF448699, AA992033, N52953, AI751412, AA041207, AA226668, AI334598, AI273080, BG118888, AI681816, AW005589, AA557710, AA125891, AI634026, AA137112, F08088, T59704, AA862232, BE856165, AW992584, W42939, BE175814, N50141, AA137003, BE763173, BE241800, AW779308, W37461, AA137030, AW134572, AI587433, N49080, AW518458, BE938814, AI765373, AA225057, AA025585, AA112008, BE001081, BF879612, BE156363, F04325, R77217, AW263897, AI476767, H64546, BE698074, AI801684, BE001762, BF216620, BF922676, AW510569, R77216, H97581, AA040753, BE001874, AI522014, AI985798, R54502, AA522871, F13336, AA083727, BG163467, AI950665, AI469785, R39609, AA584935, AI355766, AW023663, T78816, BE814685, AA114916, AA780920, AA158376, F13337, F10932, D25892, AW085745, BF815310, AW380152, AA581970, R14160, AI564130, AI928248, BG119830, AW361788, AA026947, AA136892, AW576773, H64977, AW801938, R44942, T27830, BE882921, AW747841, AI672161, R56033, BE149521, T56465, AW363296, H64646, C21384, AI089162, AA303992, AU141061, R54403, AA569174, BE000830, AR020616, AF261089, M23254, AF284441, U71320, D38117, AF015038, L09120, Y10139, AK023851, AR018788, U07850, AC021798, and AC021798.</p>
HTXAR92	2466	656935	1 - 1586	15 - 1600	<p>AV762633, AV738383, BG180976, AA916430, AV756491, AW974932, AI523316, T74524, AI890324, BG029528, AW270385, BE138594, AI254779, BE062159,</p>

09950083 091201

				AI133083, BE063437, AL036665, H07953, AI733856, AW084445, AW865946, AA501461, AW068596, AI687343, AA825827, AA456924, BF857849, AW303196, AI800180, AI457397, BF675251, AW301350, BE150793, BE968744, AW274349, AW514662, AV762430, AA714110, AV759632, AW970981, AI696962, AV760915, AA515351, AW069769, AI223968, AL118991, AI859438, AI279417, AL119691, BE139267, BE155951, AI800168, AW410354, AW510513, AA683069, AW193265, AA640430, AA631507, AI963720, AV710066, AV695478, AI912401, AV719392, BF854308, AW270258, AW026305, AA833875, AA833896, AI421950, AI419337, T50676, AI334464, AI344812, BE062478, AA640410, AA477503, AW502237, AI280504, AV754716, AA578621, N35306, AV715310, AW575000, BE501593, BE139358, AW504224, AI216990, AA084609, AC010150, AL138836, AC011497, AC005409, AL078638, AC004840, AC003108, AL035704, AC011489, AF207550, AC007956, AL034420, AL353807, AL137119, AF001905, AL135927, AC004087, AL022313, AL009181, AL121758, AC008379, AL009172, AP001330, AL445687, AC007878, AC020904, AL133289, AC007739, AL021939, AC005800, AL035409, AC006312, AP001631, AL133445, AC011464, AC002303, AC004805, AL121825, AC005318, AL121895, AC009228, AC002472, AC004526, AP001709, AC011475, AL121586, AC005052, AL049843, AC004824, AC005098, AL138894, AC007766, AP000127, AP000205, AC005031, AC008745, AC007664, AP001748, AL034548, AC004867, AC007041, AL109758, AF165926, AC007011, AL109823, AL163279, AL078477, AC004216, AL008627, AL133227, AC009516, AL139328, AC008569, AL445483, AC023880, Z93016, AC007528, AC002352, AC002395, AD000092, AP000547, AL138976, AC004905, AC005940, AC010205, AL049569, AC007227, AC004149, AC002412, AL078604, AC004963, AL391122, AL121776, AL033520, AC022148, AC008496, AC005015, AL132987, AC004983, AL137162, Z83847, AL137005, AL022159, AL022323, AL022333, AL138878, AC008747, AL031597, AC004166, AC007151, AP001711, AC020934, AP001715, AP000193, AC008649, AL022302, AC027319, AL033527, AL078581, AC005255, AC012318, AC010170, AL031291, AL121897, AF217403, AC008784, AC005378, AC010219, AP000031, AC005225, AF112482, AC005332, Z83844, AP000117, AC010271,
--	--	--	--	--

					AC009502, AL049692, AL136109, AC007220, AC004156, Z82189, AC005933, AC007731, Z84466, AC005500, AL161670, AL008583, AC008760, AC005288, AL133246, AC027689, AC004263, AC006468, AC005562, AC005740, AL035634, AC007536, AC002059, AC005184, AL133453, AC006254, AC004099, AC002378, AC024952, Z85987, AP001728, AC022149, AC006345, AL445490, AC004070, AC009032, AC008687, AC005726, AL033383, AL049570, AC005089, Z84484, AP001718, Z82244, AL157938, AC006111, AL031658, AL365505, AF254822, AB003151, AC003101, AC006455, AC005071, AL390374, AL162430, AC005056, AL035682, AC005666, AC005064, AL139188, AL136124, AF196779, AP001714, AC004813, Z85986, AC003982, AC004832, AC009238, AC006536, AL022320, AC018663, AC005081, AL137139, AL021707, AL031588, AC074338, Z82206, U80017, AL049764, AC004033, AC008050, AC007488, AL445217, AC004967, AL449217, AC011742, AL117694, AL022322, AL035400, AF243527, AC008267, AL022097, AL031311, D87009, AL132639, AL031667, AL137143, AC002080, AC005919, AC006101, AC005261, AC002312, AD001527, AC022336, U62292, AL353706, AL049829, AC018697, AP000252, AJ277546, AL050335, AL136228, AC011442, and AC009097.
HTXBS38	2467	604984	1 - 745	15 - 759	BF222872, AA746604, AA804603, AA293700, AA292944, BE719392, BE258934, R14098, AW408573, AL521473, AC010239, AC010239, AC010239, AC010297, and AC010297.
HTXBU88	2468	604985	1 - 762	15 - 776	AI494203, AV756491, AW270385, AW021917, AW976024, AV757289, AA714110, BE138594, AV695478, AL134330, AI683131, AW237905, AW798093, AW265688, AV764259, AI923052, AL079734, AL037777, AU144320, AI815210, AV758870, AW958962, AA491814, AW272294, AA225406, BG112362, AW574899, R92658, AA569206, AL038842, BF899132, AW069227, BF725884, AA602906, AW023111, AI573198, C75574, AW500250, AI620376, AW105729, AV760014, C75406, AW957502, AA680243, AI859438, AW574794, T57767, AI431513, AI061313, AA626040, BF526964, AI634187, AA622801, AI040051, AI206841, AI246796, AA515728, AI862716, AI887235, BG029528, AW499745, AA643770, AI962030, AA832175, AW471332, AV755593, AW501806, AI732120, AI312309, AI732151, AI635440, AW963463, AI791211, AI821947, AI457397, AV761107, AA302812, D44639,

					<p>AL047349, AW474822, AU143972, AI633168, AI590404, AW103383, AW897556, AI280678, BF891585, AL043878, AW303872, BE049409, AI961983, BF057578, AA525156, AI417469, AI085242, T47138, AA765925, AW512196, AI078409, AW468372, AI380617, AW020612, AI362442, AA804943, AI669589, AL449689, AA700943, BF854090, AA878140, AV735495, AI754037, C75602, AI189682, AW819125, AA664604, AI609992, AI821044, AW190505, AV691556, AV693536, AA501460, AW021399, AA714011, AA314891, AV734149, AI499954, AI687343, AV763550, AW020094, AI457313, AV702109, AV740009, AL118991, AA284247, AV742057, BG180976, AA484892, BF968874, AW008169, AC005052, AC008102, AC005519, Z95152, AL035681, AL050349, AC022308, AC007283, AL133246, AC007055, AL049780, AC008848, AP001716, AC006329, AC005412, AL137162, AL133163, AC022148, AC007510, AC011475, AC007226, AC005815, AL033527, AL157938, AC004671, AC005821, AL035420, AC004825, AC010271, AC020663, AC011495, AL117352, AP001137, AL096701, Z82201, AL035659, AC007664, AC010677, AC007541, AF165926, AL022312, AF196779, AL031668, AL020997, AC005393, AC005932, AC011479, AL109743, AL021579, AC011453, K03021, AF107885, AL133174, AL136362, AC005527, AC020955, AC020904, Z93930, AF195953, AL031664, AC010789, AC011444, AC023490, AC021876, D87675, AC007934, AL121952, M37468, AL357497, AL391839, AC005696, Z84469, AL133467, AC010083, AC004971, AC005038, AL136170, AC006483, AL117258, AC008155, AC004531, AC007277, AC005920, AC006330, AL031003, AL035587, Z68756, AP001681, AC008649, AL160175, AC007687, AL121977, AL034420, AL034372, AL023284, AL109956, AC005800, L78810, AL135844, AL035079, AC007993, U47924, AL132639, AC005529, AL354720, AP000511, AC004151, D84394, AC018663, AL035086, AC004821, AL109935, AC004223, AC008760, AC004643, AC004859, AL109627, AL133153, AL021391, AC002991, Z81364, AL353807, AC006241, AL136223, AC013436, AC005971, AP000495, AL137918, AC005592, AL163248, AL135746, AC008264, AL118496, AL354696, AP000298, AC005000, AC004033, AC004840, AC005921, AL034417, AL109914, AC016995, AC004965, AC006211, AC007242, AL049692, AP000031, Z84488,</p>
--	--	--	--	--	--

					<p>Z97632, AL450224, AC007850, AC009756, AC006449, AP000133, AL356299, AL357150, AC000025, AB023048, AL357752, AC005015, AC008747, AL121712, AC002472, AC018738, AC004955, AL133260, AP000044, AC004031, AL135744, AL121652, Z97055, AC005747, AC020552, AL353194, AP001670, AC005500, AC005969, AC007957, AC008925, Z94056, AC004222, AC002312, AJ295844, AL049761, AC009247, AP000557, AF053356, AL158198, Z85996, AC005037, AC007653, AL137039, AL356794, AL359382, AC004638, AC020550, AC007917, AC009178, AL356057, AC008537, AC006071, Z82976, AL035445, AL117336, AP000501, AC004552, AP000289, AF168787, AC015550, AC005370, AC004605, AL121752, AL096775, AP000042, AP000110, Z97630, AL022396, AC005586, AC005369, AC011547, Z81357, AC006468, AL133453, AL049779, Z99716, AL023553, Z99127, AC005099, AC005694, AC004805, AC010285, AC007878, AC006441, AC005355, AC009721, AC020908, AC004851, AC018719, AF216667, AC007956, AL121890, AL009181, AL136304, AC011445, AP001725, AL356010, AC008569, Y18000, AC011509, AC008694, and AC008694.</p>
HTXCP27	2469	892364	1 - 1559	15 - 1573	<p>AI016791, AL521555, AI769584, AW003495, BE856337, BF446219, AI949680, AI690420, AI674866, AL533716, AW129519, AA708807, AI623188, BE293472, BF035143, AI954672, AL521554, AW514319, BE249868, BF689284, BE538568, AI563921, AW771608, AL517350, BE259763, AW087864, AI432554, AI634705, AL515207, BG036122, BG236428, BG115132, AI432562, AA418416, AI570350, AA203332, AI587529, AL044111, AA938139, AI359192, BG114338, AI686700, N53494, AI802218, AI917417, AW250025, AW675447, AI634891, AI697620, AU151799, AI290844, BF115088, AW163096, AA418531, AU151100, AI619769, BF315350, AU152226, AI963691, AI150541, AI933325, AU160147, AI204201, BG231985, AI983072, AA190579, AI687672, BF341923, AA831968, AI435812, AI335028, AI587523, AW079871, AU160398, AA292839, BE886136, AI885074, BF925747, AI378193, AI991100, AA115707, AL046114, AL516376, AI057142, AI962687, BF434788, AA132539, BE305043, AA126965, AI825350, BE326854, BF313836, AI637949, N36716, BE730329, AW516535, AA460426, AI754481, AI655486, N26077, AI445408, AW370275, BE391119, W38641, AA994318, AA613447, AA133683, BE149668, AA079391, T30419, AI683064, AI207138, AW612812, W38314, AA470639, AA151641, AW136714, AW361020, BE391795,</p>

					AA465129, AA398657, AI968061, BF752099, BG257918, AI018391, AA580007, BF183780, AW806697, BE894829, BG005753, BF477892, AW605433, AA070393, BE896031, AU150496, BF342609, AA488550, C02081, BE019785, AA465723, BF476177, AA621622, AI675741, AI634337, AI909859, AI417892, AI372924, R53936, AA587456, AI909888, AA480806, AI686098, Z39444, BF476014, BF094087, BE275719, BE262361, AA992081, F22497, AI168005, T96676, AI910826, F02918, AI690302, BE410095, AA132645, BF125948, T31266, AI335789, AA127135, AA011661, BG015191, AW003770, BE273827, AA468876, BF792848, AI341290, AI989697, AA102615, T35716, BF758622, AA143396, AA599620, BE877526, AA149561, AI474225, BF310379, AA079390, AW374064, BF309325, BE305168, AI630284, AA411024, AA992038, BF685612, AI566643, R53331, F06647, BE791732, BE258093, T34872, BE298365, AI377530, BE892839, BE764643, BF183439, BF305546, BE315518, AI249805, AW243608, BE261268, BF968173, AL046319, BE397655, BF313972, BF027670, AI804129, AI091009, F06506, AL079753, BF898365, BE764651, BF081866, BF477570, AI524863, BF801833, AI370399, AW062337, AW515691, BE540507, BG169718, AI471880, BE743009, AW806702, AA133682, AI204427, AW025644, BE273369, BF315118, BF971477, BF156009, AL045879, BF448724, AI275224, BF767235, BE795519, AW627480, N72497, BF848631, BF767249, Y13936, AR083379, U42383, AX002424, U81159, AX002422, AC074117, and AC074117.
HTXCU30	2470	839846	1 - 1426	15 - 1440	AA604691, AW452507, AA341372, BG058664, AW953071, AL046457, BF852604, AW276817, BG032943, AI963720, BE047069, AL046409, BF940837, BG249643, AI284640, AV763892, AI350211, AW406162, AW276827, AW193265, BF668217, AI345654, AF063563, AI610159, AV762098, BF677892, AW975425, AV761286, BF475381, AI613280, BF736198, AI431303, AV728928, BE160516, AA720702, AL040382, BF241967, AI133164, AV728425, AV760777, BE389111, AL041690, AA680243, AW303196, AW274349, AV760937, AI471481, AV763354, BE350475, AI754955, AU147104, BF854876, AW974109, BF130107, AI289067, AW301350, AV740801, AI754658, AV725423, AW965008, AL037683, AW977303, AA469451, BF339640, AW969698, AW969694, AI801591, AW961994, AW193432, F36273, BE895987, AI281881, AV702857, AI375710, AI688846, BF851403, AW265385, BF678323, AI341664, BE042649, AL138265, AI334443,

0950083 091201

				<p> AV757607, AI379719, AF330238, AI053672, AI313166, AV682003, AI085719, AV762139, AU145711, AA610491, AV703682, AI732120, AV760466, BE908796, AW148792, AL040921, BF939954, AA470969, BF684828, AW029038, BF724767, AI537506, AI061334, AI962050, AA491814, BG109996, AI298710, BG024252, AV764578, AW338086, AI339850, AL039958, AW162049, AV762050, AI619997, AI929531, BG059568, AI340453, BF942454, AI814735, AF074677, AV760774, AW473163, AI355224, BG236735, BF806176, AI539563, AW238542, AI821271, AV764398, BF681427, AV681599, AV691147, AW467362, AV718479, AU151000, BE677379, AV759382, BF965007, AW630298, AV764490, BE872393, BF311000, AV658688, AV710066, AI270117, AI064864, AW269488, AI669453, AV719316, AI434695, AI679782, AW731867, AW511743, AW021583, AI344844, BE540527, AW302013, BF805094, AW276435, AW513362, AI732186, AV738303, BF217299, AV759204, AW103758, AV719311, AW769399, AL119691, AI017024, AA652764, AW502975, AA581903, AW600804, AW956704, AL079869, AW965237, AA521323, BF337291, AW468028, AV762067, C75026, BE042475, AW474160, AW833862, BE206443, AI674873, AV762009, AA623002, AW238278, AI341548, AV704375, AV763122, AL046205, AI801482, AW517737, AA503258, AW410400, BF841869, AV733468, AV764530, AV719422, AV761631, AI149478, D82290, AU146417, BF679274, AV762959, BF697673, AI623898, AW238583, AI801600, AA598586, AV764307, AV760042, AW169537, AA577906, AI761471, AI358571, AU144500, BF683672, AW574794, AI537955, AA594145, AI499938, AI446464, BF131362, AW419262, AL038785, AI733755, AV733830, AI339440, BG250302, BF813686, BG104686, AI246119, AI005388, AV763988, AV762535, AA177061, AI357288, AI951863, AV725431, AF254822, AC006127, AP000114, AJ003147, AL354720, AP001717, AC020904, AC006312, AC004019, AC005046, AL133500, AF001549, AF091512, AC007298, AC006017, AC018720, AC005808, AL109965, AC008569, AL157372, AC007878, AP000112, AL158040, AL356299, Z83840, AL031432, AC010422, AC000353, AC004638, AL121753, AL033376, AC008892, AC018719, AL023575, AP001716, AC004906, AC001231, AC011450, D87675, AC073593, AC005412, AC025593, AC006271, AC005089, AC004929, AP001759, AP001714, AL353807, AC007201, </p>
--	--	--	--	---

					Z93241, AP001727, AC005839, AC004990, AC002549, AC004686, AC018751, AP001705, AL031311, AC005200, AC010643, AC005245, AC008155, AC004841, AC010072, AC012351, U95742, AC005052, AC004971, AC005535, AL136179, AC011497, AC012597, AL121601, AL022315, AC010789, AL133551, AL137918, AL022163, AF186190, AC004840, AC008044, AP000402, AC022432, AC009516, AL021808, AF123462, AL035089, AC007739, AC004024, AL022476, AC006080, AC007272, AL117333, AC007216, AC008753, AL160313, AC027644, AC007051, AC022493, AF010238, AC009228, AC006958, AC004491, AL049795, AP000090, AC020893, AL096700, AC008068, AC005081, AP001752, AL354696, AL034449, AC006057, AL133399, AC003003, AC007030, AC002404, AC002310, AC011455, AP000501, AL031737, AC005921, AC008760, AC005527, AL161731, AC010553, AC006128, Z98304, AC008372, AC009247, AL031281, AC003085, AC008770, AC004381, AC006038, AC002126, AL135927, AL023882, AC011453, AP001691, AP001695, AC005815, AC002429, AL050308, AF134726, AC009087, AC003029, AP000509, U80017, AC005821, AL359457, AC011480, AL121754, AL080243, AP001731, AC004833, AC005231, AC008764, AP000553, AC016995, AL163853, AC002301, AF165926, AP001671, AL133174, AL118520, AL158167, AC005251, AC018738, AC022515, AC004812, AL035659, AL355392, AL157938, AC002451, AC005799, Z97054, AL354864, AC008925, AC005406, AP000163, AC019227, AL096819, AL049709, AC002091, AP000046, AC008277, AB020859, AC007383, AC005828, AC007227, AL022322, AF196969, AC021016, AL136980, Z84469, U72787, AC002350, AL355497, AC004099, AC007055, AL133153, U91321, Z93015, AC008394, AC011742, AL022328, AL021546, AC007676, AC004650, AC007666, AL031666, AL033383, AL078582, AC008623, AL110292, AP001760, AC004070, AC024082, AC008101, AP000350, AC009305, AL158196, AL356782, AC004531, Z98051, AC007620, AC007882, AL359238, AC005323, AL132994, AC002400, AC007226, AC005772, AP000269, AC023344, AL132987, AC006581, AL138759, AC025264, AL049692, AF111170, AL133260, AC004878, Z83841, AP000049, AL139095, AC010165, AC005682, AC005740, AC006127, AC006127, AC002369, and AC002369.
HTXCV44	2471	740773	1 - 1530	15 - 1544	AI088031, AW023939, AI821670, AA478048, AA478209, BG007575, AI678760, AA708114,

0995008 091201

				<p> BE154945, BE154911, N57913, AA988318, AA513692, AC004858, AC010422, AC006515, AC010412, AF109907, AC005102, AF265340, AC007881, AC009244, AL139339, AC007182, AC006001, AF111168, AC009060, AC009311, AC083863, AL121936, AL158196, AC007114, AC020750, AC008806, AC012076, AF207955, AC022336, AC008747, AC006011, AC008155, AC025264, AL136418, AC002115, AC008753, AC009505, AC011500, AC022150, AC074338, U52111, AP001747, AC004087, AL133260, AC018644, AC008068, AC009086, AC005050, AL121893, AC022424, AL118502, AL354696, AF049895, AC007546, AC003007, AC008372, AC004889, AC008569, AC009247, AL360078, AC005531, AC025594, AC004998, AC004103, AC008134, AL031943, AC007450, AL137228, AC006146, AL049872, AC008950, AC025593, AC004033, AL050348, AL023877, AL096776, AL163246, AL137072, AP001669, AC016602, AC024164, Z99495, AL133163, AF001552, AL022318, AP001697, AC019117, AL023693, AC007193, AC012598, AL050321, AC011510, AC011491, AC009892, AC007381, AC010267, AP001628, AC004832, AL109762, AC020728, Z83826, AL117341, AJ271736, AC003037, AC008651, AF235093, AL356010, AC008040, AC010719, AP001598, U95742, Z97053, AL139826, AC011815, AC006478, AC009516, AC004534, AC008062, AC007679, AL035610, AL133551, AC004965, AR036572, U91328, AC011533, AC004967, AP001694, AL137139, AC004448, AC010203, AL121926, AL031003, AL031767, AL355385, AP001754, AC004925, AC022392, AC007637, AL139377, AL121656, AL163533, AL034420, AL137800, AC005180, AC004491, AC003684, AL096791, AC005015, AC007731, AC011497, AC005500, AC007216, AC002425, AC008736, Z85987, AL355392, AC008403, AL132988, AC012442, AL158830, AL033543, AL138976, AL136362, AC004559, AC004840, AF124523, AL031120, AC007676, AL031681, AC006387, AC005368, AC007345, AC020917, AC004032, AL117381, AC008537, AC005288, AL022476, AC005823, U80017, AC005820, AC006502, L78810, AC007151, AC007954, AP001728, AC004000, AP000555, AC005792, AL137009, AC005722, AC018663, AL139352, AC006512, AC010311, AC010180, AC010168, AC004876, AC004617, AC004605, AC005520, AC068799, AL035419, AC004067, </p>
--	--	--	--	---

					AC007279, AL035659, AC013281, and AC027176.
HTXDJ21	2472	739510	1 - 1321	15 - 1335	AW972830, AA504990, AA489820, AA527647, AA570362, AC061991, AC061991, and AC061991.
HTXDJ75	2473	562780	1 - 1614	15 - 1628	AA225175, AU144320, AV764526, AW576034, AL034405, AC011449, Z84483, AC005000, AC005081, AL109806, AC004089, AB009667, AL050341, AL356299, AC005015, AC010412, AP001711, AP000555, AC005088, AC008403, AF030453, AL050335, AC011490, AC005049, AP000688, AC004858, AC020908, AC020552, Z93017, AF109907, Z99716, AL136979, AL136059, AL139343, AC020916, AC021999, AC002316, AL118501, AC020754, AC011491, AC006117, AL133245, AP000553, AC005520, AC005103, AL096791, AF047825, AC007225, AC002369, AC005409, AL031985, Z93015, AC004821, AP001746, AL136137, AC002375, AL031662, AC005037, AC006038, AC009079, AC008784, AL035071, AC004963, AJ277546, AC010463, AC006088, AC009311, AC011450, AL049795, AC002312, AJ246003, AC023880, AL158830, AC006970, AF196779, AL137162, AC027319, AC005180, AC018751, AC008745, AL136170, AC011500, AC007240, AC009155, AP000509, AP001631, AC003003, AC005921, AL390374, AC005837, AC009516, AJ251973, AP001728, AC003037, AP000692, AC010605, AL023807, AC004895, AL133453, AL121658, AC005736, U47924, AL117348, U07562, AL133551, AC006329, AC004876, AC006480, AL355385, AL158040, AC006111, AC004253, AC008044, AC002288, AL121601, AC005102, AC004898, AL022721, AC005874, AC005670, AF134471, AL109865, AF111167, AC005225, AC007546, U95742, AL162615, AC006028, AC002350, AL096840, AL121653, AC020913, AC007722, AL035684, AL034405, AL034405, and AL034405.
HTXDJ85	2474	590015	1 - 1943	15 - 1957	AA401804, AW880562, AA398288, AW975041, AA657797, AA657405, AA928778, W21329, AW474123, N93524, T86243, N46929, AC009420, AC009420, and AC009420.
HTXDK09	2475	754079	1 - 622	15 - 636	AA806417, AA058515, AA526939, AA526924, AW954118, AW016593, AW082771, AI208040, AW978269, AI024074, AA377401, AI632361, BG011792, AA442742, AI858633, AA531612, AW169720, AW973084, AA877913, AI817451, AA765081, BF311648, AW023592, BF311566, AA996104, AA044239, AI014883, AW771752, AA056753, AA836085, AA737566, AA812884, AA845592,

					AA443740, AA760732, AA620932, AA745072, AI142733, AA989404, AI371736, AA844999, AA620949, AI066536, N95172, AI127959, AA056227, AI284661, AA581252, AA187612, AI928743, AI300651, AI015893, AI422812, AA872495, N21256, N23312, W72874, AA047673, AI079128, AI146658, AA613296, W74335, AA603145, AI500352, AI349303, N80774, AA782772, AA588122, AA938344, AA468649, N58038, AA730107, AW873139, AI862987, AA564172, AA845424, AI267922, T53069, AI565295, AI283681, AI933233, AA760885, AA358391, AA778023, F22831, AI193246, AI187081, AA622185, AI422801, AA865485, AA598695, AA809697, AA027352, AI193923, AA810539, N29349, W42662, BE907800, BE384115, AA528130, AI582026, AA595913, H50881, AA160834, AA835106, BE727885, H20978, H26619, AV729234, AV722752, BE278811, R38340, AI017321, AA282593, H29876, BE277241, BE275267, AV704908, BE797262, BE615457, AI000313, BE791768, T54913, T48132, BE255300, AW960219, BE827415, AA643509, BF686333, BF967031, C18333, BE275998, T81971, BE827402, AI802281, T47662, BG164563, AA887311, AA484061, BE729378, BE258948, AA953140, AV705675, BF246169, BE615182, BF304834, AW300695, BF349644, AI886652, AA120814, BE898477, AW059781, D20514, BG151168, H01955, BE727895, BE388304, BE256781, BF972190, AA074602, AI868281, BE730007, AV721733, BF237736, AW581720, AA835961, AI693331, AA569887, BF206450, BE730760, AA341057, F31834, AP000727, AP000727, AC009438, AC009438, AC009438, AP001931, AP001931, AP001931, AP001082, AP001082, AP001082, AC013381, and AC013381.
HTXDO17	2476	843388	1 - 1306	15 - 1320	W92861, W92862, AA064647, AA064689, AW848901, AW848159, AW848118, AW848162, AW848485, AW848634, AW848890, AW848285, AW848427, AW848283, AV761475, AC005243, AC004895, AC002464, AL445435, AP001692, AL133516, AC005386, AL441887, AC005291, AL139327, AC003006, AC008115, AB017653, AC007676, AL034379, AC008279, AC020913, Z98752, AL133353, and AL353777.
HTXDT72	2477	778083	1 - 1507	15 - 1521	AL519585, BE901976, BE782256, BE887585, BE885885, BE259072, BE881931, BF237969, AA191744, AV695824, BG164068, AW247186, BF438703, BE019530, BF111333, AI056228, AA305318, AA158055, AI348164, AI421266, AW167923, AA453703, AA456674, AA779367, AI143900, BF877870, AW471329, BG110754, BE274433, AI369050, AI332645, BF025941, BE857542, AA826716,

					<p>AI985039, AI768210, AA931707, AI951511, AI369624, AA565685, AL134072, BE838862, AW959326, BF435552, BF091480, AA292101, AI274287, BF763827, AI335905, AA158054, BF677597, AA158464, AA158465, BF528635, AA292102, BG055903, AV703383, AI241064, BF763829, AW248501, BE388432, AA250899, H26302, AA252868, AI890115, H61718, Z39961, AA325486, AA528055, AI688367, AI688366, AI868427, N42491, AA252930, BF303919, AA432122, AA257007, AI828976, R21112, AA431109, R62428, T15511, F03139, BE302734, T55464, AW020080, AA336031, R33428, AW020137, AA354344, AA512878, BG164272, AA299418, AW277079, AI246027, Z43896, BF804269, BF027030, AA250841, AA299419, AA643557, AA864515, AA729926, AA281029, BG256942, T15597, BF125472, BE709207, T34474, R33524, BE389474, H61922, H26461, BE384625, AW247410, AA257046, AI866895, H27801, AA026266, AA026194, F06881, F06092, F02347, AA299461, AA479775, BE389620, BF950586, AA629173, AA345146, T30437, BF095479, AI128565, AA490649, BF763757, AA911731, BF799151, AW881160, AW878088, BE835056, AA620344, AA442340, BF245895, AW295377, AW264879, AI379492, AW367942, AF122819, AB014561, AC013570, AC013570, and AC013570.</p>
HTXDU08	2478	827578	1 - 1089	15 - 1103	<p>BF966668, BG110004, BF752445, BF752447, BG254287, AV651818, AV651537, AI741328, AV650956, AW377577, AV651739, BF752454, BE466129, BE551199, AW966667, BF194918, AI431917, BF114783, AA889457, BF223440, BF063359, AW170381, AI167913, AW975808, AI057135, AA524106, AA722809, AI689751, AI921842, AW450256, AI168214, BF752433, BE350130, AA806429, BE219065, AI143743, AW391182, AA576765, R49044, AI672654, AW675497, AA428269, N95174, AV682632, BE439470, BF477695, R56812, AW377491, BF887427, AW449337, BF062743, H07930, AW118876, AI932465, AI863635, AW391217, AI288119, BF437537, AW303424, AI620980, BF932784, N99509, AI742657, AW340749, BE501735, AI262110, AW006878, T34095, AA811021, AA360767, AW469794, AI203762, W21020, W21148, AI452980, AW819814, AW975855, AA532532, R50418, BG036897, H94454, AA045815, AA884023, AI648358, N93742, AW391212, Z26994, Z38154, AW957881, AA045929, AA721083, AA330766, BF511393, BE010067, R37013, AW391013, AA378981, AA731011, R34521, AA782392, W32806, BE410387, AW957802, R14615,</p>

					R18658, AA854507, N71459, AA307675, AC002094, AC002094, AC002094, and AC002094.
HTXDZ68	2479	824545	1 - 954	15 - 968	AF150415, AA521443, AA765651, AA599108, AV738892, AI804030, AC003111, AC004798, AF161369, AC004798, AC004798, AC004798, AC003111, AC003111, and AC003111.
HTXEN33	2480	589556	1 - 1530	15 - 1544	AI248767, BE246093, AR069468, U20949, AR069487, D78261, U63738, U52682, AC058816, and AC044842.
HTXES13	2481	651367	1 - 663	15 - 677	BF344394, AW959944, BE907670, BE382517, BE313765, BF338479, AI339466, BE743989, BE906305, AI570621, BE222828, AI088762, AI817835, AA434549, AA662133, AA699669, BF725056, AI333967, BF683844, AI669307, AI076748, AI310414, AI310480, AA662096, BG054990, AI333025, BE158972, AI309627, AI308854, AA740871, AI310004, AW444735, H45858, BF940054, BE393365, AA948684, AA931252, BE858296, AA552025, AI376399, AI498530, W48747, AA834581, AI289161, AI378371, N25908, AI885673, AI138476, AI301684, AA844504, AA045902, N40684, BE857636, AA676520, AI370578, AA636048, AA687448, AA948027, AA531076, AA807788, AA644403, AI928410, AI568228, AI927015, AA861046, BE782530, AA584366, AA729907, AI214670, W42852, W42853, AA452718, AA468593, AA005400, F19317, AI359175, AA687504, AA047301, AA328076, AA363383, AI300708, AI913876, AA664856, AA322016, AI185524, AW168584, AA045808, W42908, AA371588, AI244359, BE070822, AA323072, AA327326, AA401721, AA319963, T58761, AI695156, AA369484, AI247975, BF796178, AA737174, W48746, H51522, AA292888, AW577232, BF940538, AA450070, H45817, BG150586, BF526895, N98291, AA005401, T58832, AW390693, BF336794, AA336274, BF837412, AA903394, AI004554, BF845035, AA999871, AA987369, AX017831, and AL356352.
HTXFD86	2482	899517	1 - 1664	15 - 1678	BF975692, BE748819, BF663131, BE748349, AV719646, BE093777, BG260073, BF210820, BF381789, AU157441, AW129766, BE694628, AI342385, AI374623, AI142542, AU141881, AA418160, BG054909, BG057480, W06891, AW242835, AW189544, AA040130, AI253058, AI202968, T27092, AA708710, R20281, BE328237, AI700875, R51987, AA564977, R12643, BF244190, AI336115, Z44733, BF826921, AA417909, AW173471, AA976778, BE963237, AU138042, AI457632, AV655157, N55235, AI571177, AI066692, AL138752, AK023620, AC073598, AC073598, AL138752, and AL138752.
HTXGK12	2483	896690	1 - 1665	15 - 1679	AI832168, BG028154, BE791643, AI440196,

09950083-091201

					<p> AW338360, AI286157, BF434926, AA306132, BE779914, AW294196, AI970497, AI376443, AW953556, AV650233, AA852848, AA303759, BE670807, AI806589, BG114492, N68825, N92038, AV718938, AV718633, AW975618, AW964477, AV720791, D59467, AV720203, AW966053, D80164, D81030, AW978648, AV719468, AW966050, AW966062, AW964488, AV718931, D80038, AW966534, AV719188, D80227, AW949641, AW960465, AW966054, AW959062, AW959597, AV699927, C15076, D80269, AW966022, D80195, AV720731, AW966013, D59610, AW966041, AV718707, AW973474, AW964756, AW975621, AW958992, AW956434, AW965185, AW965197, AW966029, D59275, AW965175, AW966065, AV719783, D59502, AW959570, AV718800, AV720464, AW973334, AW966531, AW978634, AV724520, AW949654, D51799, D58283, C14331, AW959799, AW973541, D59859, AW959628, AW949646, AW949658, D80022, AW965184, D80166, AV718770, AV718489, AW975605, D51423, D59619, AW973485, AW978661, D80210, AW965163, D80391, AW960553, D80240, D80253, AV719822, AV718692, AW973307, D80043, D59787, AV722801, AW966075, C14389, AV719324, AV718440, AV720028, AW965177, AW973488, AV720211, AV718844, AV719557, AW973482, AW959582, AV699447, AW958993, AV723927, AW959136, D59927, AV699550, AW962082, AW949656, AW949642, AW959202, D80378, D80045, D80241, D80212, D80366, D50979, AV700229, D80193, D80196, AW975613, D80188, AW965196, AV718530, D80219, AV700889, AW964737, AV720812, AV723097, AW965158, AW960473, AV720878, D57483, D50995, AA305409, AW753053, AW959469, AW949657, D59889, AW962245, D80024, D51060, AW973330, AW956397, AW949629, AW949653, AW949645, AW949633, AW949632, AW949631, AW949643, AW949618, AW949655, AW960504, AW966059, AW960532, AW966023, AW960454, AV721386, AW966043, C14429, C14014, AW960564, AW973447, D51022, AV699746, AW752082, AW975623, AV701004, AW966030, AW962395, T03269, AW178893, AV699682, AW966032, AW973465, AV720654, AW964532, AV719913, AV720150, AV744690, AW949630, AV742001, AV742667, AV701125, AV701335, AV719628, AV701166, AV742430, AV701149, AV701043, AV701332, AV701017, AV701248, AW177440, AV742048, AV701431, AA305578, AW965176, </p>
--	--	--	--	--	---

					AW973490, AW378532, AW179328, AV720533, AV700159, AW964468, D81026, AV720755, D80522, AA514188, AV719049, AW966380, AV699866, D80251, AV701130, AV701419, AV701154, AV701422, AW964766, D80248, AW960474, AV701357, D50923, AX027925, A84916, AJ132110, AR070327, A62300, A62298, AX033851, AX047063, AX047064, AR018138, AX021518, AR087649, X67155, Y17188, AX035434, A25909, D26022, AB028859, AF058696, AX020191, A67220, D89785, A78862, D34614, AJ302649, AX020190, AR008278, AX047062, D88547, AX028130, X82626, AR077702, AR025207, Y12724, A82595, AR060385, AB002449, A94995, AJ294956, AR008443, AR074545, AB012117, AF260572, I50126, I50132, I50128, I50133, I14842, AR074141, AX015396, AR066482, AR016808, AR016514, AR088705, AJ287395, A85396, AR060138, A45456, AR066488, A26615, AR052274, A44171, AX042372, AR074139, Y09669, A85477, I19525, A43192, A43190, AR038669, A86792, AR066487, AR074136, A30438, X93549, AR054175, Y17187, AR066490, D50010, AR008408, A63261, I18367, AR008277, AR008281, AR091537, AR062872, A70867, AR093385, AR016691, AR016690, U46128, D88507, I79511, A64136, A68321, D13509, AR060133, X68127, AF123263, and AL354983.
HTXGL32	2484	732135	1 - 1411	15 - 1425	AW573194, and AI360355.
HTXJD08	2485	566885	1 - 1224	15 - 1238	AI745124, AC011464, AL136140, AL122057, AL139109, AL352984, AL355112, AL117258, AC005822, AL049829, AL353681, AC007436, AC006600, AC008498, AC011248, AC023275, AC005969, AL022151, AL355101, AL022401, AC003035, AC005913, AL359641, AF049895, AL031289, AL031286, AL356513, AC007225, AC010269, AL035423, AL138752, AL023581, AC008277, AL133516, AP001728, AC024947, AC007537, Y10196, AC009953, AL050305, AL135786, AL035090, AC008268, AC004552, AL163209, AC009411, AL390023, AL109921, AC055740, AL137881, AC008433, AC025442, AC007683, AL358293, AL031388, AC006581, Z84470, AC002519, AC018738, AL161646, AC005250, AC004385, AL139182, AL136531, AC004677, AC004025, AL355352, AC002367, AL096829, AL354833, AF205588, AL049843, AL080239, AC008033, AL033530, Z84719, AL121582, AC026704, AP001696, AC009333, AC005220, AL023574, AL035402, AC007380, AP001685, AL031671, AL117337, AL109653, AP001684, AL049830, AC005386, AC008041, AC002476, AL138743, AC021019, AC002994, AC024082, AL133162, AC005229, AP002528,

					AL355530, AC005971, AL133328, AP001595, AF117829, AC008279, AC006017, AC023344, AC021863, AC022173, AL139023, AL355520, AC004106, AC006965, AL137878, AC044791, AL161415, and AL139404.
HTXJD85	2486	840391	1 - 1103	15 - 1117	BF511613, R16403, H06356, AC005874, AF134471, AC005261, AL137039, AL031588, AL132653, AC008649, AP000689, AB003151, AC005080, Z94056, AF243527, AC016637, AL391259, AC008569, AC001231, AL354720, AL050349, AC002425, AC078797, AC078797, and AC078797.
HTXJE12	2487	862072	1 - 1626	15 - 1640	BF342706, BG035446, AI005482, AA781114, AA522519, BG235909, BF125900, AW665523, AV702780, BE504396, AA527389, AI073948, AW573307, AI338315, AW963008, AW970698, AA973588, AA988131, BF057602, AI354277, BE503992, BE551453, AI906249, BF933161, N64487, AA916538, N92221, AW361948, T23802, H11780, AI796937, AW188867, Z44433, AI678325, AI524764, AI306616, BE817526, AI310310, AI301826, R37401, Z40363, N48912, R13493, T91454, BF983587, R42177, AI300719, T97412, T98990, AA323393, BF568332, AA454136, AA844125, AA872855, N45674, BF125698, T97298, AW118112, AB018302, AL137268, AK025609, AF111169, AF275805, AF111169, and AF111169.
HTXJI59	2488	869905	1 - 2046	15 - 2060	BE614278, BG252077, AV705332, BE905132, BE786765, AI084660, AI809967, BG252003, BG108303, BE906642, BG058745, BE788380, AU152616, BE613556, AU149398, AI571308, W52078, AI581132, AI185833, AU149110, AW069045, AA479518, AU148344, BF344416, BE349927, BE208302, BE301712, AI371737, AA688219, BE906733, BG151536, AW057784, BE302277, AW276839, AA767918, AU145894, BE783512, AA151688, BE466258, AA742225, AU153022, W81272, AI590694, AA742312, BF216173, AA995820, AI829835, AA479516, AA659913, AI888398, AW168209, AW276462, AI720004, AI982668, AI335648, BE389934, AA813688, BE464798, AI343756, T49397, AA579809, AA588563, AW853510, AA157273, T57908, W81235, AW050990, AI813965, N99989, BE350474, BF593643, AA617919, AA295064, AW149090, N71715, AI131468, BE887474, AA636034, N94601, AI582572, AW606299, AA743745, BE829873, AW627366, BE046162, AI081800, AI983946, AA579797, AA987874, AA070837, AA669981, AU157606, AW630395, BF027125, AW960826, AU138424, T50498, R52961, BG057768, AI526061, C00415, AW070868, BE829930, W03414, BF154990, AU125446, AI565917, N67488, AA721064,

09950003 091201

					BG261318, AI091079, AI224467, AW082172, BF970039, AA515074, BE208786, W76594, AI935208, AA976579, AI870359, BE645732, AA308143, AA977004, BE905936, AA070068, T49396, T57991, AA994486, F28267, AA329859, W31120, AA029568, AV703268, N99664, BE829890, AA026207, AI219236, W72898, BE410811, H94596, BE829889, AI735136, AW473136, AA320271, AI080225, AI367525, AA376113, AA303748, AA954228, N89362, R17039, T50633, BE829926, AI289037, R44259, AA157311, AA522995, AI766917, BE312263, AW275160, AA610456, AV654127, AA334886, BE737052, AW364120, AA592909, BE910494, BE094847, BE698237, AI220768, BE826001, AI801589, AI909325, AI909243, AA157312, AA999699, AI909306, AI909315, AI909258, AI909322, AI909324, AI557918, BE829920, W23809, AI000802, BE903271, AA375400, AV698096, AV684976, AI815157, BF855016, AX036785, X68148, Y09846, AL034408, AC003102, Y09847, AF197017, AF196996, AX017817, AR003567, U15784, U46956, AC027440, AC034149, AC034149, AC034149, and AC021890.
HTXJJ92	2489	688062	1 - 809	15 - 823	AW575012, AA484373, AL135377, AW576374, AI538870, AA513188, AL031447, AC007405, AC010271, AC000353, AC011510, AC004985, AB023049, AL135928, AF243527, AC007308, AL035455, AL080243, AC008372, AC004851, AC018738, AC005829, AC005803, Z83845, AC004000, AF001549, AC008736, AL049836, AC007272, AC005215, AC008569, Z83844, AC002544, AC009032, AC002350, Z93015, AC003046, AP000512, AC020893, AL353812, AL109743, AL049636, AC007242, AL031777, AC007688, AC005086, AC008403, AL024498, AL024507, AL109823, AC005082, AL136992, AC005280, AC005291, AF215937, AC021036, AC005907, AL163201, AL049829, AL354696, AP001753, AC016831, AL355392, AC006330, AP001726, AC005015, AC004386, AC007597, AC006441, AC006028, AL096701, AL035405, AL031311, U52111, AC006315, AL136981, AC011495, AP001670, AL109976, AF109907, AC011491, AL049856, AC004858, AL118520, AP001713, AC006480, AC004089, AC005102, AL121653, AC009244, AC004098, Z84466, AC004832, AL035086, AP000697, AL132639, AC006960, AC009502, AC005031, AC007216, AF196779, AP000350, AC007537, Z95152, AC006112, AP001711, AC005103, AC008083, AC007686, AC004253, AL021155, AC004821, AC004887, AF038458, AC002457, AL132713, AJ400877, AL158830,

					Z98200, AL138743, AC008848, AC009228, AL121601, AL121897, AC006372, U95742, AC007993, U52112, AL353898, AC007151, AC005682, AP000155, AC005011, AC006057, AL121928, AC024561, AC005081, AC010489, AL023807, AL023284, Z82215, AL121886, AP001667, AL022320, AP001688, AP001718, AL035587, AC083863, Z93017, AP001729, AL163249, AC020916, AF047825, AL121920, U07562, Z82190, AC008064, AC006316, AL022302, AC007193, AC005901, AC004854, AL450226, AC004703, AC008543, AC000041, AC002314, AC005071, AL391122, AC007620, AC010311, AC002565, AL137039, AC008784, AC020904, AF205588, AC007225, AC002059, AC020750, Z94161, AC010463, AC010422, AL445705, AC005914, AL356299, AC011465, AL133288, AC007546, AL132987, AL031577, AL121585, AC008482, AC011455, AC010150, AC004797, AL022336, and AC006026.
HTXJM94	2490	853413	1 - 924	15 - 938	BG024444, BE896196, AW968455, AV706548, AW275036, BG105730, AW504765, AA906072, BE621482, AA446724, BE814134, AW838478, AA609262, AI123259, AI382929, AA017689, AW439934, Z24971, N24537, N33979, N43850, AA459144, AW953804, T34768, N27708, AV728950, T30082, H16873, AF077205, Z98200, Z98200, and Z98200.
HTXJV54	2491	637729	1 - 1882	15 - 1896	AI765535, BF001346, BF432356, AI701144, AI628601, AI167667, BF725081, AI167568, AI038426, AI636448, N64599, BF592944, AI635509, AI971699, AI359072, N79924, AV726317, BF725082, AV725594, AC010206, AC008119, AC008119, AC008119, and AC010206.
HTXJW06	2492	789201	1 - 1480	15 - 1494	AW952371, BF448738, BF525391, BE503178, AI436256, BE552271, AI972385, AI492252, AV707436, BE551513, BE552281, AW009753, AI183654, AI360373, W56751, AA305172, AI802417, AI590638, AI149096, AI597689, AW088684, BG180464, AI498875, AV702493, AA643746, AA446604, AI217779, AW073167, AW157326, AA468851, AA766884, AW082218, AI083941, AA932568, F29109, AA936458, AA482263, W56510, AA057187, AA804336, N29950, AA482364, T54472, AA989232, AI474387, AA325882, AA852753, BE162155, AW130456, AI470427, AA494415, AW967135, BE162159, AI923140, AW806448, AA552761, AA339266, BF725131, AW810576, AW129327, T54424, BF925621, BG250974, AA836475, T59739, AA977388, AA233545, BF989706, BG029102, BE277201, AW473387, BF091317, AW050678, BF963610,

					AW129132, AI567851, BE728640, BF762046, AW293543, BF445463, BG024609, BE294174, T10594, BE392289, D12039, BE747510, BE410373, AA446733, T10560, AL390135, AB037752, AF110776, AC007779, and AC007779.
HTXKB57	2493	844329	1 - 2822	15 - 2836	BE746332, BE250160, BG250511, BE790537, BG109733, BE794650, BE871407, BE395096, BG111951, AI806629, BE257485, BE795803, BE263975, BE742562, BE866828, BE795502, BE255875, BE783791, BF342278, BE546010, BF314278, BE897195, AW966822, AW958714, BE747754, BE542251, AA553530, BG231832, AA404278, AI940621, BF966416, AW958709, AW043740, AA306203, AI700759, AW068246, AW027960, AW028124, AW084841, AI375516, AI745008, AI940631, AW965024, AW516186, BE019217, AI122741, AA779826, AI378187, AI351744, AI831758, AA844486, AW177716, AI016653, AW439704, AI884816, AW592450, AI610610, AI188921, T86796, AI078369, BF507528, AA401285, AI360991, W86514, AI826299, AI341117, R73315, BF856400, AI299038, N33030, AA007548, AA021613, R72851, AA069819, W24871, N28447, AA007549, R50693, AI924743, BE747477, H58842, N68903, AW068155, T33029, Z43505, AI306715, BF850966, W86697, AI193692, AI193945, R18404, BF826654, T86701, AW512162, AI191487, AI040417, AA620845, H95601, BF966545, AA160461, AI868736, AA069909, R50694, BE091218, R35962, R49476, R42467, BE091232, BE746043, AA337897, H67263, C75020, BE250636, AI203308, N95714, H58790, T30599, R33854, AA021562, W00580, N42091, BF886065, AV739843, BE514668, BE074582, AV740588, BE934261, R33853, BE669899, AV736508, C03610, AI288754, BE831628, AA847535, AA019726, AA742772, AW000794, AW376002, BE258035, AW803305, BF959166, AI915677, AA368831, AW385395, AW577405, C20848, AK025825, AC005726, AF161427, X05196, AB017483, and AB037982.
HTXKH22	2494	562781	1 - 1059	15 - 1073	H06497, AI698573, R37408, AC024570, AC024570, AC024570, AC008899, AC008899, AC008899, AC025441, AC025441, and AC025441.
HTXKH40	2495	600378	1 - 1276	15 - 1290	AL521607, BE729126, BG248593, AL522549, BE744899, BE743976, BF971112, BG249430, BF688906, BG116617, BE410692, AI905440, BF570295, AW964395, BG166435, BF340242, BG108566, AL522001, BF031826, AA142829, AA206373, BE883834, BE546682, AA318302, AI905525, AI189410, AI961339, AW579408, BF968375, AI884509, AA121661, AI678707, BE741195, AI052759,

					AA039438, AI086493, AA903122, AI347904, AI624202, BE207861, BE388318, AI828786, AW571711, AI188539, BE349225, AW662459, AW663672, AI473764, AA922287, AI871802, AI720105, AI185017, BF807239, AW268633, AI302604, AW956985, AI677950, AI376533, AW182583, BE272171, BF675911, AI570190, AL521165, BG025351, AA481905, AI419954, AI696156, AI814113, AW000854, AA159025, BE047488, R99569, R45648, R99920, AW675147, AW193234, AI357664, AA954481, H53115, AI679473, AI554473, AI689084, AI560258, BF878947, AA186671, AW005618, AI432912, AA588482, AI886981, AI590039, AI439924, BF940624, H53221, AI184323, BF222290, AI288336, AI933122, H57919, BE247774, H20524, N74606, AW674526, R99107, BE736037, AA128573, AV652985, N25780, N35307, AA630400, AA285114, AV683356, AA828934, N39758, R99741, H93681, BE122842, AW086188, AA371038, BG109330, BE328332, AI699235, AA380216, AA641233, AA335172, D82554, AV694316, AA044290, BG164142, R99649, AA307924, AA843226, AA994664, AA962068, AA744950, AA029921, N33571, Z25248, AA308080, AA284957, AW087573, AW181903, Z28874, AI628527, Z39000, R99675, AW148298, BE743048, N79259, AA860227, AA946734, AA468405, AI023168, N26315, BG112068, BF753347, AA188290, T83417, AI983876, H93682, AI400866, AA633319, AW630527, AI784577, AA877318, AA927059, AA299967, BE262711, H63332, N36491, AA468445, N44679, AA330143, BE206546, BE206670, AA158660, AA337578, BE074777, H66875, AA337579, R77505, AA204882, AA357049, AA480100, AA829201, AA683569, AI885497, AW130772, AA368717, BF569778, BG015451, BG170896, BF352239, BF908186, AW380841, AI572762, BE857669, AA029920, AA044117, AA128695, BF909062, AI982839, BG059418, AA873793, BE086531, BF734256, AA453577, AA191345, AA039437, AA962504, BE931331, H20615, AL521608, T32419, AA976742, AA143372, T81599, BE742694, AA368324, BF945977, AA233960, AA927402, T56764, AA961698, N92149, AA612675, BE719355, AA781272, AL522002, AL532064, Z42865, BE909669, AI905465, BF743866, R25678, AF151829, AF132942, AJ243649, AJ243650, AJ271671, AC004832, AJ243651, and AC005585.
HTXKK76	2496	858755	1 - 1615	15 - 1629	AI628541, AW022897, AA584482, BF828714, BF748990, AV762220, AV732497, AW961160, AA931974, AI821596, AI475114, AL537797, AV704375, AA173983, AA179163, BE747049, AA261953,

095005660" 09101

					<p> AW059943, AV760760, AL044441, AA478048, AI821670, AI346122, AA478209, AC010328, AL138759, AL034380, Z97054, AC006511, AC008155, AC009244, AL442167, AL163285, Z95331, AL139396, AL163203, AL137220, AC007597, AL163246, AC020717, AP001671, AL031005, AC005261, AC016620, AC004075, AC073539, AF001905, AC004089, AL355838, AC004965, AL445490, AC008114, AL163209, Z83841, AC004166, AP001726, AC011479, AL136303, AL096701, AC018719, AL031229, AC005089, AL049795, AL117381, AC004966, AC020916, AC004913, AF070718, AC007782, AL354696, AP001256, Z98044, AC005399, AC011510, AP001601, AC011484, AL133406, AC006064, AC002565, AC025430, AL109804, AC002115, AC005210, AC005952, AL121586, AL162430, AC005969, AC008474, AL365505, AL121601, AC003070, AC009756, AC003043, AC005071, AC007216, AJ003147, AP001678, AC010677, AP001711, AL445248, AC002467, AL034405, AC008403, L44140, AL356244, AL031293, AL023807, AL035461, AC007051, AL035458, AC007684, AL138743, AC005288, AC005764, AL035404, AC005274, AL022476, AL133387, AL138836, AC008372, Z98043, AC005940, AL133353, AL391114, AL034372, AC011491, AC020904, AC010271, AC020552, AC002470, AL080243, Z83845, AF254983, AC005080, AC004167, AC008812, AX039602, AC007272, AC009123, AC011449, AF001549, AC008738, AL118520, AC006285, AL161670, U52111, Z82244, AC010363, AL138787, AC016601, AC003684, AL117693, AL049911, AC005084, AC005519, AC008392, U95742, AP001753, AC006312, AC025594, AL137230, AL033397, AL132775, AC007620, AC004837, AL035659, AL117382, AC007541, AC006345, Z83822, AP000513, AC011497, AP001727, AC005280, Z95124, AC006483, AL163201, AL163248, Z84466, AC004887, AC006971, Z94044, AC002492, AL033525, AL008726, AC011473, Z85986, AL163283, AC006120, AL135901, AC020913, AC005036, AF165176, AL096840, AC007326, AC008521, AL035422, AC004967, Z93015, AL121574, AC008018, AC007739, AL031311, AL133289, Z97181, AC016395, AL136300, AL160231, AC003006, AC021016, AC018639, AP000046, AC004846, AC000035, AC007366, AC006125, AC005011, AL356652, AP001705, AL035249, AC008745, AC002352, AL022163, AP000688, AL162615, Z83844, AC018663, Z98742, AL136137, </p>
--	--	--	--	--	---

					AL161757, AL133320, AC005086, AL035089, AC007242, AL161731, AL133286, AL121903, AC005907, AC005920, AL035587, AL050349, AC019014, AC087095, AC005822, AL450226, AC004686, AC008892, AC000353, AC087094, AC004841, AP001054, Z99127, U96629, AL117336, AC009060, AC007919, AC006277, AL050318, AC002425, AL159977, AC007842, AC004458, AC004812, AC004000, and AC004150.
HTXKL53	2497	783141	1 - 1596	15 - 1610	AA121863, AW972157, AI005330, AW246779, AI553793, AI076501, AA889963, AI245539, AI298575, BF732476, AA483640, AI123318, AI298397, AI829879, BE645309, AU159951, AA627412, AW250974, BF063304, AI094218, AI160351, AI523173, AW731821, N70566, BG259765, AA721451, AA877594, AW376863, AI032434, AA564388, AA723454, AL530857, BE865465, AA404537, H07869, F13805, BF840014, AI055951, AA122217, AI457636, AL518848, AI474982, AI678622, AW189626, W80766, BE740884, AA706470, AW591598, AW571641, AW514292, AW474453, AA741065, AA480437, AA122216, AW051083, BE793925, AW605095, T08020, AL355392, AF161516, AK023992, AF152097, AF177477, AL355392, AL162505, and AC023015.
HTXKS11	2498	688063	1 - 1931	15 - 1945	H56294, H56092, AW470421, AB048998, AC069204, AC069204, and AC016937.
HTXKS27	2499	877820	1 - 1441	15 - 1455	AW976623, AW960541, BE217989, AA808905, BE218518, BE875190, AI279824, BF445188, BE672631, BE467324, BF110736, BF433873, AU153961, AI912984, AI698016, AA831644, AI656363, AA973823, AV756098, BF939052, AW136350, AI384081, AA776403, AI247530, AI024521, AI348295, AI417567, AW505328, AA243344, AI869547, N66641, BE464856, H52842, N22498, AI825211, AI580661, AW207497, BF196037, AI685865, AI080592, AI272182, AI784098, AI611158, C03505, AW205873, AI288137, AA303634, BE468085, AI049588, AW248483, T67112, AI925755, W25985, AI128707, T67113, T12050, Z45926, BF588536, AW516480, AI620704, AA721452, AA748861, R63515, W01319, AI869722, AI916080, BG150041, T12049, AI263474, C03415, AI446201, F04834, BF091587, Z38932, AI560595, BF738730, AA485075, N98667, BE513505, BF734769, AK023258, AF208848, AL359593, AC068385, AC068385, and AC068385.
HTXLC05	2500	843523	1 - 729	15 - 743	AW779536, AI830607, AW293632, BE254656, AI672798, AW389052, BG058550, AA359942, AW605580, AW882500, AC012251, and AC012251.
HTXLC45	2501	834930	1 - 701	15 - 715	
HTXLT36	2502	843477	1 - 1026	15 - 1040	BF038509, BE907863, AW450318, BF064162,

					AA846547, BF090743, AI743758, AW452088, AI038546, AV721433, AI124936, BF221659, AW961339, W88912, AI038218, AI468531, AI401419, AA312153, D12023, AA361016, AI024136, AI308237, AI763043, AA722378, AW043671, AA304380, AA329556, AI917526, BF507852, AW901911, AA483331, AW969635, AA280698, AI524566, BE898992, C75284, AC006160, AC006160, AC006160, and AC006160.
HTXLY94	2503	844165	1 - 3497	15 - 3511	AA020911, BE788018, AL043774, AA019977, BF056613, W28989, BF436846, BE222901, BF593644, BE745686, BF362452, BG169901, AA722786, BE781921, AI479294, AW953288, AI081914, BF892861, AW340385, BF892856, BF059634, AI693001, AI362051, H15465, AI582303, W92504, AI393893, W28775, AW801662, AA972096, AI350787, AA085086, AA716695, AI246148, AI167703, AA176911, AI272111, AA846714, R44017, W22289, H10893, AA983733, BF892850, AA019947, AI655328, BF771956, AI361104, H11171, AI129209, AW883899, H09577, W27911, AW303345, H15409, AI183495, H06910, AI917751, T77487, R12554, AA630822, AI202926, N35843, AW005038, N27096, T24063, AW451315, AI873102, AI359408, H09010, F13366, AI199126, AW276553, R37456, AA179065, BF362464, T71056, AA350792, AI825414, R19553, AW801443, AW594421, F13261, AW883658, AA335582, F10959, AA644598, AI222148, AI221744, Z44777, AA470413, R19030, BF892105, AI341391, AA350791, AI961948, W28409, F09096, AA906000, T71127, AA315168, AA991920, AW801737, N62428, BF736932, AI933848, N40504, BE706305, BF763743, R38376, AA335806, BG057975, F10863, T10127, W28251, AW974911, T10126, H11083, AI656021, AI687663, AI424575, AL043690, AA659600, Z38415, H10849, F05434, H06869, AW951967, BE939030, AI696592, AW753706, T24459, AW753716, BF846116, BF362403, AK025533, AL158822, AL158822, AL158822, and AL353636.
HTXNV66	2504	840597	1 - 2044	15 - 2058	BE792248, AW954130, BG256142, BG107704, BE387653, BF570074, AA020886, AW408062, BG116672, BE747864, BF968877, AI955209, AA868560, AW403898, AI948454, AI017433, AW170522, W74645, BF196572, AA443135, BE466188, AW504451, AI811668, AW452937, AA631600, W28761, AI818285, AW071094, AA912905, AW205374, AA693936, AI636861, AI758902, AI347305, AW270748, AA033756, BE736097, AW571705, AI090605, W15149, AA886310, AW385052, AA486822, AI707616, AA634914, BF764059, AW235273, AA504537, BE263896,

					AI399736, AW504095, AA033755, AW196362, AW408843, AA454931, AA626291, AI805061, AA070902, AA046736, W89118, AW500041, W28310, BE183532, AI183676, AA934729, BF971691, AA505993, AA679417, T65413, AA663909, AW874641, W80974, AI671510, AI819949, AA007507, AA018598, BE783299, AW503803, W89117, AI925696, W92606, AI014474, AI198564, AA046459, AI818952, AW627397, BF081864, BE245973, BF899608, AI818934, R56755, W80973, AI744463, F09622, F11972, AA018034, AA226465, AI817959, AA780831, AI359075, BG058499, AI972639, AA827181, N78660, AW500794, BE561259, W95231, BF368840, AW515819, AA385420, AA333726, BF204094, BE007859, AI086405, AA504463, AI478470, AW503743, AA609139, AA013289, AI418016, AW948546, D11614, W27923, BF742267, AA807696, AA034986, AW183394, AI620079, BE397066, T65487, W94191, AA878990, BF087353, AI134613, AW505187, W94192, AI969910, AA013028, BE268738, R56907, AA034956, AW505080, BF030539, AI381967, AA630618, C00613, BE269411, BE267529, BE832560, BE177758, AA621088, BE267651, R15547, BE269122, AA725322, BE560221, T92724, AW403709, BF889130, AW797095, BE397067, BE268765, AI342134, W22425, BE295992, AI368615, AA779634, BE673414, D31434, H17565, AA353173, BF593146, R23415, H08407, R34989, AA976522, R19683, AA779863, AW571895, AA378856, AA252777, BF569587, BE766711, BE939445, R33966, T63060, AV749275, AA307157, BE770191, BE392477, BE698798, BF742287, AW059828, BE382928, T58123, AK026350, AB029003, AC002400, AK024619, AF165531, AF233522, AF190863, and AL135924.
HTXOL30	2505	838462	1 - 826	15 - 840	BF883491, BF884916, AA768255, N53352, AA985662, Z82215, AC073898, and AC007620.
HTXOW27	2506	839280	1 - 2373	15 - 2387	U55777, Z78337, BE300034, BF026557, BE563006, AL526718, AL526598, BG029578, BE394636, BE856395, BE396082, AV709362, BE276200, BE296060, BE265498, BE966334, BE736038, BE296312, BE296083, BF982930, BE614635, BF972662, BE791686, BE269523, BF341340, BF345556, BE613770, BE393395, AW205654, BF348051, BE895571, AI762983, BE782600, BF795878, AI762523, BF513996, AI676109, AI859583, AW956261, AW960990, AA843899, AW938743, AW248619, AW081190, AI394519, AI435270, AW402704, AW375353, AW072978, AI339596, BF770167, AI498777, AA713835, BE299628, AI131133, AI039680,

0950083 091201

				AA349398, BE047665, AW295939, AW769539, AI273303, AA825495, BF374364, AA825309, AA399251, AA236925, AI357024, BF842267, AA644225, AI003333, H82888, F27994, R17799, AW029170, BF925721, AA045280, BE244494, R61592, Z78338, R43004, AI365458, T53074, AA854424, BF094318, AI362458, AW393482, AA325016, BE700701, BF338138, BF990571, AA702452, T60176, AA226790, AW370637, R61536, AA631391, AA761720, T60082, Z38952, BF898597, AI040474, BF924499, BG252158, BE243966, T53073, Z42809, AW014935, T30802, R34863, AA364491, BE895477, AI140371, AW362331, AW806658, AA769524, AA227495, AI194026, AW014940, AA046561, BE298541, T24506, AA338532, AW386008, AA236741, AI940467, AW751399, AW387144, AI940442, R34965, AW806488, AA412234, H82887, AA378482, BF855331, BF083714, BF335387, AA569384, AI587121, AI918449, AI949960, AW149876, AI783530, AW008048, BE965121, BE254113, AI872423, AI540674, BG104506, BE896769, AI632851, BF812936, BF750879, BE963838, AI271786, BE966729, AI267725, BE047833, AI582932, AI287233, AI569637, AI961589, AI702073, AW020397, AI890507, AI926800, AI866469, BE620084, BG109221, AI241901, BG253412, AA806534, AV756956, BG178351, BF856052, AI281757, AV764180, AW238688, AI581033, AV714485, AI587209, BF965701, AL515089, BE965022, BF982923, BG180996, AW162118, AI866770, AW827144, AA641818, AW090206, AI802372, BE964617, BF752864, AI433157, AW827227, AI612750, BE964614, BF871314, BE011880, AI623828, BE965724, AI611810, BF812961, BE843239, AI559586, AW130068, AA449768, AW051088, AW057937, AI633125, AL513907, AW079432, AW151136, AI698391, AA502794, AI380176, AW265004, AI915291, AW152182, AI859991, AW087193, AI500061, AI889189, AI521560, AI699056, AI521040, BF812426, AW161156, BF313411, AI801620, AL036634, AW193872, AW024275, AW021717, AI917963, AW834302, W74529, AI432570, BE886827, AW150762, AW827249, AW148478, BG027653, AW088691, AV729632, AI890907, Z21709, AI886753, AB031069, AF149758, AJ132338, AJ132339, AJ236590, U01145, L13297, U72620, I46765, AK025383, I89947, AC005992, A76335, AL122100, AL137529, A08910, X59414, A08909, I48978, U00763, AR038854, AK025084, AL133640, AL389935, U80742, AK026649, AK000310, AL050155, AL137550, AK000418, X72889, AK026613,
--	--	--	--	--

					<p>AF079763, D16301, U95739, A08908, A08916, A08913, AK026927, AL390139, AF217982, AL133606, AJ003118, AF175983, AF285167, Z22828, X82434, AL161953, X76228, AK025857, E05822, AF056191, AB048954, AX040974, AB026128, AF260436, AK024538, AL137275, AB044547, AL080156, AF119899, AF132676, AF118090, AF061836, AF111849, AF090886, AF130077, AL110159, AF113694, L19437, AF119896, AL117649, AF130110, AC068496, AL133560, A77033, A77035, Y10936, Z82022, AJ242859, Y14314, U35846, AF125948, AF091084, X06146, A08912, Y07905, AL050138, I89931, AJ299431, AK000655, AK026630, L24896, AR087170, A18777, AB016226, AL050277, AK026480, AF116610, AB050410, AF192557, AX017991, AF142672, AK025015, AF061573, AL110218, S78214, AF266204, I48979, AF026816, AF314091, A83556, AL137523, AK026744, AB052191, A93350, AX010492, AL133113, AL117435, AL137480, S76508, AL133112, AL137479, AL389939, AK026462, A18788, AR013797, AF087943, AF143723, M27260, S77771, AL110225, AL389982, AL137463, AF116602, AK024594, AF218031, AF183393, AR038969, AL050172, AX019230, AL117460, AF008439, AK026532, AK026542, AK026741, U75932, AL117648, AF116688, S78453, S36676, AF067728, AX042059, Y11254, AL080159, AK000636, AL137271, AK027146, AK026659, AL117457, AL137712, AL137533, AF039138, AF039137, AB032264, AK000690, L04504, AF182215, AL110280, AK024546, Y10080, AL110196, AL122050, AK027082, X70685, AF119871, AK025119, AK025491, AK027160, AL049452, AF130087, A07647, AK025410, AF130082, A45787, AK026434, AF111112, AJ010277, AX019229, AK026506, AF230496, AR011880, AK026865, AF116682, I89934, AK026534, AK026408, AL162002, AK026464, AF113019, AK024992, A65341, AL049283, AL050024, I33392, E02349, AF031147, AR053103, AF116646, AK025312, AL050149, AK000323, AF177401, AF061981, AL080148, U42766, AF106862, AJ005690, U72621, AF116676, Y10823, AK026528, AK026057, AL390154, AF113699, AL137560, AF217987, AK026762, AK000083, AL122098, AF090900, AF262032, AF058921, I09499, AL137488, AL137292, AL162004, E12747, AK026593, X99717, X63410, AP002393, AP002393, and AC015962.</p>
HTXPD86	2507	834772	1 - 2050	15 - 2064	<p>AL521101, AL522518, AL521102, AL522517, BF222844, BE245220, AW959835, BG149650, AI609812, AL524379, BG122179, AL040969, BE387667, BG122113, BE546056, BE388954, AI341394, BE311623, AW007709,</p>

					AI991603, AA312852, AI824857, AA147144, AA737797, AA682285, AW515765, BE245192, AA452519, AA421313, AA040246, AI199145, F25756, AI368839, BF343965, BE264849, AL524378, W90483, AA379756, BG107010, AW819052, AA505052, BE787154, AA421266, AA723336, AI693795, N53821, AW819090, BE264656, AI825728, AW954581, BG024734, R89681, R36282, AW015437, AA261983, AA852677, AA364349, N91752, AA040245, T81937, F26379, BE018777, AA488962, AA490411, AW589761, BE244475, N64739, AA897257, AA147087, AA355856, BE542220, AW800193, BG009929, AA350464, AI569772, T12627, BF740555, BF845272, AA612875, BE062916, H22847, AW410081, BE297783, BE269909, AK000552, AJ011376, AK025810, and AK024484.
HTXPT57	2508	897834	1 - 1113	15 - 1127	AA565918, AW192718, AA719835, AL044340, AL527073, AV719941, AI076616, BE207261, AC005995, AC007956, AC005696, AC011484, AC009032, AL050349, AL022721, AC008474, AC006544, AF111168, AC005288, AC005821, AC008403, AL136137, AC005037, AP000553, AF001550, AC007686, AC005180, AL117382, Z93015, AL031427, AC004686, AL096701, AC005940, AC002302, AC010677, AC005015, AC005318, AL138752, AC004867, AL118520, AL034429, AC005102, Z98200, AC010553, AC008521, AC007969, AL049694, AL356299, AC007850, AL109743, AC005722, AL096840, Z84480, AC004765, AC002400, AF254822, AL034420, AC008569, AC002425, AL009172, AC004477, AL031577, AC005666, Z85996, AC010203, AC004929, AC008738, AC005089, AL160175, AL133263, AC005529, AF134726, Z97985, AC011497, AC011895, AC010170, AL157938, AC011465, AC002390, AL109797, AC004087, AL139353, AL031276, AC004967, Z93023, AC006120, AL132653, AC004644, AL355392, AC005822, AC011500, AL136300, AP001728, U80017, AC004253, AC005231, AP000501, U47924, AC005874, AF134471, AC011811, AC004216, AL121890, AL139286, AC000379, AC002563, AC005005, Z97053, AL031311, AC007536, Z82180, AC004000, AL024507, AC006111, AC007637, AP000115, AL157838, AC006160, AC007114, AC005952, AC009000, AC002301, AL023284, AC005212, AC009087, AC005057, AC004971, AP001694, AC004099, AC007240, AC006057, AC011442, AC007225, AL022320, AC027644, AL031295, AL117336,

					AC008760, AC004656, AL133246, AP000556, AC009079, AC002316, AC011495, AL136162, AC006960, AC009516, AC006050, AC005399, AC004878, AC006512, U52111, AL080243, AL136228, AC004821, AC073517, and AC073517.
HTYSJ88	2509	598718	1 - 758	15 - 772	AA412030, AI208174, AW961493, AA318943, AI732151, AW069227, AL079734, AV762633, AW504485, BF725761, AW474168, AL042373, BF668217, BF725844, AI625604, BF526964, BE138594, BF918640, AI687343, BF868994, AW302909, AV710482, AW500684, AV762354, AW973992, AI300054, AW270385, AA715814, AI267356, AW813668, AI755214, BE062478, AI380617, AI754105, AI754567, AL043878, AI369580, AI267450, AI792575, AI282253, AV760391, AI821714, AI792133, AI791913, AA535216, BF528591, BE062476, AI345681, AI345675, AW327624, AW576251, AI654738, AI081147, AI336054, AV760389, AA487226, AI634187, AI696955, AC006965, AL122015, AL133545, AL049795, AL355392, AL121754, AL139232, D87675, AL354720, AL163208, AP000493, AL009181, AL136295, AC004594, AL035461, D83253, AC010150, AC007065, AC005370, AC000025, AC004884, AC022392, AL035659, AL139378, AC004148, AC005013, AC010282, Z85987, AC006057, AC004259, AC009600, AC004686, AL031276, AL139123, AC004600, AL031847, AP000150, AC008616, AC083871, AC002377, AC006160, AL035091, AP000009, AL354773, AL117351, AC008806, AP001429, AL035086, Z82206, AC008764, AC005189, AC009331, AC005358, AC004973, AL353807, AC018633, AC020663, AF254983, AP000302, AC004067, AC016995, AC010422, Z82215, AL163201, AC006211, AC010200, AC006511, AL163248, AL022165, AF126531, AC011497, AC002349, AC010677, AC004089, AC008569, AC006120, AL157938, AP001443, AC003683, AL035246, AC006130, AL022320, AP000997, AC011811, AF168787, AC005736, AC006561, AF031078, AC008623, AC005340, AC005952, AF030876, AC005103, Y10196, AC004851, AL135927, AC007227, AL353680, AL121897, AF196969, Y14768, AC005015, AL008732, AC009511, AC004707, Z98051, AL158046, AL049709, AL031297, AC018719, AL137059, AL022318, Z69719, AP000088, AC009087, AC009958, AF047825, AC004846, AL078476, AL049776, AL118501, AC004840, AC004876, AL121989, AC008509, AC007383, AL157702, AL031657, AL445258, AC006441, AL354766, AC004477, AL049870, AC003687, AL136124,

					AC005081, AC004971, AJ277546, AP000505, AC007030, AL157791, AC011462, AC004694, AL358293, AL031281, AC009032, AL024509, AC006011, AL133246, Z97196, AC008812, AL031767, AC011526, AP001694, AC004672, AL162430, AL117381, AL391415, AC005255, AL049650, AL139351, AC011480, AL035681, AL034372, AL353194, AC015973, AC011453, AL049696, AC002072, AC009247, AL136992, AL096819, AC007458, AC007878, AC007308, U91319, AL121886, AL109976, AP000558, AC007201, AC005953, AC005998, AC002091, AL136162, AC009086, AL121578, AC008733, AL160231, AC008772, AC005084, AL442064, AC002400, AL049697, AC022148, AP000506, AC004685, AC005145, AC004028, AL109915, AL354836, AC006958, AC009480, AF129756, AC016602, AL135785, AL109804, AC002470, AL031589, AC004887, AC012089, AC005225, AL022399, AC002350, AL031584, AC012309, AL136094, AC004381, AC013429, AL121601, Z80896, AP000112, Z97985, AL138836, AC024561, AL353748, AL136179, AL031432, AC007098, Z83846, AC011362, AC004845, AP001670, AC007683, AL022316, AP000356, AL353658, and AL353658.
HUDBE20	2510	637731	1 - 1000	15 - 1014	AL158089, AL158089, AL158089, and AL158089.
HUDBK47	2511	669688	1 - 1628	15 - 1642	AL117327, AC004859, AC006411, AC006411, AC018946, AC018946, AC027478, and AC027478.
HUFAB57	2512	587275	1 - 1520	15 - 1534	AI075765, BF511556, AI700584, BE047285, BE350093, AI300389, AI350116, AI092464, AI672219, AI810545, AW451849, AI949748, W17246, AI199260, AV746568, AI349327, BE815322, AA528022, H89427, AI284787, BF798855, AA524323, AA994801, AA634039, AW074763, AA923362, AI307448, AI862259, N78843, AI224260, AI349625, AI312256, T29866, AI223617, AI312337, AI309447, AI313341, AI306266, AI284614, AI312336, AI305757, AI609857, AW302009, AI271061, AI335152, AI279036, AI345078, AI223510, AI224767, BE041933, AA650336, AI309381, AW302987, AI271412, AW268753, AW661990, BE041588, T63436, BF885328, AA609425, AI284368, AA732526, BE893005, AV733824, AW811135, BG119916, BG255030, AW079432, AI471909, BF569448, AV704928, AI887775, BF726146, AL033526, AR088270, AL049824, AF042386, AF104013, AL033527, AR068879, I58414, AK024413, AR009513, AR031487, AR001352, AR053029, M97016, AC006059, AF184110, AC010311, and AL353680.
HUFAL17	2513	741072	1 - 843	15 - 857	AA524604, AW169183, BF823372,

					BF823375, BE242879, H70187, AL079682, AL031285, AL080249, AL163201, AC068658, AC005353, AL034405, AL109897, AC018764, AL138743, Z82215, AL133245, AJ277557, AC004922, AC006071, AL357374, AC011480, AC012531, AC004707, AC005516, AL034344, AC074331, AC022816, AL356747, AP001748, AL033527, AC009314, AC005736, AL121952, AC007238, AC011445, AP000210, AP000132, AC005262, AL391259, and Z93244.
HUFAO92	2514	608175	1 - 805	15 - 819	BF508637, AI808815, AI457550, AI911077, AI658931, AI916359, AW009684, BE962972, AW072228, AA579578, AA622141, BE873629, AW973509, AA552628, AA594836, AA551833, AI167645, BE829868, BE829869, BE829871, BE829870, BE829875, BE829863, BE829935, AA576815, BE829887, BE829862, W23220, BE829872, BE829896, AA295027, BE829925, BE829922, BE829884, AF114127, BE869579, AK025424, AB014603, AL137668, AC022165, AC022165, AC010732, and AC010732.
HUFAO94	2515	668267	1 - 725	15 - 739	BF981862, AI148644, BE867924, BF311998, AI141717, BF509112, BE314616, BG110292, BF311948, BF313743, BF312883, BF311119, AI355280, AV722471, AI814247, BF526877, AI633658, BE273931, AI452731, BE740008, BE560662, AI435066, BE259880, AI139019, BE904125, AI857914, AA581889, BE315038, BE730028, BE269698, BG115129, AA617971, BE907044, BF127883, AW026378, BE018876, BF435025, BE878105, AV705691, N36775, AA830266, BE878373, AI375292, AI023389, AI332719, AI432438, BE206561, AA622478, AW964913, AV753433, AI292252, AA977732, AI092111, AI986398, AI302659, AI285818, AI560415, AI379121, AW051470, N35923, BF350372, AI123258, AI075383, AI075385, AI916505, AI369803, AA286687, BE878298, AA063584, AW072015, AW050793, N42201, AI348562, AV722817, AI474997, AA037218, AI241935, AA242888, R67060, AI081160, AW381262, BF058648, BE772928, H29834, T64303, AI376198, H87649, W00490, AW381259, AA513317, AI985043, AI186406, AA082867, AA064615, AI873125, AA910546, R66180, AA832506, AA036653, H19829, N45936, AA252242, BG026118, N72214, AW404601, AA064673, H89959, T64688, H87148, AA121400, AW352312, AW614467, H19859, AA627610, AA320571, AA365156, AW407121, AI186455, AA814718, AA085339, AA760794, BF900713, AW001262, AA283941, AA563841, AA325877, BE265587, AI865765, AA303566, AI090967, AI341845, BF476282, AA922816, BF842746, BG055494, AA581137, BE843392, BE830401, BE830390, BE830400, AA262055,

09950083-091201

					AI971859, AF112202, AI133534, AF252856, AF025506, AF120162, AC010616, AC010616, and AC010616.
HUFAP33	2516	600379	1 - 1523	15 - 1537	AI492875, AI749838, AI796676, AA918144, AI814590, AI923531, AW188476, AA524255, BF513992, AA847476, AI720725, AI748873, AI150879, AI279072, AW190984, AA483519, AA775055, AI284510, AI492104, AW612904, AI141231, AA613554, AI917655, AW662148, AA662104, AW769047, AA565985, T89953, AI473980, AI623640, AA377113, AW769091, AW581798, BE934311, AA295763, BF064110, BF063227, AW974745, AA662070, AA075641, AA075642, BE829414, AW593671, BF760645, AA309786, BE829378, AW612888, AI626076, AW769569, AI796678, AW149876, AW366372, AI690813, AI559752, AW079432, AL121496, AV736475, AV738825, AI336633, AV737685, AV735391, AV743177, AW028416, AV735682, AV736885, AV740451, AV744590, AV735798, AV743713, AV739235, AV745038, AI345567, AI254226, BG029399, AI567582, AI224463, AV692771, AA421957, AL037558, AW161202, AW302816, F28295, BG114724, AW117652, AI288050, AI621341, AV737481, AW576176, AI590043, AI627714, AW245729, BG180605, AV682035, BG113299, AI345143, AV713517, AV710608, AW167229, AV701560, BG180273, AW058233, AV743831, AW827285, AI352326, AW026905, BF676775, AA207067, BE393784, BF680133, AI612885, AW081606, BF814449, AW023338, AI814087, AW021373, AW090206, BE895585, AI888956, BE966699, AI681952, N22276, AI591407, AI434134, AI473451, AI564765, AW089844, AI624529, AI440263, BF038002, AA243996, BG008941, BG180614, AW105455, AI218156, AW162194, T66952, AI272038, AW090128, AV733336, W48671, AW089390, AW083826, AA614517, AI815162, AV682523, BF698359, AL041150, AI473208, AW020397, AI625516, AI472767, AI923061, AL047826, N63128, AI923190, AI244105, AI572717, AI670895, AI345688, AW673679, AL118781, AI889728, BE906419, BF337602, AI247298, AW148841, BE876029, BG251415, AI811785, AI536601, AI826225, AW050725, AI203903, AI933992, AV723064, AI683099, AI348847, AI336634, AW102902, BG110741, AI830259, AI250646, AA587590, AI879064, AI250627, BF792136, BF811808, AW834282, AI537677, AI699020, AW194014, AI802695, AI114703, AW583182, BF828245, AI696626, AV701707, BG029829, AI954422, AI698391, AV702994, AI251205, AI345013, AI633061, AI335363, AL043239, AL044192, BF969494, AI298026,

				AI623941, AI683395, AI499913, AI890907, AI682903, AI868204, AA641818, AA848053, F34309, AW167222, AI431909, AV757528, AL048499, BF341287, AW082532, AI917963, AW827206, AA580663, AL036652, AI564448, AW080279, AI590423, AW900453, AI538980, AI249946, AI889168, AI538885, AI927233, BF792678, AL515403, BF341334, AI432570, AA857847, AI310940, AA987875, AW020419, BE892572, AI932458, AI689702, AI340600, AJ243224, A90822, AF159456, AJ243212, AJ243211, A90827, AF043112, I72544, U32681, AB020851, AB020850, AB020848, AB020849, AF105035, AK027164, X54971, AL050280, AF254119, AF080622, Y17327, I48978, AL110159, U72621, AF113690, AL050277, AL133565, AL389935, AK000603, U86379, AK027146, AR022283, AF132730, D16301, AL137459, AK026603, AK000612, AL390154, AF218005, AF130092, AL137478, A18777, X76228, AK000137, AL117443, AK026857, AR038854, AF100931, AF162782, AF146568, A15345, AX040958, AL357195, M85165, S36676, AF111849, AL080074, AB024524, AK026526, X53587, AL137526, E12579, AK000655, S77771, AF130075, AL137712, AL080137, AL133113, AL117629, AK024538, AK026647, AL050024, I89947, AL050172, AL110296, Y10823, AY004290, AL133558, AK025391, L40363, AF069506, AL050092, AF115410, AF111112, AF119883, AF119894, AL133093, Y13350, AK025549, AK000636, AL133081, AK025084, X52128, U55017, X67688, AB049880, AL359601, AF090900, AK025356, S68736, AL137658, AL137480, AL137665, A57389, AF113677, AR075041, A07588, AB047623, A08913, AK000257, AK000753, AL122093, L13297, AL137479, I48979, I89931, AB048975, A08912, AL133016, U42766, A08910, A08911, M84133, AR087170, A08909, AK026480, I33391, AF116639, D89079, AF107847, AR070212, AK025491, AK026630, AL133557, AF090886, I30339, I30334, AF177401, AF321617, AR034821, AL117432, AF182215, AK026608, AK024622, A08907, A08908, AF119875, AF087943, AF031147, AF044221, AL137705, S76508, AF061943, X72387, I89934, A32826, A30330, A32827, A30331, AK027111, AB016226, AL049324, AL117463, E05822, E02349, AL359583, AF192557, AK025407, AK026613, AK026746, AF094480, AK026494, S69510, AK024944, AK025113, AL122045, AK025484, AL442082, AF004162, AR068466, AL133049, I68732, AL137550, U49908, AK000476, AB049849, A65341, AL049283, AK026597, X60786, X55446,
--	--	--	--	--

					AL080127, AL359623, AF036268, AL389951, AJ010277, AL353935, AF091084, AK026542, AB048994, AF067728, AX042059, AF215669, X99257, AL050208, AF054986, AF166267, AK024974, AF200464, AF119860, AK025798, AF017152, AL137529, AL117457, AF119856, E01963, AK026038, AL110228, U35846, Z48796, E15582, AF260436, AF130059, AF116631, AB047869, AK000718, AK027113, AF114170, AF017437, AF082526, AF175903, AL049464, AK026086, Y08616, E04233, A90844, AR075044, U90884, E12580, AK026797, AL117460, AB044547, AL122098, U96683, AL050116, AR083283, and AL137656.
HUFAU71	2517	638836	1 - 2132	15 - 2146	BF340737, AI765521, BG114935, BF593742, AI497757, BF743683, BG121981, N40717, AI761233, AW467938, N53455, BF000670, AI818496, AI343785, BF476138, AL043306, AA682443, N24761, BF593836, AI147248, AI086795, AA151316, AI768274, T15981, H17331, H17318, AA132787, AA151317, N35901, N72458, BF909329, H16864, AI749561, AA058371, AA719371, AW959402, AW205506, AW083548, AI142035, AI769860, AI784548, N27951, AI800679, AA595210, T95298, H99958, AI127616, AV652948, R56598, AI041733, N21027, H17360, AA678249, AI459902, AA132895, D58461, T95379, R14499, H62754, R59632, R22689, AA301293, BE327641, BF948700, AI267251, AA133062, AI865829, R59573, BG014697, R40829, H29327, H29420, T08304, AA132582, AI254736, AI302433, AI744176, H16755, BF844589, R43395, R56599, AI241825, AA027907, AA058446, AA027842, H17342, AB032773, AB032786, AC012555, and AC012555.
HUFBK95	2518	683039	1 - 1370	15 - 1384	AW452004, AA056314, AA019290, R35058, AA054208, AI365062, and AC017081.
HUFBP77	2519	596839	1 - 1360	15 - 1374	BE739384, AV710021, BF676739, AI458066, BF061560, AW083173, AA780374, AW474167, D62469, AA662183, AA334613, AW974772, AA580883, AA662139, BE867532, BE738837, AI753636, and AW152474.
HUFBV62	2520	742892	1 - 729	15 - 743	AA738437, AW262213, AI291500, AI038166, AI913142, BE351047, AI800898, AA149855, BF198140, AA403160, AA947723, AA760904, AW976794, AA247914, AW085529, AA905798, AA115111, AA983378, R55017, AI139388, AA872872, AA149828, AI379673, AA114978, AI681324, AA249615, and AK024939.
HUFBY96	2521	838624	1 - 722	15 - 736	BE218028, AW055220, AW015575, BF674149, BF056653, AA482033, BE777992, AA058511, AA723887, AI808961, AI288770, AA436822, AW468020, AA805154, AI279159, AA074967, AA872194,

					AW058239, AA287779, AW043971, AI610856, AI887662, AI371079, BF445269, AA488177, AI623836, AI005140, AI127720, AI393524, AI276307, AA771783, AI418169, AI073775, AI128631, AA582253, BF445266, BF195368, AI287359, AA770180, AA644422, AW806015, AW806000, AW806033, AW806031, AW806010, AW006744, AW806048, AW806041, BE090639, AW806037, BF858519, AW806047, AW806042, AW806012, AW806023, BF379048, AW806007, AW806001, AW806024, AI684205, AI445159, AW806091, AW806111, AW806020, AW028973, AI383732, AI478460, AI032355, AW771014, AW806094, BF379054, AI242222, AW806077, AA720014, AW806071, AW806085, H05978, AW953793, AI262086, AI859087, AA771878, H11474, AW806113, AW959550, AA776516, T16286, AW241190, BE934804, T34752, BF431187, AI701319, AW806086, AW806074, AW160363, R65670, AI718448, BF448031, AA054471, AI985890, BE934801, BE672780, BE090443, AI474395, BE090449, AA528439, BE090458, AI985898, R80363, H91578, AA011175, AW866103, AI422514, R70746, AI565890, BE090451, AA759008, BF998712, BE701936, AW519236, BF665991, AA091455, R32961, AW806115, AI284998, AA936164, BE008295, AA442578, AA249530, BE701923, AW866111, BE180126, BE090638, BE710341, BE180311, AW806027, BE180108, AA653374, R70747, BE180124, AW806068, R65669, AA948526, AA010747, BE180428, BE180354, BF083112, BE180427, BG113360, AW806095, D20602, AV713329, AL162085, AX013702, AF156165, AK000299, AX013997, AC008450, AC008450, AC008453, AC008453, AC010309, and AC010309.
HUFCN72	2522	664789	1 - 789	15 - 803	AC016568, AC025439, AC023134, AC023134, AC023134, AC016568, AC016568, AC025439, and AC025439.
HUFEF79	2523	704092	1 - 996	15 - 1010	N52398, H71335, N72804, R59061, AW295411, AW516424, BE464620, AI288309, BE221550, AW327762, AV736989, BG059419, AW022119, AW393469, BE268088, AA644128, AU154118, AU153558, BF445796, AI926467, AI278389, BE048838, AW403199, AA847471, BE208376, AA620434, AA205539, AI625269, AW269874, AA232096, BF350990, AA176375, AU151143, AW080883, AA071316, F20501, AW768374, BE965953, AI698309, AU149997, AW510430, AU151961, AU150857, AU151862, AU154039, AU153659, AU154406, AW078564, AW770439, AU151448, AW168160,

					AI565494, AA224373, AA558419, AA196494, AI637678, BF027009, AI652397, T29566, BE243387, BF129490, BE962363, AW027477, BE561814, BE966554, BE905977, AA577965, BE208009, AA199870, BF129310, BF128082, AF113699, M97856, and M37893.
HUKAD46	2524	604986	1 - 1540	15 - 1554	BF432546, BE886845, AI864293, AA219173, H69004, AA299902, AW001729, AA219484, AW137258, AI671534, AW242472, AL039919, AA649642, AW419201, AV759204, AW886169, AI334443, AW970896, AW886165, BE350475, AI307201, AI830390, BF797630, AW261871, AI349850, AI284640, AI305547, AW020094, AI814735, AW833862, AL119691, AW406447, AW796255, AA493170, AI613459, AW193265, AI431303, AW021399, AA829225, AA513181, AW474160, AI471543, AV764389, AW419262, AA525807, AI571512, BE147093, BG249643, BE047069, AW673241, AI859742, AL037683, AA779599, AI345654, AA502103, BG058664, AA226153, BE044000, AA649705, AI124798, AW511743, AU158130, H88666, AW276827, AV760937, AW276817, AI963720, AI281881, AI350211, AI733523, AV763540, AI962050, BF885741, AW238583, AI053672, AI355224, BE147827, AI744827, AA134367, AI887468, AV719422, AI471481, AI688846, AA601376, AI061296, AI613280, AI281903, AI085719, AI038990, AL137954, AI282832, AW193432, AI061334, AV763418, AW979269, AI624698, AW274182, AW502975, AV763354, BF724372, AA349854, AV762050, AV762139, AW500125, AA482768, BE047693, AL046409, AW513015, AV762781, BF475757, AV728928, AA810370, AW965008, AW192199, AA758934, AV719506, BE327924, AA664535, AA634101, AV759935, AA513850, AI375710, AW131457, AI358501, AI889781, AV762098, AB023061, AF133588, AC000102, AC000029, AC000080, AC004659, AP000692, AC005663, AP001725, AL031281, AC005602, AC016026, AC011450, AC024082, AC011469, AL031228, AC078899, AC007845, AC018642, AC005971, AJ003147, AC004826, AC007226, AC005245, AC009311, AL008582, AC005011, AC007383, AC009086, AC005082, AP001418, AC003007, AL109743, AL133519, AP000289, AC006275, AL121905, AC011484, AL136137, AL133545, AP000042, AP000110, AL049870, AC010326, AL020993, AP000555, U95742, AC004257, AP000008, AC008379, AL121580, AL109935, AL353807, AL049745, AL121983, AC004695, AL445687,

					AC006236, AC005295, AL031295, AL135744, U89337, Z85989, Z83838, AC008687, AL031311, AL049557, AC009247, AC007406, AB023051, AP000347, AC006130, AC007037, AP001727, AL137802, AC008848, AL109936, AP000704, AC009244, AC006329, AC005740, Z93241, U91321, AC005803, AC006483, AL121578, AC005632, AF165926, Z82188, AL050335, AL133448, AC005154, AP001716, AC007676, AL049547, AJ009611, AC006345, AL031680, AC006165, AC002350, AL139120, AL117377, AC011491, AL031670, AC005696, AL035659, AC005210, AF225899, AL135839, AC004134, AC003035, AC007216, AP000512, AL158040, AC004816, AC005585, AC009154, AC004854, AC006928, AC006501, AL033529, Z97054, AC006561, D84394, AC009516, AL050338, AL022476, AC003956, AP000556, AJ295844, AC005736, AP001732, AP000105, AP000037, AP000552, AC005041, Z82184, AL162430, AL161936, AL034380, AL118497, AL136362, AC007316, AC005225, AC011742, Z97985, AL022336, AF243527, AL109984, AL050332, AC005529, AC008443, AP001720, AL133467, U47924, U52112, AC015550, AC011490, AL157915, AC068323, AF111169, AL137139, AC020916, AL359833, AC006120, AL157877, AC011495, AL157938, AC002347, AL022238, AC005377, AC005921, AF279660, AC005670, AC073316, AC008403, AL390768, U62317, AF196779, AL138787, U91323, AL022326, AL021918, L78833, AC002432, AC011442, AL445483, AL109804, AC004019, AC027319, AL121653, AP001040, AC005822, AC005251, AC004383, Z82190, AC026431, AC006511, AC012627, AL020997, AL031985, AC006360, AL136969, AL161422, AC011479, AC009362, U91326, AC006254, AC006581, AC007899, AC002988, AL021391, AC020603, AL138807, AP001714, AL121928, AL138702, AC005913, AL121601, AP000349, AC020728, AL034372, AL050328, AC009756, AC006211, AL023575, AC005020, AL109653, AL050349, AP000501, AL096701, AC007956, AL049748, AL118510, AL008629, AL049872, AL109797, AC006006, AF053356, AL133382, AC006515, AC010489, AC005288, AC026341, AL354816, AC004802, AC004983, AF168787, AC011523, and AC006312.
HUKAI28	2525	843240	1 - 1686	15 - 1700	AW970124, AI927902, BF726581, AA514595, AA024569, AI522300, AW085977, BF515984, BE265766, BF516552, AA024681, AI655128, AI342667, AA909007, AA453980, BF772864, AA453472, BF772010, R19012, R44083, R16327, Z38629, AA299821, R41403, AK026733, AP001021, and AP001021.

HUKAO50	2526	805975	1 - 2044	15 - 2058	AV722338, AI431788, BF435333, AV722940, AI681264, BF514353, AW518687, AA458581, AI040847, AW364986, AA302862, N28394, AA179069, AA837053, AA809755, N94870, AI273305, AV752097, W39636, BE244937, AW297331, BF746111, AA767777, N29522, AI186482, T30167, Z43594, AI520861, AA971245, AI830344, AA258244, AA179320, R25275, AI800261, Z39659, BF436198, BG014470, AI055986, AA973838, BE896547, BG014481, AA351029, D61168, AA096257, AV749923, AA094345, BE172430, AI394005, AA911658, AI150929, AA191277, AI050897, T17427, AA746683, T89350, BF037031, AA291353, AL513907, T89365, AL514627, AL513597, BF726322, AL513803, AL515375, BG180996, AA247994, AV758179, AL514473, AW162071, AV755290, AV655645, AI436456, AI064830, BF248233, AV682521, AV682266, AV682249, BE785905, BE047863, AV706777, AL135661, AL121270, AV682809, AL119049, AI818683, AI687376, AL047042, BF343172, AV682672, AI679724, AW827203, BG260037, AW268253, AL514791, AW071349, BG058208, AI349772, BG179993, AL514919, BE048319, BG107847, BF054789, BE048071, BE880190, AV756770, AV755613, AL515041, AV757158, AI815383, BG250190, AV708119, AV681857, BF724691, AL045500, AI868831, BF795712, BG036846, AV710479, AI349645, BE964812, AL514129, AL036396, BE877769, AL514793, AL120854, BE964700, AV757797, AV681785, BF970162, AV681951, AL046849, AV758806, BF883916, AV729890, AV755614, AV710608, AL514691, AV681630, AI433157, AL515047, AV682252, AV704928, BE964633, BE968552, AV682222, BG105099, BG108147, BF882343, AI969601, AL036146, AV757327, AI433976, AV704350, AL514359, BG033403, AV682772, AL036802, BG027204, AV682476, BG168696, AV756477, BE048163, AI349614, BF793644, AL120736, BG257535, AV682289, AV756342, BF971016, AL514929, AV758217, AV681668, AV723953, AV682330, AI920968, AV755311, AI500553, BF981774, AV727506, BF970446, BE048081, BF970731, AI608667, AV758110, AL121365, AI687728, AW827211, AV758668, AV763915, AV682351, BF968493, BG259801, BE881155, AI538716, AL513553, AI625079, BE777769, AV757943, BF812933, AI500077, BE048026, BF673434, AV682441, BG179633, AV682074, AV756703, AL047763, BE018711, AV733397, AI567351, BF340031, AV711509, AV757455, AI906328, AL514803, AL036759, AL513643, AL514935, BG252929, AV726951, AV733470,
---------	------	--------	----------	-----------	--

09950083-0101

				<p> BF726297, BF969494, BF792099, BF348329, AW999049, AV715462, BF968041, BE965556, BE965095, AI934036, AV723772, BE964486, BE048135, AA613907, AV755581, AL513911, AL513631, BE047859, BG114104, AI568870, BG108324, AV682479, BF036115, AW238730, AW467961, AV681716, AI349598, BF724198, AV755207, AF116639, I48979, Y11587, AL442072, AF116644, AF116631, AF090943, AF130059, AR079032, AL133016, AF116691, AF090900, AF078844, AL050393, AF113013, AF130105, AL133640, AF118064, AF116646, S78214, AL389978, AF130075, AF090934, L31396, L31397, AL117457, AF130082, AF116602, AF113691, AF118070, A93016, AF125949, AL157431, AF090901, AF130104, AX046603, AF138861, AL049938, AL110196, AL442082, AL137527, AL117460, AL050146, AF090903, AF104032, AJ242859, AB048953, AL080060, AL162006, AK026608, AL133606, A08916, AL390167, AL049452, I89947, AF113690, AF113694, AF218014, AL110221, AK000212, AF119878, AL359596, AL359601, AB049758, AF116688, AX019230, AF113676, S68736, AF130092, AX019229, A08913, AL122050, AR059958, AK026865, U42766, AF113689, AF119875, AF111847, AK026741, AF090896, AK025084, I89931, AL050108, AB047615, AL050149, AB048964, AK026045, AB019565, AL133075, AB041801, AK025958, AK025339, AL096744, X84990, AF106862, AL050116, Y16645, AF113677, AL122093, AL162083, AL049314, AL049466, AL133557, AF113019, AF017152, AL133093, AL050277, AF219137, AL080137, I48978, AK026744, AL080124, AK026855, AL359618, AL133565, AF113699, AL389982, AX006092, E03348, AF116649, AL122121, AF119899, AR011880, AL137557, AL133080, AF091084, AL137283, AL133258, AF158248, Y11254, AK000618, X63574, AL137459, AK025772, AF097996, AJ000937, E07361, AK027096, U91329, X82434, AK026784, AF314091, AL122123, AK026583, AF111851, AF017437, AF119909, AK025092, AL359941, AK000137, AK000083, AL049430, AF125948, AF146568, AB048954, AL117394, AK026542, AK026647, E07108, AK026533, AF207829, AF079765, AF242189, AL110225, A65341, AK024538, AF271350, AL050138, AL353940, AK026452, AL049464, AK000652, AR087170, AL133560, AF177401, AK000445, AK026592, AL117583, AB051158, A03736, AL137550, E02349, AK027113, AF119871, AL049300, AF225424, AK026927, AK000614, A08910, AK026086, AF116682, AK025491, AK026504, E05822, AK026353, AB052191, A08912, AB047904, AK026532, AK026534, </p>
--	--	--	--	---

0950083-091001

					U00763, AJ238278, AL117435, AK026959, AB048974, AF130077, AF130099, AL359615, AL049382, AK026642, AL117585, AF130066, AK025391, AF177336, AB052200, AF130110, AC002467, S61953, X70685, AK027204, AK026480, A08909, AK027164, AL050024, AK000432, AK025967, AK026528, AK000323, U72620, AF067728, Z82022, AK025414, AX042059, AC006944, A77033, A77035, AF116610, AF260566, AB049892, AF175983, AF118094, AF130087, and AX026824.
HUKCS86	2527	826468	1 - 1767	15 - 1781	AL523744, AL518418, AL532448, AL518417, AL520420, AL522351, AL523745, AL518139, BE616497, BE616605, BE733368, BE738588, BE799416, BE791749, BG260098, BE732685, BE731572, BE614414, BE730661, BE615071, AL516298, BE273481, BE900141, BE871390, BE615333, BE391259, BE615677, BG250164, BE273477, BE545185, BE868600, BF343630, BE787044, BG121292, BE311771, BE872564, BF314993, BG253624, AL516299, BF343495, BE901752, BE798255, BG177283, BE784359, BG252058, BE785642, AW966070, BE536452, BE790200, BG169715, BE888542, BG255223, BF340374, AL536080, AW245914, BF339152, AW245656, BE784381, AW378155, BF306529, AF121180, BE544651, AW604406, BE613802, AW370146, AL522350, BE547030, BE615584, BE545666, BG260825, AL520419, BE297286, BE785317, BG106890, BE889515, BE298547, AW954718, BE787356, BF307304, BE382543, BE696066, BE546574, AA522530, AW370144, BG168704, AW073128, BE696081, BE299139, BE907496, BF683301, BE785357, AL536079, W29115, BF724212, BE787716, BE544219, BF341544, AA001456, BF307862, BE548627, AA779698, BF684396, AI435833, BF526830, AW378241, AA419606, AW574777, AW574776, AA604253, AI417546, BF589266, AW378176, AI095039, AA576999, BE546410, AW378136, AW748330, AA102640, BE311602, AA016189, AA856578, AA190486, AI937287, AA056329, AA669335, BE543362, AW006110, AW378238, BE545344, AW748329, AA520984, BF343787, AI525555, AA031658, AI498922, AW027280, AA156568, AI884745, AA838222, BE870933, AW378233, AW378139, AA548372, AA053912, AW378240, AI083719, AW378173, BF527505, AA643059, BF807840, AA618602, AA135206, AI224159, AW403561, AA533627, BG236134, AI459590, AI609698, AA411709, AA599035, AI149145, AA016188, AA968939, AW366242, BE793287, BE787846, AA826910, AI280684, AW352433, AI298959,

					H40708, W92466, BF209941, AW370162, W96135, AW366248, AA430505, AA417650, AA973742, AW352434, BE880249, AA428585, F20504, AA021123, AA729826, AW876927, AA568471, BE301108, BE882000, AA017476, BE543190, BF902746, AA668780, H06492, BE390481, AA748351, AW403563, BF149107, BE548653, AW378167, AI301333, N71755, AA429539, AV708418, AI273214, AA100403, AI933378, AI423965, BE737605, BF215342, BE301209, AA923484, AI275808, BF477479, AA568809, AA028068, AA019443, AW169241, AA059113, Z41712, AA953033, BF182663, AW802862, AA021122, AW955038, AW802936, AA670172, AI287863, AA447746, AI274001, T33341, AA430623, AA236317, H06550, AI765626, BG117204, AI765512, T30788, AA227565, H38380, H86543, BE044381, AW378231, AW361385, H40677, Z45102, AA381982, AA256047, AI721035, AA782736, AK000507, AX015530, AX015378, U91679, R40742, R40742, H30564, H83860, H84004, H86072, H86210, H86409, H86569, H86631, H92138, H92139, H92504, AA013378, AA013379, AA017475, AA021310, AA021419, AA028067, AA035064, AA035574, AA054248, AA059112, AA059194, AA059193, AA135319, AA157740, AA227687, AA253218, AA256134, AA514540, and AC044780.
HUKCS86	2528	844245	1 - 1767	15 - 1781	AL523744, AL518418, AL532448, AL518417, AL520420, AL522351, AL523745, AL518139, BE616497, BE616605, BE733368, BE738588, BE799416, BE791749, BG260098, BE732685, BE731572, BE614414, BE730661, BE615071, AL516298, BE273481, BE900141, BE871390, BE615333, BE391259, BE615677, BG250164, BE273477, BE545185, BE868600, BF343630, BE787044, BG121292, BE311771, BE872564, BF314993, BG253624, AL516299, BF343495, BE901752, BE798255, BG177283, BE784359, BG252058, BE785642, AW966070, BE536452, BE790200, BG169715, BE888542, BG255223, BF340374, AL536080, AW245914, BF339152, AW245656, BE784381, AW378155, BF306529, AF121180, BE544651, AW604406, BE613802, AW370146, AL522350, BE547030, BE615584, BE545666, BG260825, AL520419, BE297286, BE785317, BG106890, BE889515, BE298547, AW954718, BE787356, BF307304, BE382543, BE696066, BE546574, AA522530, AW370144, BG168704, AW073128, BE696081, BE299139, BE907496, BF683301, BE785357, AL536079, W29115, BF724212, BE787716, BE544219, BF341544, AA001456, BF307862, BE548627, AA779698, BF684396, AI435833,

					<p>BF526830, AW378241, AA419606, AW574777, AW574776, AA604253, AI417546, BF589266, AW378176, AI095039, AA576999, BE546410, AW378136, AW748330, AA102640, BE311602, AA016189, AA856578, AA190486, AI937287, AA056329, AA669335, BE543362, AW006110, AW378238, BE545344, AW748329, AA520984, BF343787, AI525555, AA031658, AI498922, AW027280, AA156568, AI884745, AA838222, BE870933, AW378233, AW378139, AA548372, AA053912, AW378240, AI083719, AW378173, BF527505, AA643059, BF807840, AA618602, AA135206, AI224159, AW403561, AA533627, BG236134, AI459590, AI609698, AA411709, AA599035, AI149145, AA016188, AA968939, AW366242, BE793287, BE787846, AA826910, AI280684, AW352433, AI298959, H40708, W92466, BF209941, AW370162, W96135, AW366248, AA430505, AA417650, AA973742, AW352434, BE880249, AA428585, F20504, AA021123, AA729826, AW876927, AA568471, BE301108, BE882000, AA017476, BE543190, BF902746, AA668780, H06492, BE390481, AA748351, AW403563, BF149107, BE548653, AW378167, AI301333, N71755, AA429539, AV708418, AI273214, AA100403, AI933378, AI423965, BE737605, BF215342, BE301209, AA923484, AI275808, BF477479, AA568809, AA028068, AA019443, AW169241, AA059113, Z41712, AA953033, BF182663, AW802862, AA021122, AW955038, AW802936, AA670172, AI287863, AA447746, AI274001, T33341, AA430623, AA236317, H06550, AI765626, BG117204, AI765512, T30788, AA227565, H38380, H86543, BE044381, AW378231, AW361385, H40677, Z45102, AA381982, AA256047, AI721035, AA782736, AK000507, AX015530, AX015378, U91679, R40742, R40742, H30564, H83860, H84004, H86072, H86210, H86409, H86569, H86631, H92138, H92139, H92504, AA013378, AA013379, AA017475, AA021310, AA021419, AA028067, AA035064, AA035574, AA054248, AA059112, AA059194, AA059193, AA135319, AA157740, AA227687, AA253218, AA256134, AA514540, AA618500, AA769191, AA887478, AA903993, AA977546, AA019442, AA170840, AA453677, T26327, AA909244, and T27303.</p>
HUKEA22	2529	702029	1 - 561	15 - 575	<p>AI768894, T81961, AA778796, AW445027, AI692721, AW451656, AL039827, BE218913, AA827367, AI672409, BE218606, AW964995, AI240965, BF512195, BF515912, BE179871, AA643565, BE004534, AB037740,</p>

					AF263462, AL365436, AL365436, and AL365436.
HUKEL79	2530	762857	1 - 632	15 - 646	AI200811, AI273480, AA055069, AW016208, AI867694, AI148522, AA534902, BF431437, AA946681, AA789103, AI968204, AW157475, AI308245, AI631357, AI147158, AI701539, BF063204, AI693343, AI766014, BE219048, AI248990, AW072792, AW270739, AA988740, N42555, AI653355, BE350834, AW572753, AI307759, AU145644, AI651782, AI631170, AI634428, AU153787, AU160160, AU160335, BE550299, AA834076, AI699519, AI475617, AI983108, BF592834, AI352525, AI867579, AA954040, AI765943, AI433435, BF001833, AI401138, AA806246, AI654889, AI814278, AW242151, AA454037, AW976100, AL043498, AA748166, AA700237, R51838, AI394292, AU155898, AI962791, AW418623, AW183387, AA100491, AA779116, H56156, BF002709, AA933754, AI632658, AA853481, N71927, AW467931, AI654423, BF057193, AI623190, AW004041, AW181946, AV717799, AW383482, BF476146, AI669537, AI674550, AA595598, BF028267, BF149183, AI630405, AA654813, BF770445, AW383650, AI684887, AI990184, AA055096, U70322, U72395, U72069, AF059614, AC020960, AC020960, AC008515, AC008972, AC008972, AC019104, and AC019104.
HUKEX37	2531	602275	1 - 515	15 - 529	BF326962, AW996879, and AF165140.
HUKFC71	2532	130074 0	1 - 949	15 - 963	BE378712, BF341602, AV707602, BG031712, BE266129, AA609958, BG116981, BE792230, BG035119, AV751438, BF666333, AI188204, BG256134, AA576035, BF116020, BF035496, BE542860, AA488220, N42261, BE407535, AI810755, BE220391, AA740463, AW592483, BE390292, AI735032, AI358863, AA512922, W74161, AA226209, N32817, AI074905, W31772, BF666644, AI352374, AA722666, AI684218, AI265847, AI830218, AW406347, BG149656, AA618058, AI279507, AA910358, BF679049, AI692622, BF057368, AI858044, AA488254, AI189604, AA226545, AI186453, AA504330, AA745478, AW516888, AV707960, BE645301, AI140320, AA078878, AA526293, BE392600, AU144868, BF732233, AA488167, AA521209, AI804219, AI918044, BF446053, BF310592, H40724, H46968, AA609344, AI200596, AW139530, AI373500, AA455452, AI523658, F18818, AI139576, AA908395, AA936118, BF908639, AA303191, BF590871, AI754433, BE646187, AI581489, AI086462, BF909518, AW796612, AI344465, W16486, AA488197, AI344463, AI147647, AI922977, AI348254, AI081276, AA857299, AI304803, AA831686, AI219376, AA335720, AW206542, AI348479, AW796421, AI341936, AI143196, W31201, AW072105,

					H41002, AI080056, BF155936, N90921, AI147651, AI127019, AA715375, H46969, AA911834, AI000842, AI801923, H59214, AI243928, BE675821, BE270754, BE277044, AA155778, AA147783, AW291842, C18565, BF770025, Z20855, AA483925, AA455451, AW302426, AI251169, AI309388, AI254685, AI611893, AI345689, BE122653, Z21081, F25511, BF768943, N92459, W04671, AW274188, AW274326, AW302312, AI252593, Z20733, AA143742, AI610059, BE138468, BF761202, AI337080, AW002979, AA078877, AK021626, Z56310, A75065, A74315, and A74959.
HUKFV37	2533	844644	1 - 1560	15 - 1574	AL527662, AL525578, AL527114, AL530639, AL525527, AL527971, AL530640, BE613157, BE744419, BE730254, BF974904, BG164178, BG260459, BF034551, BE741238, BF976711, BE877307, BG118709, BF981222, BF974176, BG170280, BF982815, BF033023, BE898281, BE264567, AW245879, BG168005, BE735396, BE867863, BE909093, BF219799, BE393924, BE905461, BE908482, BE901548, BE744815, BG170056, BE275013, AW954693, BF124919, BE617394, BE298595, AW575072, BF036070, BE396290, BE746927, BG165516, BE410451, AL527127, BG115529, BE616937, BG106832, BE870316, BE612995, BE548524, BE617852, BE264389, AW935158, BE617360, BE747690, BG116465, BG176553, AI885402, AW410641, BE962359, AW819594, AW575125, AW410640, BE962247, AI991184, BF976039, AW574756, AW170092, BE674969, BF115964, AW574792, AW387107, AI829944, BE676516, AI921923, AW512364, AW008540, AA745265, AI924538, AI074307, AA873160, AA465253, BF439439, AW337968, BF220187, AW387090, AA477599, AW028819, AI829318, BF437001, AA402134, BE891033, BE208227, BF034119, AI139793, AA864636, W95516, BE617598, AA857987, AA402378, AI479060, AA292270, AI005213, AI091416, AI684266, BF037640, BF037848, AW242087, AW512623, AW512911, AI336176, AI362816, BG013080, BF875069, AI075402, AI222764, AA399135, BE264829, AW363174, AA609827, AI972124, AI161220, AI538743, AA443894, BG014395, W79630, AA456798, AW474208, AI090476, N54144, BF744898, AI446095, AI096558, AI086820, AA873436, AI598268, AI609956, AW194155, BF879837, AI151109, AW934814, AI139048, AI027063, AA536177, AA488987, AW473682, AA937584, AI028016, AI805600, AI469710, F37944, AW469206, AI074246, AI187204, AW117881, AA770528, BE049246, AI434504, AI619841, BF876032, AA431854,

102160" 68005650

				AI348212, AA463799, AI304813, AW473135, AI872263, AI241070, AA838514, AA576443, AA832041, AA394128, AI827052, W79486, AA704924, AI355882, AI635832, AA872815, AW583441, AW516942, AA460236, AI613224, BF765177, BF874815, AI368869, AA477555, AI187174, BE617828, AI248658, AW438489, AI285018, W81550, AA724420, BF881820, AA916469, AW401992, AW591674, AI276959, BF880959, AW169451, BF881809, AW572650, F33861, AI346208, AW801679, BF880954, AW517487, H62110, BF881616, AW628189, AI247720, AW403654, BF880935, R72698, AW771733, AI961866, AI266343, AW192153, BE544077, AI365011, AA488766, BF880930, BF744997, AI359194, H20630, AW006966, AA897757, AI351451, AI620538, AA632947, AA975671, F26639, AA555138, AA305218, AI371332, AA468707, AI904061, AI860610, AW105713, AA973012, AA531066, N93383, AW403402, AI537125, N95718, AW583384, AJ002308, AR069079, AJ002313, AF039085, AF151985, AJ002307, AJ002312, AR069086, AR069082, AR069081, AR069080, AR069085, AR069087, AR069084, AR069083, AJ002310, U77594, AX040958, Y11587, AL389935, AL080148, AK024538, AF217991, AF130055, AK026613, I09499, AK000432, AK000212, I89947, AL137557, AF028823, AK026534, AK026626, Z97214, AF188698, AF065135, X61399, AK000074, AK025465, AK025541, AL137529, AR034821, AL137480, AL137665, AX040974, AF225424, AL137556, X80340, AF032666, AL389939, AB047941, AF111851, AK025375, E01614, E13364, AF116631, AK024594, AF113222, AL137574, AF044323, AL110218, AF130056, I48978, AR038854, AK026408, AR013797, AL117416, M92439, X72889, AB016226, AL359622, X66871, AF119899, AF119875, AF087943, I33392, AK000137, AF132676, AF061836, AF116609, AF031147, AK026746, AF111849, AF094480, S69510, AL133619, AL137547, AL359620, AF106657, AF130110, AJ005690, AK026532, AL137627, A08913, AF119843, AF017790, AF102578, A08912, AF271350, AL133637, AL049339, U92068, AB041801, AJ299431, AL117460, AF061981, AF139986, AF169154, A18777, AF118094, AL080159, A08916, AF106697, AK000250, AL389947, AJ003118, AK025484, AL157433, S76508, AK027116, AF047716, A21103, AL122110, A65340, AK026959, AK025209, AK025092, AF132730, S68736, AL133049, A08910, A08909, AK026086, AL110296, Z82022, AK025632, AK025015, AF205861, S83440, AL050170, AL110222, AR038969, A08908, AK026164, AF090900, S77771, AF002985,
--	--	--	--	--

					<p>AR020905, AL050277, I89931, AL049324, AL353940, AL049314, AF262032, AL137550, AB047904, AF180525, AL133072, U35846, AF079765, AL122121, S78214, AF227198, AK026593, AK024570, AF130104, AR087170, AK000652, AF119883, AK000614, AF113019, AK024992, S36676, AF067728, AX042059, A65341, AK026480, U95114, AK024747, AL389983, A77033, A77035, AL359600, AK026630, AL122098, AL122100, AL137711, AB050431, U87620, AF254119, AF061795, AF151685, AF274348, AF260566, AF274347, AF039138, AF039137, AJ010277, AL353956, I32738, A03736, AF057300, AF057299, AF116682, AC023880, AL133067, AL122049, X66862, AL137478, AB047878, A93350, AL162062, AX010492, AL137273, A23630, I89934, AK024588, AF069506, AF119865, X84990, X55446, AL117578, AL117440, AB047609, E02221, X83508, AY004290, AL162083, AL117587, AB048919, X72387, AL080154, AK000636, AK026885, I17544, X92070, AL133075, AL137256, U78525, T47328, T52071, T71104, T77418, R08599, T96113, T96193, T96194, R01124, T26382, R61783, R72627, H20537, H52662, H61205, N23034, W25402, W37484, W52694, W81549, AA074639, AA458500, AA282885, AA465145, AA525442, AA531532, AA587714, AA588105, AA622357, AA635625, AA635797, AA577351, AA687321, AA834496, AA856643, AA878152, AA885559, AA888543, AA931570, AA961789, AI014631, AI081520, N83457, AA092490, AA402983, AA431853, AA897375, AA971776, and D19664.</p>
HUKFY09	2534	784086	1 - 2721	15 - 2735	<p>AW968859, AV729349, BG166696, BG109533, AW167360, AI267409, AL044039, BE771015, BE882592, AA481275, BF724809, AW964621, BF116223, AW957228, BG058493, BE044455, AI920923, BG028972, BF589265, AW968662, AW341018, BE138870, BE717076, AI803017, AI827728, AI568483, AI741238, AW205267, AA446366, BG153438, AA701544, AI096997, AI354796, BE770995, AW967837, AA133347, AI631041, AA129733, AI340006, AW068733, BF941245, AA775828, AI701159, AI743676, AA789137, AI079897, AA937783, BG055388, AI862837, AW068473, AI694192, AI678459, AI189330, AW169957, AL041504, AI813594, AW085200, AA631768, AW236635, AI190188, AA058759, AI393486, AA570016, AA028141, AA491176, AI910943, AA101174, AI298033, W40456, AA782525, AA831766, N78775, AW339110, AW628334, AA766811, AA284394, AA512984, BE328753, AA917419, BF109466, AI969188, AI185111, BE138834, AI249182, AL041503,</p>

					AI272716, AL121396, BF592373, BF592740, AI190317, AI915796, AI374931, BE548455, AA688050, BF063878, AA563721, AI241121, AA825376, AA991848, AA857984, R51682, AA322968, AI470495, R17294, T40513, AA326583, D51829, AA319660, BG055695, H52339, BE247776, AW016123, AA322218, AA322915, R51594, AW103900, AA716669, AA333546, BE246888, N93954, AW888726, AI817750, BF354697, AA287007, T39302, AI694949, AI884382, BF110569, H52242, T71507, AA057393, BF795799, AA913674, AI419551, AI559589, AA865611, AA332395, AA909847, AA339517, AA868436, AA028178, AA806671, AI696783, AA027302, AA101311, AI472353, R41974, AI795880, AW661978, BE974135, Z44341, AA027301, Z40289, AI499254, AA027305, AW605429, AL044855, W07057, BF803712, AA747265, AA636080, D20480, BE247681, AW630200, BE769002, AA089759, AW237240, AA923313, AW952401, AB007930, AL158846, AL158846, and AL158846.
HUNAL39	2535	559448	1 - 1107	15 - 1121	N64385, AW817795, AW835380, AW902139, BE393346, BG030759, AW902138, BG030748, AW844141, AW834580, BF919509, BE245825, AL134565, BF953663, BF931573, AU136138, BE408894, AW816056, AW834551, BF344952, AW834576, BF876571, BF999859, AW502127, AW881669, AI086283, AW834553, AW811905, AW886473, BF954856, AW937815, BE145170, BF964123, AW802786, AW937721, BE145121, AL138046, BE145717, AA505013, AA505002, AW853448, BE148745, AW963588, BE151171, AW834598, BF815705, AA613356, AW834620, AW834550, AL157598, BG104743, AL157534, AA598424, BF767029, AI809468, AA486944, AA296591, AV732399, AW816130, BE158629, BF929360, BF343369, AW890940, BF991141, AI920994, AW896907, BE157410, BF343443, BE143181, AV651504, BG000571, AW834676, BE565164, AW806938, BF968259, AW853657, AW811963, BF841217, AW811904, BF843550, AV750964, BE172768, BE172774, AA736433, BF727068, BE172591, H93766, AW999731, BE620370, AW811815, N23244, BE894112, AA983442, BF852806, BF999851, AW861625, AW861778, BE064249, BE144522, BF853074, BF815710, BG035655, AA838082, BF746496, AV730945, AA018943, AW935976, BF576449, BE083698, AW983779, AA375960, BE267153, BE144399, AV731374, AA129985, BG000577, BF992174, BE179746, BE781027, H77857, BE074268, BF881870,

09950083 091201

				AV709506, BF987379, AW999460, AW834504, BF792143, BF766739, AV654648, AA326048, AW857800, AW176251, BE158632, AW853505, BF107358, AW892932, AA505003, AA324578, BF744137, AW853504, AL042469, BE145719, AW502325, AW853602, AW857767, AA349831, AL042237, T57665, W02666, AA074660, BG112019, BE142616, BF956959, BF914856, BF848183, AW855753, BF949724, AA718911, BE080852, AW176291, BF923272, AW875095, AW853445, AA383975, AI886095, BF528265, BE161499, AA505014, AW983839, AW853356, AW572478, AF074693, W03161, BG012092, AI940253, AW962193, AW855699, AW131775, AV731285, AW867012, BE152201, BE157408, BF340679, AV712055, AV709241, AA775755, BG009277, BF947907, BE064233, BF999853, AW105462, AA077404, AU117048, W03511, T77350, AW813858, BG000833, AW858847, BF829806, AA504312, AI253389, AL044792, BE562885, AA131562, BE062392, AL031313, AC004384, AC007402, AC005246, AL133247, AP000546, AC083860, AF198097, AL390039, AJ271735, AL354750, AC021070, AC009489, AC010176, AC025472, AL139277, AC004470, AC013410, AL355540, AC006984, AL031671, AC004043, AC006288, U73465, AC009404, AC005023, AL157765, AL022575, AL355352, AL353140, AC008473, AL109941, AL359854, AC007965, AL121930, AL049734, AL049875, AC008250, AC002416, AC007094, AC002523, Z84470, AC004385, AP000952, AC008173, AC005951, AL030998, AL357507, AL157360, AP001670, AL121882, AC007966, AC004074, AL121591, AF130351, Z98255, AL136309, AP001606, AC002564, AP002532, AL022726, AC044791, AL391374, AL159980, AC018360, AC004636, AL136970, AL354943, AC006143, AL360272, AL121958, AC002429, AL353667, AC007282, AC016623, AL050401, AC008716, AL033521, AC025470, AC008175, AC004053, AL121871, AL121654, AF238375, AB045357, AC069077, AL110502, AC006313, AL163248, AC006971, AF001905, AL359197, AP000079, AC018452, AF207955, AC008171, AL355612, AP002015, AL357153, AC016635, AC002085, AC004014, AL135780, AC007392, Z98754, AP001699, AP002027, AL135936, AC061958, AP002026, U80459, AL049732, AC004140, AL137016, AL390965, AL353804, AC004936, AC007628, AC079033, AC002038, AL096864, AL050309, AC004704, AL163203, AC010682, AF198099, AL121986, AC004021,
--	--	--	--	--

0950083 09101

					AL110505, AC004822, AC007065, AL008987, AC008180, AC019050, AC009289, AL356513, AL031000, AC016968, AL357141, AP001346, AL035593, U82828, AC002385, AC008739, AL139090, AL049564, AL122057, AC007743, AC003082, AC005994, AP001669, AL161449, AC008463, AL138702, AP001677, AC009332, Z78022, AC025226, AC008506, AC008471, AL031114, AL121841, AC023480, AC026197, AC013357, AC007051, AL353999, AC026770, AL158201, AC007092, AL450169, AC006928, AC010889, AL133480, AL023806, AC008276, AP001666, AC011198, AL157775, AC012405, Z82203, AC004746, AF011889, AP001686, AL163281, AC009312, AL357499, AP001960, AC004673, AC022443, AC008268, AF117829, AL135960, Z93019, AL035258, AC025920, AC007671, AC005609, AC009319, AL136441, AP001684, AL109766, AL161916, AC024162, AC007158, AL390027, AL022577, AC009405, AL139109, AC024947, AL079343, AL109799, AL139093, AC020599, AP001671, AC008162, AC006374, AL365276, Z92547, AC025467, AL158193, AC007631, AC022014, Z83850, Z95328, Z76735, AL049565, AC007486, Z99571, AC018468, AL390237, AL096704, AC005297, AC008178, AL445306, AC026168, AC008572, AC068713, AC007241, AC005341, AC003992, AF280107, AC015978, AF130342, AL160237, AC009411, AC012077, AC020717, AC020698, AL022153, and AL161645.
HUSAO04	2536	877889	1 - 1957	15 - 1971	AL520131, AU123464, AI653357, AA747428, AA622511, BE799465, BE784158, BE747257, AI761558, AW511493, AU143609, BE279116, AW237063, BE881857, BE384474, BE782452, AA625298, AI417969, AI989806, BE783020, AW964830, AA773026, BF966501, BF590565, BE388329, AW081433, BF063660, BF032640, BF939126, BF062975, AA305890, BE536992, AA460246, AU160677, AV751512, AI992092, BF593194, AA626751, AA576313, AA056478, BE243834, AV709838, AA305940, AW206260, AA447204, AI280277, BE468152, BE872682, AA719986, AA903035, AA663785, AA514792, AA069453, AA400835, AI672368, AA357075, AI420309, AI656766, AA393651, AI200199, AA463810, AA742236, BE873633, AA947985, N74113, AI208423, BG029468, AA341406, AI658738, AA056381, AA639420, BE670905, AV739990, N90490, AI861961, AW075450, AW582780, AW606631, BE844234, AW953252, BG166595, AL120719, AW838395, BE378136,

					BG259634, BE243385, AW582447, AW575808, AI630399, BG259550, BG259545, BE785320, AF132939, AK026619, AK024057, AC008114, AC019269, and AC019269.
HUSAO04	2537	870791	1 - 1957	15 - 1971	AL520131, AU123464, AI653357, AA747428, AA622511, BE799465, BE784158, BE747257, AI761558, AW511493, AU143609, BE279116, AW237063, BE881857, BE384474, BE782452, AA625298, AI417969, AI989806, BE783020, AW964830, AA773026, BF966501, BF590565, BE388329, AW081433, BF063660, BF032640, BF939126, BF062975, AA305890, BE536992, AA460246, AU160677, AV751512, AI992092, BF593194, AA626751, AA576313, AA056478, BE243834, AV709838, AA305940, AW206260, AA447204, AI280277, BE468152, BE872682, AA719986, AA903035, AA514792, AA663785, AA069453, AA400835, AI672368, AI420309, AA357075, AI656766, AA393651, AI200199, AA463810, AA742236, BE873633, AA947985, N74113, AI208423, BG029468, AA341406, AI658738, AA056381, AA639420, BE670905, AV739990, N90490, AI861961, AW075450, AW582780, AW606631, BE844234, AW953252, BG166595, AL120719, AW838395, BE378136, BG259634, BE243385, AW582447, AW575808, AI630399, BG259550, BG259545, BE785320, AF132939, AK026619, AK024057, AC008114, AC019269, and AC019269.
HUSCA09	2538	853417	1 - 1972	15 - 1986	BE731434, BE780760, BE796215, BF980381, AW854036, BG164231, AA907782, AA904282, BF530021, BF919011, AV712099, C75403, N63991, BF919389, BF919390, C75526, AA837611, AA247631, BF919194, BE935394, BE716123, AA226442, AW275432, AU157093, AW969824, AA456976, AA468505, AW963444, AA640260, AW439558, AI791659, BE439854, AI253987, AA394271, AA584655, AW272815, AI809818, AA180775, BE739074, AI281881, AW303142, AA535494, AW805539, AI884522, AI802804, BF725504, AW814024, AW162288, AV718585, D86984, AL117378, AC005288, AC005031, AP001207, AC004999, AC000026, AC002059, AC009501, L48038, D87012, U80017, AC007551, AC004785, AC020663, AP000502, AC002488, AL121873, AL121586, AC016554, AL353807, AC003684, AP000105, AP000037, AE000658, AL031666, AF001552, AL023553, AP000496, AB001523, AC008569, AC004983, AP001714, AC005773, AC010150, AC011489, AL078461, AL353777, AC004941, AL020997, AP001725, AC004522,

					AC004655, U85195, AC008267, AC004647, AL096800, AC008068, AC008170, AC010102, AL357560, AF134726, AP001670, AC020916, AP001746, AC027319, AC007488, AC005225, AP001711, AC006530, AP000275, Z93241, AL356575, Z98750, AL355392, AL035450, Z86061, AC004212, AC018639, AC008038, AP001753, AP002028, AC005527, U63630, AC006312, AC016399, AC004993, AL133215, AL161731, Z83822, AC016397, AL355385, Z98047, AL049795, AC007850, AB023050, AL162584, AL356354, AL356299, AC011479, AL008718, AL035587, AC012099, AP000082, AC005839, AL389886, AL157938, AC010092, M84371, AC005254, AP000115, X15723, M26434, AL121601, AC005529, AL135927, AC007227, AC005736, AC011455, AP000426, AL133230, Z86001, AC005911, AC009060, AL445590, AC002470, AL138733, AP000432, AC020904, AL139331, AL109627, Z73359, AL137162, AC004913, AL359272, AC025594, AC008764, AL050349, Z93023, AC010328, AC006120, AC003695, L13176, AL157372, AP000308, AP000355, Z84469, AL034422, Z80897, AC006211, AL133448, AC010289, AL117348, AL022163, AP000964, AL162424, AP001732, AL049776, AL022476, AC011462, AL445483, AC007365, AF111167, AL022328, AC002480, AL132653, AL121652, AP001717, AC009470, AP000048, AC002984, AC004234, AC009516, AL117332, AC025588, AC004185, AC009946, AL117258, AC007956, AC008745, AX039602, AL136450, AL133350, AC018663, AC005531, U82828, AC004836, AL078583, AC002429, AC002351, and M89651.
HUSCJ01	2539	740775	1 - 1326	15 - 1340	BF979101, AV709325, AW469184, BF511108, BE539806, AA150364, AA236207, BF436941, AA150188, BF435328, BF435139, AI417169, AI986415, AI472773, AA812205, AI699342, AW189108, AA256878, AA905150, AI246457, AA808712, AI921676, BF028438, T17368, AA514616, AI263744, AW084726, BF734581, AA255472, AI138535, AA292484, BF911956, H53524, Z39453, D63323, AA909498, AI383278, N33845, AA236206, AA193510, AW970130, AI744083, F13101, R54121, H53525, T76933, AC002357, and AC002357.
HUSGB23	2540	704094	1 - 705	15 - 719	BF350952, BE068804, BE707038, AV690059, AA828834, AI345334, AA169245, AI689198, AA599080, AI355080, AK024996, AC008806, AC000025, AC005527, AC022148, AC005529, AC005057, AC008372, AC004089, AC008403, AP000045, AC006285, AC008392, AC005412, AC007283, AC006160, AC005015,

0950083 091201

				AL354720, AL137039, AC004531, AP001717, Z95115, AL133395, AC010363, AC008736, AP000113, AC007773, AL135927, AC007227, AL136303, AL450224, AL139123, AC011491, AC005911, AP000300, AL133246, AC020754, AC011465, AL009181, AC018642, Z97054, AC011475, AL035420, AL031291, AC008753, AL136137, AC007272, AC018644, AL354864, AC005736, AC009079, AC003688, AC005086, AF205588, AC010582, AC004655, AC003029, AL138743, AP001427, AL158830, AC011742, AC008745, AC011464, AC005881, AL035400, AP000555, AC020552, AL079304, AC005103, AC005839, AF047825, AC006538, AC011489, AL022318, AC083866, AL357497, AP001751, AC008569, AC004966, AL133243, AL078581, AC002558, AL050335, AL356575, AL355392, AP001730, Z85987, AL035684, AC007225, AC003101, AL035413, AC008543, AL136418, AC006236, AL121981, AC008379, AC009408, AL121712, AC005907, U91318, AL132659, AP000691, AC006483, AC008747, AC022150, AC012309, AC020663, AC021016, AP000017, AC005305, AC005071, AC008474, AL135905, AC003006, AC021999, AC008114, AC011514, AC006038, AL031295, AC005484, AC002565, AC010205, AL031311, AL008719, AC004019, AC004491, AC004098, AC005280, AL078645, AC002400, Z69917, AC010200, AL157938, AC003038, U91326, AL035458, AC027319, AL137119, AC004167, AC005052, AC004858, AL022237, AF168787, AL135839, AL022320, AL133153, AC008997, AC004106, AC004905, AC000035, AC005089, AC010150, AP000116, AP001412, AC005972, AL157818, AL132825, AL355305, AC005233, AC005291, AC007000, AL031228, AC005932, AC010412, AC009470, AC005562, AL033529, U91321, AC002312, AC005919, AL031281, AC006312, AC018751, AL135744, AC009516, AC009086, AL138976, AC005971, U47924, AC005500, Z83840, AL109743, AC010618, Z68284, AC005212, AC004893, AC007193, Z83845, AE000658, AL049872, AL356354, AC005585, AC011530, AL158141, AC020550, AC009477, AC005288, AF053356, AC008155, AL136093, AL049713, AC002996, AL158206, AC005081, AC010553, AC007993, AL132653, AL078461, AC020904, AC003982, AC074331, AC005747, AC005300, AC078899, AL121655, AL050321, AL050318, AL158040, AP001598, AC005512, and
--	--	--	--	--

					AL354935.
HUSGJ09	2541	608133	1 - 674	15 - 688	
HUSGQ57	2542	610258	1 - 1926	15 - 1940	BF228454, BF794050, AA886888, BF576050, AI922414, AW196808, BF575711, AW044240, AI076736, BF197013, AA599294, AI954433, AI640323, AW117617, AW137422, BF000029, D63213, AW966475, BF576866, BF435544, AI985086, N41297, AA384564, AA441947, H02121, AV709100, AA370750, H98134, AA195668, BF679630, AI568072, BF156024, AA174162, AW966469, AA370584, BF796544, AL132778, and AL132778.
HUSGY15	2543	759888	1 - 1512	15 - 1526	AV759702, AA311915, AW972047, AA953779, AA521381, BF241426, AW995845, AI806245, AW361598, AI911194, AI580187, AA884217, AI872423, AI859991, AI610362, BG181066, BF725463, AI473536, AI538850, AI866469, AW054964, AI366900, AI802542, AW118518, AI440239, AI923370, AI926790, BF812960, BE786043, AI886181, AI890907, AW148363, AI568138, AW051258, AI611738, AI860003, AW983832, AI241923, BF814527, AL514829, AI570807, AW026882, BG222103, AI632408, AW302973, AW268060, AI635067, AI698391, BF812938, AW983703, BG116926, AW081036, AW983691, AI889189, AI866770, AL119319, AI866820, AW148294, AI619502, AI677796, AI627988, AI648454, AI915291, BF970652, AI433157, AI702073, AA806720, BF812961, AI633125, AI868931, AW152182, AI524671, AL513913, AL514843, AW008353, AI433976, AI288305, AI308035, AW983829, BF795712, BG257535, BE963035, AI521596, AI640704, BG110517, BG260144, AI609409, BF817402, BG029667, AI624293, AI499463, AI559586, AI963846, BF727091, AI554821, AW858243, BG252929, AW151136, AI539771, AI932794, BE897632, AI537677, AI494201, AW051088, AI500659, AI824576, AI815232, AI801325, AL515195, AI500523, BF812438, AI499963, AI270706, BE885490, AI582932, AI923989, AI284517, AI500706, AI491776, AI445237, AI620003, AW151138, AI521560, AI500662, AW172723, AI284509, AI889168, AI440263, AI564719, AI866573, AI633493, AI434256, AL514773, AI270183, AI805769, AI888661, AI284513, AI587606, BG031815, AI888118, AI613038, BF528717, AI436429, AI889147, AI889376, AI371228, AI921248, AI440252, AI670009, AL047422, AI620089, AI866786, AI696570, AI610557, W74529, AI569975, AI634345, AI887499, AI471909, AW086113, AA449768, AI635467, AA641818, AW167021, AI670002, AI620284, BF868489, AI564259, AL513977, BF982923, AI591420, AI687127, AI590227, BF792961, AV743631, AI648509, AI352497, AI651840, AW131294,

0950083-091201

				<p> AW073865, AI673363, BF895953, AW198090, AI956080, AI694157, AI636588, AI537261, AI371251, AL515413, AL042551, AW151714, AI432030, AL039086, AL514799, BF338002, AI559296, AW238688, AI536638, AL514887, AL045500, AI889133, AI452560, AI254731, BE784387, AI635016, AI500061, AI537837, AI686817, AI890833, AL514757, BE544111, AI683173, BF987104, BG166490, AW079409, AA788861, AI638798, AI569309, AA225339, AI933589, BE886728, BF727072, BF726237, AW149925, N33175, AW150794, AI659795, AI796743, AW079572, AI445829, BF814453, AI828583, R32821, AI860897, AW166583, BG178669, AV709679, AW161156, AI355827, AI445965, BF965814, AI499890, AI349967, AL514359, AI784252, AW190194, AF233322, AX040974, I89947, AR038854, I48978, AK026408, AF087943, A77033, A77035, AF026816, AF116688, AL353940, AX040958, AL080159, I48979, AL137533, AL389939, Z82022, A08913, AL137550, AL389982, AK026462, AL137480, AL110280, A08916, AX026824, AX026823, A08910, A58524, A58523, AL049382, A08909, AK026744, AF061981, AF113019, AL353956, AL137488, U35846, Y14314, AK024538, AF113677, AL080074, AF183393, AF113691, AL137271, AF130055, AL137478, AL137560, AF225424, AF185576, A08912, AL359941, AL050024, I89931, AL137538, A21103, AR087170, AF111849, AK027164, U96683, AF162270, AL117435, X82434, AL122049, A65341, AL162062, AF056191, AL050366, AL359583, AF262032, AF271350, AF090934, AF119899, AL122050, AK026506, AK025798, AF026124, AK026593, AL080148, A03736, AL137292, AF008439, AF090903, AF116682, AK026534, AF285167, AB052191, AK027096, AL162003, AK025632, AL023657, AF177401, AK026532, U80742, E02221, AF314091, AF139986, AF119875, AL390154, AF116654, A93350, AL133075, AF130077, AX010492, AL117440, AL050138, X72889, AF230496, AF116691, AK000418, AL049938, AK026528, AB048953, AJ299431, AK026959, AF217987, AK025435, AL050149, AL122118, AL133560, AL133072, AF032666, AL122110, AK024992, AK026480, AL133080, AK026642, AL133665, AK025414, AX045627, AK027204, AB048975, AL117460, AL110221, AK026927, AL133077, AK025484, AJ012755, Y07905, AL137476, E07361, AF113689, AL137557, AK025084, L30117, AK027146, AF116646, AF130099, AL050277, AL137463, AK000652, AK024588, S36676, AF067728, AX042059, AJ000937, I33392, AK025339, AB047904, AK026542, AK026885, AK027160, </p>
--	--	--	--	---

					AK025312, AK026551, AL133067, AF017437, AF116676, L19437, Y16645, AB051158, AB041801, AL133640, AL137529, AB050431, AL050116, A45787, AL157482, I00734, AF116631, AF106862, AL049283, Y11254, AB050410, AB048919, AF130100, E02349, AL137459, AB048954, AK026762, AB048974, AF130105, AK026630, AF116602, AF130082, E00617, E00717, E00778, E12747, X65873, AF155148, AL389935, AF130104, AF067790, AF119337, AK027114, AL137526, AF118094, AL049452, AL049430, AF113699, AF111851, AL117583, AK026741, AK000083, AK026784, AF153205, AL117585, Y09972, AK027116, AK025524, AF090901, AL133113, AF130110, AL050393, AL122123, U49908, E03348, I03321, Y11587, U67958, AB050533, AF090900, AF061795, AF151685, AL133016, AL050092, AF104032, AF119865, AK026045, AB050534, AF177336, U58996, AF119909, A08908, AL117432, AK026855, AL162002, AC011953, AC027300, AC025036, and AC026985.
HUSHD41	2544	866498	1 - 2562	15 - 2576	BF975230, BF683936, BF972280, BG177292, AW571669, AW276243, AW452044, AW450817, AW080419, AI523987, AI357545, AI223336, AW953367, BG056723, AI421911, BF507478, AI364803, AA354094, AI937821, AI699989, AI624920, H45305, AA251791, D19672, AA922638, AA444017, AA443995, AI655588, AA928313, AA339800, R74251, H45245, AI433894, BE293485, AI023763, AI040104, AI949593, AB046039, AP000763, AP000763, and AP000763.
HUSHK65	2545	847188	1 - 6691	15 - 6705	AU118183, AU118665, BF115928, AU132493, BE879879, AI741712, BF116220, BF590208, BF206819, AW194977, BF058123, AA775455, AI819729, BF112082, BE856768, BE856756, AA127691, AI760642, AI419870, AA099129, BE383584, AW297737, BE218878, N47688, AW473131, AA057388, AA683489, AI831261, AW881091, AI191464, AA931763, BE549811, AA702145, AW298292, BF111985, AA047104, AW008471, AI478485, AI143087, AA034447, N90777, AI797608, W72221, BE644689, AI129223, AI150714, AI934906, N54402, H99253, AW085755, AW029384, BF059616, AW880832, AA057640, AI241539, AI394589, N29046, N28005, AI139365, N21345, N46649, AI363151, AI761556, N79568, AI818035, W19201, N20176, AI480190, AA703696, N34554, BF918682, W92085, AA234942, H29681, AI023775, BE465754, BE218527, AI984281, AI168358, N20668, BE746232, AA862846, AA644365, H98775, AW207126, AI272809, N26125, W77965, AA418398, N31777, BF088496, AI090568, AI752094, N33546, N53300, AA047241,

					AA057680, N31684, AA418495, AI283701, N24184, AW886108, AI360042, BE833906, AW369652, AI277679, N29077, N30533, AI492161, N26156, R99592, W16605, AI673532, R19850, BF365464, N20808, AI457533, BE674312, AI764961, AA156964, R61144, AI124041, R99697, H97683, N31512, AA486096, R98621, N23497, AI091105, AI130881, AI752095, R99444, AI401234, AI970658, AA099166, AI564816, AW612814, W92118, BE833933, BF919960, R23539, AW297512, AW297408, BF763331, BF431063, AW880947, AA724297, AW880829, AA599995, H52434, AI673230, AA127790, BE833936, BE833932, AI092811, H89784, H89687, AA463360, W16999, H51139, AA081096, AI919601, AA857005, AW884885, AW880836, BE833935, AI434215, BG054626, AI351435, BE833943, W45267, H72110, AW339798, N24408, AA295257, AA377802, AA486159, AA298424, AW297821, H29595, AI282185, AI147043, BF087751, N36627, AW137808, AI274033, AW298745, BG231793, T74891, AW449732, BE833930, N25760, BF115581, R67834, N48154, AI207144, AI301764, AA628558, BG036067, AI971868, T96296, AW294313, AA373689, AI333804, AW886714, AA976518, R44289, T35750, AI276217, AW900171, N41524, R61862, H99620, N34920, BF754145, BF732653, R67833, N26379, H01050, Z42070, T96297, AA046600, AA055473, T99900, AA974775, AA081254, AW881860, AW351434, BE504942, AA621145, AA360837, AI696272, AI707913, BF086741, AI718749, AI127943, BE670072, AI474260, AA258408, BF436683, AA009871, AW294130, R36523, AW297619, AW296186, R98622, BF940509, AI919413, BF511675, AA361970, AC007561, AF016098, AF022860, AF022859, AF280544, AF280545, AF280546, AF022855, AF022856, AF022861, AF022854, AF022857, AF022858, AF016297, AC007362, AF281074, R37519, H01803, H72212, H89721, N25995, N44559, N47687, N50190, AA046728, AA126383, AA255756, AA627411, AA828181, AA093313, AA234943, AA463866, AA479309, AA479410, AA703467, T16994, Z38353, and Z42090.
HUSIK45	2546	853932	1 - 1401	15 - 1415	BG120786, BE874219, BF382244, BE566989, BF106051, BF540740, AI955244, BF431256, BF029862, BF130237, BF031915, AI031927, AW955664, BF770314, AI818579, AI086765, BF239706, AI298226, BF382551, BF109411, AA714566, AA829068, BF209498, AA894447, AL047103, AA826215, AA935581, AA856833, N22620, AA330112, AA554879, AI587354, AV659467, AI378756, AW853805, D62365, D62126, D62182,

					D62092, D63132, D79479, D62337, D61725, BF881894, AW614498, D62238, AA330352, D62300, BF229176, D79853, AA962320, AA296363, D62304, D79464, AW386014, BE326435, BF229164, AA564864, BF745392, BF743919, BE005532, BE812564, BF745826, BE392628, AW194502, BF813751, AC004010, AC073309, AC004010, AC004010, and AC004010.
HUSIO57	2547	886206	1 - 911	15 - 925	AI859297, BE383156, BE732153, BE384854, BE736656, AI819813, BE384900, AW301122, AA625654, AA610590, AI601169, AI521678, AA770693, BF970430, AI148402, BG166253, AI797531, AA043176, W56364, AA399245, AI224414, AA412225, AA760787, F25596, AA044224, H49207, H49393, AA044343, AA483644, F35724, W56338, H41195, R83432, H46471, H45052, H46298, T71570, R45220, H46110, H47011, H41194, AA810028, H46033, H50393, R00715, AA593682, H50355, Z40409, AA043175, H45090, BF802334, Z44510, R24958, AF151901, AC037465, AC037465, AC037465, AC018785, and AC018785.
HUSIP17	2548	608647	1 - 685	15 - 699	AI289386, BG259587, AI431323, BG122277, BG111590, BG167830, AI798359, AI924051, AL514791, AI500523, AI538850, AI887775, AI582932, AI284517, AI923989, AI872423, AI590043, AI500706, AI445237, AI491776, AI289791, AI926593, AW151138, BF811804, AI521560, AI889189, AI285417, AW151974, AI500662, AI623302, AI284509, AI539800, AW172723, AI582912, AI538885, AI440263, AI889168, AI927233, AI866573, AW058275, AI633493, AI434256, AI866469, AI805769, AI434242, AI888661, AI500714, AI284513, AI888118, AI285439, AI436429, AI859991, AI889147, AI355779, AI623736, AW194509, AI581033, AI371228, AI431307, AI491710, AI440252, AI872315, AI440238, AI697243, AI567971, AI866786, AI610557, AI860003, AI431316, AI242736, AI828574, AI539260, AI887499, AW151979, AI431238, AI539781, AI889191, AI702065, AI539707, AI885949, AI285419, AW089557, AI559957, AI521571, AI469775, AI866581, AI567953, AI815150, AW074057, AI446495, AI867068, AI952433, AI866691, AI801325, AI225248, AI358271, BG029128, AI932620, AI698352, AI282249, AI371229, AI371237, BE885490, AI815232, AI866458, BG166490, AL047398, AI890472, BE883591, AI434222, AI624681, AI554821, AI866465, AL514209, AI828585, AI434229, AL513647, BE779152, AL047422, BF796402, AI476694, AI815239, AI500659, BF726204, BE622452, AI804505, BF812963, BG116926, AL514833, BF337604, BF345716, AI702062, BG027368, BF530630, AI494201, AI499535, BG261116, AI537677, AL048951, AI804524, AL513983, AI440260, AI432644, BE897632,

					AW151132, AI500683, AI539771, BG035653, BG180788, BE904746, AW151136, AI924024, BG252929, AI289747, AI561170, AI888022, AI440236, AI469764, AI690946, AI648567, AI433157, AI499478, AL043196, BG120398, AI590024, AI687588, BE618455, BG029667, AI362495, AI801286, AI798571, AI539774, AI682915, BG113493, BG110517, AI371243, AI538615, BF812438, AI431315, AI285432, BG257535, AI687607, AI567947, AI493559, AL039390, BF795712, AI274759, AW858522, AX040974, AX040958, AL049423, AX041002, AL133655, U30290, E13998, AL133084, AL133070, AL133051, AL133076, AC023885, and AC023885.
HUSIR70	2549	838163	1 - 1222	15 - 1236	AL526850, AL526820, BE795841, BF792949, BF036122, BE880934, BG251998, BE893579, AV712345, BE906587, BF689193, BE256963, AL526060, BE881392, BE964987, BE385913, AW960397, BE747698, BG236341, AI032386, AI125617, AI754666, BF343485, AA179549, BF344744, BE879755, AW876811, AW410336, AA287697, AA775837, AI140761, AI884917, AW005061, AI754907, AI435846, BE395657, AA179409, AA709451, AW162623, AA308436, AA552922, AA130982, AI199766, AI928052, AA604095, AA630255, BE745980, AI064790, BE896123, AI827124, BF593966, BE301056, AA725395, AI954895, AA190882, AI142022, AA468905, AA602522, AA908661, BE047817, BE205857, AI750996, N40320, AA256171, AI744976, AI333054, BE257257, BE746804, AL526007, BF738998, BF696999, BF736575, AI683509, BF032887, N32975, AA455403, AW872811, AA604721, BE740430, AI750686, AW262641, AA256230, BF736574, AI041566, AI797305, AA737976, AI624186, AI858904, AI128163, BG167584, AI289038, AA172072, T15970, AA774045, AI866633, AI460134, H56016, AI208259, BE727215, AA644404, AI367810, BE301051, AA694445, H47768, BE246011, AI598082, AI205248, D53267, AA190850, AA932982, AA130977, AA172232, AI811737, W63689, BE468239, T30196, AI041229, BF904072, AA861665, AA164222, BE901587, AI880574, BF958860, AW057645, AA975875, AI091229, BF761775, AA632936, BE082047, AA057572, AI215900, AA164221, H94409, BE045086, AA243691, H12669, AW068326, AA478826, AA479994, BE045102, AA814694, T08377, AI904545, AA715150, AW083911, R37399, BF904058, BE044984, BE292811, AA478948, BF475721, T73292, R62443, AA989411, BF891529, AI961461, AI810794, AW897867, BF768825, BF980742, R63990, T90361, T55377, BF893923, AA974843, BF893175, BF773199, AA374277, BF893176, T84396, BF891499, BF768813, Z19171, BF892414, BF853058,

0950083 091201

					BF773256, BF852879, AA456043, Z39456, H82670, W02762, T48031, BF893915, AA853298, AW964949, AA588596, BF893145, AA400849, F09862, BF893912, AA304658, AI991422, AA838598, BF773198, BF891527, BF768833, R79677, N69339, AA393476, BF893904, BE243284, AA335036, AW023303, AI784067, R84583, AI868528, AW515655, H09341, BF921244, AI356217, BF893898, T93282, BE161269, BE161272, AA513362, R54399, AW606892, R36847, AW206784, AW606945, BE161260, AW606890, BE161250, AW606895, AW606903, AW606908, BE161249, BE161259, AW606894, BE161254, AW606900, T73306, BE161214, AW606889, AW606884, AW606902, BE161270, AW606951, AA902396, AW606913, W24791, AA632605, AW606907, BE161258, BE161253, BE161252, BE161262, BE161256, BE161263, AW606896, AK026866, AB033767, AK025834, AL035661, AB029636, AF090935, and AL035661.
HUSXP50	2550	561962	1 - 399	15 - 413	BE933735, AL514793, AL515375, AL514731, BG260037, AL514701, AL514885, AL514129, BE047852, AV681902, BF793187, BE887982, AL514829, AI312428, AW103371, AW827228, BF089711, AA225339, BF970449, BF792961, BF904258, AL135025, AL036146, AL513961, AI815855, BG249582, BE729660, AV686346, BE895585, AI340582, AL513763, BG151247, BG108324, AI538085, AA427700, AV727226, AL514929, AI539808, AW946806, AA640779, BG026746, BG179993, AI313320, BG110517, AW105601, AI475394, BE047737, AI571909, AI569583, AL036631, AL036403, BF792469, BE889178, AI537677, BG180996, AI308032, BG058398, BE048135, AV723772, BG030364, BG105895, AL047042, BE048071, BG110684, AW827206, AI802542, BF752252, AI564247, AL079963, AV716568, BE048087, AL119863, AL036901, AW071417, AW151138, AI872545, BE966188, AI345735, BG036846, BF339322, BF812933, BG030785, AW301505, AL135727, AA572758, BE965192, AV681586, AW806761, BG165051, BF527014, BF924882, BG112718, BG109270, BE047952, BF348329, AI872711, AI289937, AL040243, BG180034, AL135661, AL045266, BF343172, BE965432, AV746964, BG168696, BF751308, AI349004, BF882343, AA508692, AI554344, AV723953, BF812960, AW074993, AI800453, BG031815, AI800433, AI521012, BG118829, BE964812, AI633419, BE966388, AI824746, BE047606, AI280747, AV721967, AL036802, BG026428, AI312152, BE544111, BG257535, AW169653, AA470491, BG257920, AW150578, AL514803, BF968205, BF885000,

095008-0401

				<p> AW075084, BF726198, BG164371, BG033403, AW268220, AV757355, BE620444, AW059837, AW302992, AW238730, AI702406, BG027280, AI497733, AI343059, AV682264, AI491852, BF526020, BG026447, AV682809, BE895003, BG112879, AI349933, AI340519, BG259801, AW827276, BG179633, AI554245, AI348897, BG168549, BG222103, AV723204, AI590118, AI621209, AI247193, AI420521, AI922901, AI680388, AI572787, AW999049, AI282281, AW075351, AI925156, BE887488, AI564723, AI610362, AL036361, BF885675, BE894455, AI801325, AI869367, AW148320, AV723062, AI784252, AV682218, BG035511, AW075413, AI570781, BF695032, AI500077, BE172767, AL036980, AI274728, AI498579, AV756393, BF038131, BG034550, BF868928, BF339594, AI862144, AI439717, AL045500, AI873604, AI362637, AL041150, AI433384, BE535358, AI567360, AI207510, BE909398, AI611348, AI344785, AV681857, BE964263, AV734765, AI340627, AI567351, AI344928, AI349614, AI431424, AI343112, AW193134, AI340603, BE781369, AL515047, BE613880, AW268253, AV720938, AW301300, AI567612, AI349598, AI890806, AL036664, AW023590, AI589267, AV682697, BF344652, AI955906, BE886728, AW827249, BF792099, AI612885, AV756838, BF037097, AL022147, AL356015, U72620, AF118064, I48979, I48978, L31396, L31397, AK025339, AF130059, AL122093, AF130105, I89947, AF130075, AF113013, AB047615, AB051158, AL117457, AL137550, S78214, AL133557, AL133606, AR011880, AX019230, AK024538, AF116691, AK026353, AL049314, AF138861, AL050393, AK027096, AR079032, AF146568, A08916, AL133075, AX019229, A08913, AF116639, AK025958, AF116631, AL137459, AF130092, AF090901, I89931, Y11587, AF090943, A65341, AF118070, AF218014, A08910, AK026959, AJ242859, AR087170, AF158248, AL157431, AL442082, AK026542, AF078844, AL137527, AF130104, AF017437, Y16645, AF119875, AL122123, AK000647, AF119878, AL133640, AF111851, AK026741, AK025092, AL117460, AL122098, AF090903, AL050116, AL096744, S68736, AC024171, X82434, AF119899, AL122050, AL110221, AL117394, AL049452, AF104032, AF090934, AL359615, AF116688, AB047904, AK000445, AK025084, AL133016, AL359618, AK026045, AF225424, AF116602, AF116646, AF130082, AF113676, AL110225, AF113694, AF113690, AF116644, AF113677, AK026086, AB048964, I33392, AK000137, AF119909, AF130099, AX046603, AK000432, E02349, AB048954, AL359601, AF242189, AL390167, </p>
--	--	--	--	---

0950087 091201

					AL353940, AL133565, AL122121, AB048953, AL050277, AL050108, A08909, AL442072, AL049938, AF113691, AJ238278, AF090900, AL050146, AB019565, AL049464, AL110196, AL133080, AK000212, AK025772, E07108, AF125948, AB049758, AF177401, AK026592, AL080060, AL359941, AL162083, AL137557, AK025967, AJ000937, AB041801, X70685, AF017152, AL050149, AB052200, AK000618, AF314091, U42766, X63574, AJ012755, AF113019, AX026824, AX026823, A58524, A58523, A93016, A77033, A77035, U00763, AL359596, AF125949, AL080124, AL389982, AK026855, AK026504, AF113699, AK026784, AL162006, AK026532, AK026534, AF111847, AL080137, AK026865, AC006371, AF091084, AK027113, Y11254, AK026583, AF207829, AF183393, AK025491, X84990, AC027716, AC027716, and AC027716.
HUSXY93	2551	658733	1 - 625	15 - 639	AA180188, AI568265, BF750806, AI142552, AA180498, AU157319, D79888, D61752, AI313467, AW589289, AL513597, BE875713, AL121270, AW827211, BF883916, BF968493, AV711509, AV682074, AW071349, AL515173, AL515041, BG260037, BE047863, AI349772, AL121365, AV757737, BE048026, AW827203, AL513907, AI064830, AW268253, AL036146, AL047042, BF971016, AV763915, BG168696, BF054789, BE785905, BG058208, AL513803, BF726421, AL514919, AL119049, BE048081, AV681668, AV682466, BE048071, BF724691, AV655645, BG108324, AL514627, AI868831, BE964812, AV681951, BF795712, AV710479, AL135661, AL119791, BF793187, AV733397, AI608667, AI436456, AV708119, BF970162, AL040169, AV681630, AV682330, AV758822, AV682252, AV758592, BE048163, BG179993, AW303152, BG108147, AI687376, AI349645, AI500077, BF793644, AL045500, BF791874, BG110797, AI815383, BF343172, AI567351, AV717299, AV729890, AI349004, BG036846, AV682476, AW089572, AW162071, AI433157, AV758110, AV682809, AI440426, BG151247, AL036396, BG250190, AL120854, AI521012, AL514691, AL515047, AV682249, AI207510, AI433976, AV682772, BE965095, AV756393, AI312152, BE887488, AL047763, BG257535, AL046849, BE964700, BE048179, AI349937, AV734318, AI873731, AV681857, AV758179, AW074993, AV758217, AI687728, AL513553, AV757598, AL513911, AV723772, AV756122, AV758806, AV764282, AV755290, AI568870, AI285735, BG259801, BE877769, AV682222, AV756477, AI702406, AI538716, AW117882, AV755613, AV729334, BF969494, AV704928, AV682441, BF673434,

09050083 091201

				<p> AV682521, BF812933, AA640779, AI343112, AA613907, AL514791, AV723953, AI349256, AL036802, BG104782, AL514803, AV711924, AV723204, AV682672, BE048319, AI906328, BF348329, AL513643, AV706777, AI580190, BG252929, AV755207, AV710608, BG109270, AI340582, AI349933, BE881155, AI679724, BG178809, BG179633, AL513631, AV756703, BG180996, BF726322, BG121222, BE965556, AL514261, BF726297, BE964486, AV682479, AV682266, AW166645, AV705644, AV756560, AV681638, BE881061, AL036980, AW071417, AI282655, AL514935, AV757639, BG107847, BE018711, AV757943, BG029399, AI349614, AV755581, AV681586, AV723062, AV709517, BF981774, BE613622, AW080838, AI475371, BG252914, AV682289, BF037097, AV756619, AV682218, BF882343, AI920968, BF970446, AL513693, AW827249, AV682082, AI500553, AV682351, AV757797, BG109125, BF968041, AV757158, AI934036, AV695052, AV733326, AW238730, AI699857, AV727776, AV757455, AI907070, AI609592, AI583316, AV681872, AV755614, BG105099, AV755668, BE783707, AI469532, AW301409, AI863014, AW467961, AI281779, AV681869, AI349598, AL514299, AC011891, AF130059, I48979, AF116644, AF116691, AL133606, AF130105, AR079032, AF118070, AF078844, AK000212, AF130075, AL157431, S78214, L31396, AL442082, L31397, AF116639, AF116631, AF116646, AF130082, AL133640, AF090900, AF113691, AJ242859, AF130104, AF090934, AL442072, AL050393, AF118064, Y11587, AF113694, AF116602, AL133016, AF090943, AL389978, AF090903, AF113013, AF138861, AX046603, AL049938, AL390167, AF125949, AL049452, AF104032, AF218014, AL117457, AB048953, AF119878, AK026608, AF090901, AL080060, AL110196, AL110221, AL117460, AL050146, I89947, A93016, S68736, AL137527, AF111847, AF113676, AL359596, AB049758, AL359601, AX019230, AF116688, A08916, AL122050, AF113690, AX019229, AF130092, AF106862, AB048964, AF090896, AR059958, AL049314, AK026865, AF113689, AL050149, AB050510, AF119875, AK026741, I89931, U42766, AB047615, AK025339, X84990, AL050116, Y16645, AB041801, AL133075, A08913, AL050108, AK025958, AL162006, AF113677, AK025084, AL162083, AB019565, AF017152, AL133557, AK026045, AL049466, AL133080, AL096744, AL080124, AF219137, AL122093, AF113019, AL050277, AF113699, AK026744, AK000618, AL133093, I48978, AL080137, Y11254, AR011880, AK026855, AL133565, </p>
--	--	--	--	---

0950087 091201

					AF119899, AF097996, AF146568, AL137283, AL049430, AL137557, AF111851, AL137459, AX006092, E03348, AF116649, AL389982, AJ000937, X63574, AL122123, AK027096, AK025092, AK026784, AF158248, AL122121, U91329, AK000083, AL359618, AF119871, AL117394, E07361, AF091084, AF119909, AF314091, AL133560, AK025772, AF207829, AK026533, AK026542, AF079765, AK000137, X82434, AK026452, AL110225, AL049300, U00763, AF242189, AF125948, AF130099, AF271350, AL050138, AL359941, AB048954, AK026642, AK000614, AL133258, AF177401, AK000445, AK026927, A65341, AB048974, AK026647, AK026592, AR087170, AB051158, E07108, AF017437, E02349, AK026534, AL050024, AK026583, AJ238278, AB052191, AK024538, AK026959, AF130066, AK000652, AF177336, AK025491, AL359615, AL353940, AL137550, E05822, AF225424, AL049464, AB047904, AL117583, AK026353, AL117585, AF116682, AL122110, AF067728, AX042059, S61953, AK027113, AK026504, AK026480, AL049382, AK026528, AL353802, A08910, AF116610, AL117435, AK025391, AF130110, AC002467, AK026532, AK025967, X70685, A08912, AF091512, AK000432, AK026086, A77033, A77035, AF130077, U35846, AK025414, AC007390, AF183393, I33392, AB049892, AK000323, A03736, AC006371, AL122098, AL133344, AL359583, AK027164, AF130087, AB052200, AF061943, AC011891, AC011891, AC011891, AC073232, and AC073232.
HUSYG26	2552	709709	1 - 730	15 - 744	BG251809, AI380185, BE856114, AI654549, BE139193, AI656314, AI654829, AI783769, AI656635, AA853364, H54036, AI968574, AI799978, BF821711, AI831492, AW009524, AW858522, AL038878, AL045327, AL135012, AL134524, AL045328, AL042898, AW207831, AL047163, AL045891, AL047611, AW601637, U46344, AW979234, AW577201, AW577199, AL043089, AL043321, AL134110, AL042655, AL042488, AL046356, AI431321, AI432644, AI431307, AI431316, AL042802, AL043239, AL047037, AW975391, AL042515, AL047092, AL042931, AI431238, AI432666, AW971301, AI304382, AL041862, AL043091, AL042533, AL042853, AL043196, AL119324, AW979127, AL042729, AL042787, AW970540, AI432653, AI431323, AW976043, AL079977, AI440238, AI623302, AI926593, AL040207, AW975657, AW973652, AI567971, AL047675, AI539260, AI610557, BG252929, BE897632, AL040459, AI500658, BF084778, BE883591, AW969988,

09950083-091201

				<p> AI436429, AW972292, AW971404, AI952433, AW974909, AW972662, AW970576, AI866465, AL042745, AW974098, AI433157, AI432656, AW151136, AI539771, AW968204, AI537677, AI494201, AI866786, AW172723, AI500659, AW975929, AW971066, AI815232, AI801325, AI500523, BG029667, AI225248, AW973397, AI582932, AI923989, AI284517, AI500706, AI445237, AI491776, AW151138, AI521560, AI889189, AI500662, AI284509, AI889168, AI866573, AI633493, AI434256, BE886728, AI805769, AI888661, AI284513, AI888118, AI889147, AI371228, AI440252, AW968355, AI648567, AI690946, AW089006, AI440263, AL042523, AI561170, AW976010, AW074057, AW973377, AL042939, AL038529, AW975638, AW970966, AL515375, AI538850, AI887775, AL042420, AI590043, AI872423, AW973717, AI289791, AI633468, AW974795, AI804505, AI582912, AI539800, AI538885, AI554821, AI927233, BG110517, AI866469, AV684075, AI434242, BE964089, AI500714, AI285439, BE885490, AI859991, AI623736, AI355779, AW968347, AI581033, AI491710, AW979134, AW972927, AI890907, AI860003, AI242736, AI828574, AV682326, AI887499, AW972649, AI866691, AI539781, AI702065, AI539707, AI885949, AI285419, AW089557, AI559957, AI521571, AI469775, AL045620, AI866581, AW020381, AW971269, BF338002, AI493559, AI567953, AI815150, AI446495, AI867068, AW971167, AW968212, AW979105, AW970070, BG257535, AI872310, BG113493, AI431315, AV736134, AA768018, BF795712, AW975152, AW975858, AW979080, AI698352, BF812963, AL042508, AL039011, BG104735, AW022682, AW976024, BF811804, BF797072, AI697243, BE788683, AI371229, AA279293, AW858243, AW971290, AW971737, AW975169, BF343394, AW968336, BE886773, AW972594, AW151979, AW877209, AR076189, AB007144, AB022341, L10353, AL133053, AL133049, AL122101, AR066494, AL133074, A93931, AX030435, X61399, AF178432, AL137254, S61953, AL137627, AX040958, A85203, U30290, AK026642, AF124728, Y17793, AX040974, AF067420, AL122049, A93923, AL133070, AK000636, AL110159, AL353935, AF260566, AL157433, AK025383, A18777, AF207750, AF028823, AL137478, U42031, A93916, AL137479, I89947, AL353625, AL133067, L40363, AK026522, AL133076, AL117583, U96683, AF130099, E01314, I25048, I25049, AL080124, S75997, AL389978, AF119871, AF132730, AF030513, I41145, AK000421, </p>
--	--	--	--	---

					AF116649, AK025772, AF067790, U67958, AF038847, AK000432, AF242376, AK025239, AL389935, Y14314, AK026865, X63410, AF130055, AL359583, AK025209, AF113676, AF199027, AL117435, X79812, A65341, I26207, AF031147, AF305835, AF111847, AL080126, AF182215, A08910, AC024247, AF111112, A08907, A08909, AL359622, S36676, AK025906, AK027103, AB048881, X84990, AK000753, AL122118, U77594, AF002985, AF013214, AF199509, I48978, A08908, AL049382, AF130100, AB047947, AB048974, AX020124, AL162062, AK026434, I66342, S77771, AR034821, AR081734, AF061943, U49908, AK026591, AK026534, AL133084, AF116644, L13297, E03671, AF203473, AK025015, AF130082, AK024622, AR050959, AK000652, Y16258, Y16257, E02756, Y16256, AL137526, AL050277, AL133093, D44497, E01812, A08913, AK025708, AF217991, AF180525, AF078844, I89931, AK024992, AL137271, Z82022, AK025407, X52128, AL122098, U68233, I92592, AF116646, AL137711, AF090886, AF274348, AF274347, AK026038, AL137658, AK026551, A12297, A08912, AK000647, AL137292, AF057300, AF057299, AF035161, A08911, AR038854, AR087170, AF003737, AK024594, AK027160, AL049466, AF026008, E06743, AF004162, AL133113, AK026649, I00734, AR016469, AF119883, AB016226, AF114170, AF100931, S63521, AL137557, AF225424, AK027161, AF111851, AK026885, L30117, AK027164, AK025435, AF185614, AL133075, E00617, E00717, E00778, AL389951, AR011880, S76508, AL161802, AC004227, X81464, AK000450, AK026592, AK027105, AJ001838, AF082526, AX046716, AL117626, AL080154, AB051158, AF110640, AK024974, AR059958, AK027217, AL117585, AK026603, U72621, AK025573, AF026124, E12580, AL117457, AK000323, AL133072, AF019298, Z97214, AK024546, AF116682, I89934, AK026633, X66862, AL157878, AF119899, AL109672, X87582, I33392, AR029490, E02349, AB025103, AF176651, AL117416, AK026797, AK000655, AC011488, AC011488, and AC011488.
HUVCQ68	2553	561963	1 - 664	15 - 678	BG169321, BG258652, BE539719, BF663612, AV759029, BF239202, BF056223, AV716151, BG254787, AV759032, BE265426, BE561524, BF344287, AW274631, AV715869, AV716258, BE395535, AW602195, AI784535, AA994106, BF590282, AI813617, BE545672, AV761057, AI743389, BF839256, N90862, AI277996, AA906652, AV738454, AI313268, AA469453, AV735720, AA526368, AV749084, AI079353, AI087923, AV739098,

					BF738960, AA677113, AI860597, AA677095, AW860443, AW860480, BF680311, AV741458, AA577146, AV736779, BE931596, AW610366, AI367976, AV757929, AV747366, AV742892, AV749082, AV712419, AI370907, AV743113, N26385, AV742865, AA305331, AA962294, W17085, AV736074, N30813, BF930605, BF351056, N40654, AW821817, AV735693, AV744019, BF906764, AV738002, AV712773, AA976489, AA938500, AV738417, AW610347, BF359055, T62963, AA165027, T49805, AW860654, AI423799, AA808220, AV741758, AV735472, AV757936, BF095707, AW860655, AW015834, AW862529, BF845386, AI279747, AW406446, BE091434, AA760863, AW840037, AV745065, AV735634, AV735823, T49806, AW404421, AW934901, BE810650, R01789, N41587, H67554, AV744825, R65657, AA467782, AA361428, T63214, AI245538, BF211812, AW406745, AA244301, AW869801, AA688351, AI907982, D63243, AI760888, BE810551, AA371270, BF738943, AI766627, AA299296, AW834749, AA630775, BE810585, AV748156, AW674915, BE934944, AI766607, AV743011, BF737038, BE070301, AA634593, AA176219, N41720, AW103363, AW083134, AI128418, AW957387, AW071816, BE810575, AV743629, AW264479, AA357680, AV743545, AV750838, AW148506, AA922790, BF815921, AV749802, AW079315, BE702488, T26925, AI305511, AI309416, AI611892, AI252475, AI371614, AI252598, AI251486, AA973571, AW075042, BE138512, BF594565, AF053233, AF053724, AF045661, I56674, and AF132812.
HUVDG58	2554	895443	1 - 2771	15 - 2785	BG109143, AV756057, AC012312, AC012312, and AC011205.
HUVEG53	2555	797556	1 - 2149	15 - 2163	BE386016, AW958825, AI924134, AA847884, AI290465, AA149630, AA825487, AA149634, AA128265, BF849323, AA149498, N36935, AA744599, AA744600, AA825175, AA306759, AI280357, AA253422, AA789124, AA128306, AI222734, AA361836, AI298984, AW876085, AW028634, AI916423, AI382702, BF542170, AA149502, N46208, AA779865, T52521, AA046646, T52599, AI817864, AA993703, AA158418, BE385026, AF282265, AL121585, AC016526, AA252884, AP002793, AC021454, and AC021454.
HWA AH11	2556	815548	1 - 1264	15 - 1278	AL535070, AL533539, BE748937, AL527179, AW083127, AW327290, BE205822, AI798681, AW410645, BF063649, AI797284, AW183981, AI096678, AI092936, AI144400, BE867172, AA236980, BF035835, AI685511, AI860643, AU146089, W46771, AI074319,

09950083 091201

				<p> AW513360, AL527769, AW273299, AI801043, AI521427, AI885247, AW300673, AW295119, AI457885, AW327628, AI399827, AI095832, AI924126, AW137947, AA745647, AW295118, AW029034, AI381953, AI393856, AA813851, AI811198, R43938, AI458562, AW151914, AI823612, R16113, AA651935, T62960, AW473030, AW198081, T16212, BF940829, AA430674, Z39341, F09974, AA884361, AW197763, T63211, AI493592, AW971745, AW662317, R40698, AI926977, BF002645, AI420802, AA861954, BE042395, F02261, BG055863, AA664368, T32088, AA853151, BE613349, BE612945, BE622828, BE613070, BE613136, BE614307, BE622931, BE621911, BE614230, BE614237, BE622925, BE074372, AI473459, BE618811, BE622868, BE620742, BE614451, AA235918, AW961017, AW263814, T99938, C01020, BF759319, BF939920, AV645401, AV645332, AV645383, AV645377, AV645379, AV645385, AV645389, AV645369, AV645380, AV645321, AV681527, AV645390, AV681501, AV645400, AV645392, AV645393, AV645404, AV645353, AV681458, AV681492, AV681475, AV645358, AV681500, AV681509, AV681529, AV645334, AV681510, AV681528, AV681474, AV681514, AV645339, AV681477, AV681512, AV681471, AV681491, AV681526, AV681472, AV645336, AV645317, AV681507, AV681465, AV681488, AV681525, AV681495, AV681505, AV681502, AV681504, AV681486, AV681483, AV681464, AV681462, AV681487, AV681506, AV681523, AV681519, BE620756, AV681497, BE066319, BF742482, AV681531, AV681461, AV645343, AV645344, AV681468, AV645350, BF516089, AV681513, AA404569, AL134524, AW861944, Z99396, AW804686, AW858525, AW877209, AW604723, AL119324, BE085487, AK021918, AR080280, I25027, AR054109, I44515, I26928, I26930, I26927, I25041, AR069374, AR069375, I44516, AR035224, A49045, A94046, A94054, AX009487, AR093384, AR009151, AR027099, AR083151, AR091518, I85513, AR009152, AR093392, AR093383, AR038321, AR038307, I05393, A10617, AR028792, A01324, A01323, AR034783, AX030966, I63120, AR067733, AX009486, AX029455, AR064322, AR064323, AR064320, AR064321, A94048, A94061, AX027785, AX003194, A32110, A83642, A83643, AX035462, A70359, AR019094, A92666, A92668, I49890, A92667, A92665, I05430, AX032758, AR028791, </p>
--	--	--	--	---

					AR028793, AR019098, AR067734, AR069417, AR018924, AR018923, A48774, A48775, AR000006, AR015960, AR000007, AR015961, AR051652, AR067731, AR067732, AR091393, AR020199, AR020200, AR001287, AR020198, AR020197, AR029418, AR029417, I89986, A91752, AX008865, AX008867, AX008868, AX024906, AR069413, AX033488, AX033489, AX033490, A92081, A92080, A92077, AX033474, A92078, AX033486, A92079, AX033487, AX030369, AX030368, AX036660, AX036661, A91751, AR091571, A29109, A32111, AR068550, I58322, I58323, A23373, AR068551, AR062871, AR068508, AR068510, AR068509, AR085090, AR003585, AR051957, A91754, AX026821, A63067, A51047, A63064, A63072, AR031375, AR068507, AR068506, AX001323, AX001324, A85308, AX044426, AX044425, A60213, AX046332, A44171, AX027908, AX027904, AX027906, A49700, A60207, A60208, I58669, A58521, AX004550, I09121, Y14971, AX030435, AR031374, I07209, I07249, A63954, A93444, AX046357, A46342, AB026436, A46343, AR066494, AR055065, AX040911, AR054110, AJ251859, AR060234, A30600, AR055066, AR032878, A81671, AJ279014, and AR069079.
HWAAQ28	2557	806588	1 - 1635	15 - 1649	AL522894, AL519650, AL522935, AL533662, AL522936, AL516787, AL529748, AL516786, BE747239, BE617640, AL519651, BE900926, BE253756, BG107583, AL522224, BE796747, BE408146, BG032407, BF984483, BG106698, AL522223, BE902939, BE882144, BE901801, AV716026, BE903519, BF795428, BE788045, BG259257, BE744493, BF796663, BE870681, BF531083, BE735617, BE780655, BE745064, BE780676, BE782159, BE616997, BF574609, AW363345, BF690512, BG259249, AV733450, BE541553, AW857780, BF033038, BG027712, BE298540, AV734792, AV701418, BE297543, BF306362, BE273544, BE389895, BG027225, AA778572, BE278438, BE259346, AW411199, BE785598, AV762649, BE882851, BE789480, AI859450, BE276662, BE882706, BE935907, BF129054, BE548402, AA313653, BE893454, AW578901, N25766, AA147730, BF974999, AV701113, BG120751, AA557185, N53489, AA402500, AA176091, AV737877, AV701039, AA148269, AI291236, BE540583, BE871734, AI338866, N41826, BG026635, BE246137, BE167827, BE778520, BE735402, AA148268, W84337, AA775692, BG121372, BE501636, AW673432, AW411200, AW169109, AW792918, AA399165, AW403612, BE674822, AA310007, N20938, AI754518, AA147927, AW148838, BF689607,

					AA411829, BE392081, AI888692, AI264552, AI925596, BF131873, AI336357, AI352436, BF848204, N27806, AW381860, W47285, U56657, BF155187, AI366728, AI422692, AA223352, BF780107, AI280252, AI140153, AI215713, AI187089, W69118, AA789050, AI355599, BF197763, N98638, AA401899, AI458428, AI143209, AI090686, AV659444, AI801394, AI274124, BF337688, N23199, AW402897, AI056723, AA259129, AA010489, AI814051, AI308755, W00505, AI302555, AW467941, AA403236, BF698327, AW128893, AA259168, W46942, AW241367, AI802697, H16280, AA983554, AA443079, AI089372, AI207337, AA176331, BE244114, AV735304, BF431994, BE246202, AV659570, AI928874, AI361880, R86971, N24694, AA182539, AA424931, BG170548, AI750314, AA250748, AI187741, AI085164, AA931922, AA251055, AA404960, AI262880, N66091, AI184131, R91324, AW403547, AA837851, H95502, AW513449, W60526, AA947001, C18057, W69243, R86891, AA291507, AA182584, BF743772, AA223252, AA631660, AA305586, AI190144, N23575, N42504, H50935, U56658, H98852, AV751245, N36673, AI688894, BF088492, AA594292, BE902663, R85340, AA927346, AA814039, N70482, AW675759, AA357201, AA962443, AA458576, H64241, AV727253, BF886646, H70416, H62700, AA393658, T64619, AI750315, H17996, N72226, AA854016, AI150407, AA718956, BF091940, W03839, AW050797, H78347, AI472076, T78672, AI805357, AA312825, AF161518, AX013066, AK000481, AK000586, AL109806, AK026118, AF161513, AF117231, AB047911, Y17793, AC011530, T55382, T55810, T93718, T96182, R18951, R44065, R44065, H17995, H45107, H45416, R85341, R91285, R94337, H51598, H58870, H59028, H62786, H70417, H81902, H81994, H95006, H95011, H94981, N28011, N36483, N53392, N72492, N75347, N93569, N93712, W01855, W21099, W21565, W25640, W56213, W93366, AA011229, AA027185, AA027184, AA027978, AA027977, AA035623, AA054329, AA054239, AA062963, AA064913, AA064914, AA464280, and AA425869.
HWAAY60	2558	745917	1 - 1713	15 - 1727	AL530992, AL530991, BG251146, BE395713, AW955083, AI685470, BF037254, AW195758, BE395793, BG167395, AA477698, BG255897, AW025151, AA206118, AI684900, AW071954, AI042479, AI076835, AI052688, AI925453, AW299605, AI638447, BE551207, BF194856, BG149868, BG057307, AW299615, BE896641, BE392847, BE546040, AA477699, AL526556, N41705, W85698, BE007568, AI014560,

					<p>AI655938, AA243713, AA579661, AA858009, AW251059, BF197312, AI081769, AA827676, BE671162, N54544, BE045652, AW138173, AA026397, AA233066, BF971815, BE220154, AI932354, AI660197, BE504629, AI912697, N23942, AW025431, BE901604, T69708, AW263820, AW166895, BE504641, AI200982, BE504651, AA461050, AW151190, AI379930, AI298201, AI079400, AA975242, AI083591, AA810573, AL522056, AW293145, AI086200, AA704156, T62655, N49202, AI831159, AW188188, AU150237, AI269728, BE042520, W79510, AI805447, AA811276, BF975273, BE898722, AI342865, AI200807, R49471, AI866234, AA009526, W92371, AI907500, AA576912, BG250763, AA769354, AI972137, AA284866, AA121798, AI631573, AA034488, AI624884, AA833829, AI096559, AI637897, AW272844, AA232909, AW016376, AA890160, AI567962, AI361574, AI969404, AI341367, AI147028, W79609, AW182166, AA744718, AW014569, AI635075, H08393, AA228932, W46662, AI766531, W46571, AW002747, AI539697, AW952410, AA255548, AA977178, AA858173, W78930, AI916921, AI392713, AA494131, R96990, T16336, BF448099, AI754998, AW579441, AA551555, AA652299, AA365158, AA626307, AI074338, R33109, BE047100, T62509, AW062453, AW374921, BE735465, N54838, H10058, AI830620, AA352728, R38835, BE614757, AI393236, AA476422, Z38719, BF309688, AA055575, BE617654, AI683165, N76247, AW062437, AL522055, N56492, AI797298, AI833138, BE622395, BE904767, R35675, T73753, R33206, AA460482, AA428438, AA256662, BE827794, AA009841, AI500338, AW946922, AA732166, AW946741, AW946850, BF336785, AA026453, AA328088, AA625258, BE165782, AI685438, BE171649, T70365, AA969716, BE622838, AI422223, R31637, BE737245, R35786, BE827776, AA070038, AI670707, BF307717, AI244096, AA095522, AW295268, AI539150, BG235892, BG153026, AI769916, BE896979, N72615, AA644486, BF308421, BF924005, BF205005, AW938856, AW627518, BG260881, AV736507, W92370, AA009874, AI870275, BE706786, BF350564, BE706658, W28025, W22956, AU128965, AL533305, AW798754, AL390195, AK022860, AF052183, AL390195, AL390195, AL390195, and AL356387.</p>
HWABR43	2559	823367	1 - 2300	15 - 2314	<p>AI240662, H52198, H52199, H47421, H47334, H63365, AW500029, AW503420, BF681619, AV710482, BG236628, AW504485, AI732151, AA574442, BF768846, AV758790, BF677892, AA610509, AA402129, AI357823, BG029224, AU152561,</p>

				<p> AL120343, AI354423, AW805539, AW069227, AV762741, AA634837, AA191418, AL118991, BF681607, AW021917, AA584489, AI753488, AL079734, BE139267, AV760918, AV762633, AA533176, BE147833, AA410788, AW518220, AW814024, AW270258, AW855803, AW855730, BG180976, BF965775, AW974932, AV711465, AV711430, AI040051, AA904275, AI963720, BE138594, AV761745, AI345157, AI279417, AI056177, AI754037, AW468372, AW438542, BF965290, AW805547, AW327624, AV761862, AW302903, BF769368, BF984807, AI085242, AW271904, AW271917, AA490183, H05940, AA833875, AA833896, AA491284, BE019467, AA084766, AW168433, AW069769, AI537020, AV763847, AW190505, BE294700, AW963904, AW270256, AV710066, AI270117, AI431303, AA581903, AI799607, AU118852, AV758903, AA226584, AI625604, AW302315, AW964231, AA682912, AI791150, BF839844, AA654321, AW975049, AI267356, AW191886, AI267450, AA457249, AI369580, AI908575, AV763026, AV763058, AI754767, BE062478, BF760919, BF477463, AV761106, AI004591, AI859946, AI865905, AW504299, AA862243, AL037632, AU140464, BE395137, AI278089, BE062476, AW517743, AW833862, AV764340, AU147162, AC006388, AC005089, AC005015, AL138741, AC005102, AC002425, Z82206, AC011290, AL096840, AC005995, AC005399, AC004596, AC008569, AC020913, AC005778, AL355385, AC010510, AC005081, AC002301, AL121897, AL138807, AC008753, AF111167, AC001226, AC005666, AL445248, AC005484, AL022476, AC011465, AC011895, AP000557, AC007731, AC024571, AL353748, AL109976, AC004098, AL139100, AL109825, AC005330, AX039602, AC005619, AC009086, AC004905, U95090, AL135928, AC005231, AC006430, AL121890, AL024507, AC004089, AL138752, AC009087, AF109907, AF196779, AC007383, AC002492, AC005236, L78833, AC004707, AP000065, AC005500, AC008745, AC011497, AC005488, AP000689, AC005037, AL031432, AB003151, AC006262, AC005004, AC004858, AL135744, AL136137, AC005377, AL121586, AC005747, AC009953, AC011479, AF196972, AC016637, AC018633, AC004531, AC006013, AC006441, AL008719, AL031311, AC021016, AC020955, AL133551, AL139188, AL049697, AL109840, AC084732, AL159168, </p>
--	--	--	--	---

					AC007376, AC020898, AC002558, AC016025, AL034379, AC006449, AC005274, AC007136, AL049553, AC010271, AP000113, AP000045, AC083863, AP001745, AC010605, AC003070, AL096870, AC010328, AC004882, AL135927, AC007227, AC010422, AD000092, AL121751, AL357560, AL035555, AC003046, AL356299, AL109758, AC011464, AC010206, AL109627, AC005899, AL158040, AC007421, AC008736, AC005971, AL121747, AB023049, AL035422, AL022323, AL031577, AC007722, AC018758, AL078633, AL031778, AL031447, AL031602, AC010618, AL035704, AC004151, AC006238, AL022163, AL139286, Z82178, AL080243, AC022154, AC005837, AL161670, AC004884, AC021999, AP000503, AF134726, AL136418, AL354816, AC004019, AC020754, AL136223, AC011455, AC003108, AC009244, AC010102, AC007956, Z85986, Z98200, AC004386, AP000692, AC005920, AL034420, AC005071, AP001717, U78027, Z84466, AL132639, AC006055, AL109935, U66059, AL021155, AL356801, AC007199, AL049552, AC005480, AL138717, AL139384, AC004638, AC005664, AC007963, AL096791, AC005291, AC002369, AC002477, AC009756, AL035461, AL031650, AL008627, AL109984, AC006064, AC003071, AJ400877, AC005295, AC004815, AC021036, AL161802, AF243527, AC005216, AC053467, AC002073, Z95115, AC005261, AL135902, AL117381, Z93015, AC020947, AP001759, AL049793, AC002565, AL138706, AC078833, AC007739, AC008403, AC005387, AL163301, AC005911, AC006547, AL021546, AC005736, AL021808, AL138721, AC010282, AC007282, AP001630, AL035659, AP001619, AL136969, AC006014, AJ277546, AL050341, AC007030, AL162272, AC004408, AL138976, AC004150, AP000030, AP000212, AP000134, and AL049766.
HWACH06	2560	799620	1 - 1147	15 - 1161	BG112811, BE540349, AU160354, BE877776, AW964647, BG248203, AW080842, AI126237, AI738708, BF437231, AI288602, AA033525, AI200312, AA410504, AW293916, AA033526, AA070152, AW182578, AW674004, AA961647, BE049558, AA808104, BE301368, T82022, AA410505, AA884100, AI635980, AA580797, AA748547, AA322668, BF811398, T71568, AI004853, AA975024, BE093091, BF871651, BF128606, AI674097, AA988905, BF804322, AL137420, AL133609, AK025653, AK024038, AC016642, and Z94722.
HWACZ33	2561	827311	1 - 1448	15 - 1462	AI992047, BE247516, AI380304, BF848604, BF872804, and BF872803.

0950083-091201

HWADV90	2562	897732	1 - 2379	15 - 2393	AI793141, AL042110, AA682890, AI268385, W46143, AW294829, AW517042, H12201, BF511080, AI027880, W59940, AA284296, BE300890, AW572488, AI277625, F02781, H16962, BF243115, AA558997, W59939, AA026524, AA584482, AW069762, AI080618, AA577906, AW955706, AA593471, AI792578, AU152722, AI054419, AI919265, AW327624, BF678990, R15790, AL037632, AA583394, AA651639, AI284640, AW407578, H14377, AW960468, AI580250, AA533176, AV762979, AA469327, AW675551, AV756848, BF991208, AV755512, BF934742, AV710482, AW021917, BE908602, AA191418, BF681619, AL048925, BF839800, AV761156, AI243789, AA574442, AI872216, AA904275, AI371429, AW020088, BE150580, AI357823, AA078849, AU147529, AA610509, AW970915, BE393367, BE968744, H15652, BE085098, AA528546, BF849430, AI306028, AW969928, AI339845, AA134335, AA829065, AW022897, AV763847, BE019467, AW438542, AW979096, AA229975, BF965775, AA749235, AW327591, BG236628, AL079734, AW069227, BE883893, AA644090, AI732865, AI056177, AV715301, AV728912, AI862716, AA682912, AW970078, AW021583, AW963117, AP001761, AP001478, AC002055, AF270529, AF229525, AL078621, AC004908, AL031259, AF327129, AC009410, AP001724, AL117345, AC008555, AC025809, AP001695, AL391867, AP001760, AL136137, AL163280, AC005020, AL049871, AC005186, AC006961, AP001688, AP001671, AC006150, AC020912, AL365475, AC005304, AC027269, AJ229041, AC011288, AP001694, AL035704, AC002492, AP001727, AP001698, AC007969, AL163248, AC020754, AC010605, AC021999, AC002994, AC007298, AC009303, AL137141, AL159977, AL031432, AP001699, AL133467, AL157688, AC009123, AL163301, AC005047, AP001435, AP001715, AL121869, AP000704, AC022448, AL163278, AP001707, AP000566, AP001706, AC004887, AF131216, U91323, AC007677, AP000246, AL133379, AC007225, AP001596, AC006231, AL356103, AP001696, AC004963, AP000008, AP000036, AJ252312, AP001710, AL022329, AP000099, AC020647, AJ252311, AJ252313, AL442167, AP001676, AP000093, AC002456, AC015971, AL163285, AC010209, AC007883, AL135746, AP001700, AP001601, AP001730, AP001689, AC006210, AC012000, AL133243, AC017002, AL354766, AC011890, AL031431, AP000237, AP001678, AL357519, AC008556, AP001254, AL021368, AC009405, AC005539, AC022212, Z95152, AL121975,
---------	------	--------	----------	-----------	--

					AC002369, AL021407, AL034402, AL136111, AC004815, AL137191, AL133395, AC020552, AL035669, AL021939, AL163283, AC003682, AC009263, AC073651, AC017047, AL109967, AC010363, AC006257, AC009116, AL109918, Z93015, AL354858, AP000108, AP000040, AC026126, AC010451, AL359457, AL355540, AP000144, AL078583, AL161445, AL133372, AL354984, AL359174, AL356288, AC005066, AP001683, AF064865, AL109758, AC002351, AL354674, AF265340, AL163268, AL137230, AP001256, AL359238, AP000567, AL135787, AC012039, AL033392, AL035671, AL050350, AB041512, AC005587, AL138976, AC009481, AB020863, AC024947, AC007034, Z93931, AC009087, AL117694, AC013434, AL121656, AC016554, AC002299, AC005701, AC019155, AC008066, AP001667, AC000075, AC024085, AL137251, AL009177, AC005227, AP000208, AP000130, AP000282, AP000207, AP000129, AL118517, AC006399, AL359382, AL390027, AC002559, AC007564, AL158040, AP001708, AL359641, AC004999, AC006948, AP001705, AC034242, AC004063, Z82179, AC018698, AC005034, AC004953, AL157879, AL121761, AP002530, AL031393, AC019140, AC006337, AC006530, AC009508, AL117356, AC007850, AL158141, AP000472, AL049760, AL136136, AC009502, AL163246, AL078603, AC005598, AC005181, AL365367, AC078843, AC003009, AF241728, AC004012, AC005519, AL133238, AC073898, AP001718, AL161731, Z92547, AC010348, AC004908, and AC004908.
HWAEB52	2563	873207	1 - 2179	15 - 2193	AI657485, AL532472, AL532471, AU123421, AL040049, AA152049, AW960660, AI123690, BF724895, BE890141, BF569925, AI671820, AW959495, BF590288, BE646576, AW779655, AA524015, AW015538, AI743339, AA398719, AU159563, AW058623, AI675998, BG248946, AI858508, BE782389, BE547088, AI638603, AU150408, BE710171, BF196797, AI640308, N23869, W92992, W68625, AU154693, BE169553, AW994753, BE813521, AI671374, AW516075, BE813670, BE169351, AW518935, AW771500, BF063773, AI668815, AI738979, BE169552, BE546935, BE169730, AI743066, AI306547, BE932233, AI867416, BF224229, BF087906, AI636926, AA149820, AI339396, BE883189, W68626, AI475630, AI937656, AI808394, AI936620, AI268424, AI470087, BE217978, AA203500, BE710141, AA055470, BF995655, AA602512, AI085310, R69944, H66476, AI745272, AA749312, BF745582, BE932304, AI189504, BE813728, AA305216, AI147197, BE813697, N51913, AV706826, AI769119, AI699496,

					<p>AW020350, BF995501, AW086038, H59673, AW579059, H85493, AA393390, H92627, AA707630, AI918818, BF770668, AW960662, BF770665, H96084, H92410, AI991427, BF945598, AI129883, D79378, AW008020, AI307151, AA971636, AI650973, R92009, AA431716, BE902337, R62846, AI380238, AA988827, AA362712, R92771, AW749913, AW749905, AW749958, AW749918, AW749914, AW749951, AW749916, BF989914, BG005357, AW749908, AW749952, AI474389, R63844, AI525907, H62659, R37909, H66475, AW406064, BG005353, R37826, AA297824, R24456, BG005354, R87652, R84289, T77910, AI802420, AW593746, BF811262, R64458, AA303679, BG008910, AW839281, AI908514, AA311599, AW131892, AW402961, BE883450, BF946388, BF946375, BF947681, T77826, BG230830, BE155648, BE542412, AA055737, BG008047, H58202, BG258450, AW877209, AL119399, AL119319, AL119324, AL119457, AW392670, AL119443, AW804686, AW604723, BE695785, AW858526, AW861944, AW858525, AL119497, AL119363, AL119444, BF755702, AL119391, U46350, U46347, U46351, BF868684, AL119484, AL042544, U46341, BE705903, BE705906, AW577135, AW372827, AL119483, AW363220, AW971745, AW384394, U46349, AW861889, AW858455, Z99396, AL119439, AL119355, AL119418, AL119464, U46346, AL134531, AL119522, AI142131, AL037205, AL119401, AL134920, AL134538, AL119341, AL119335, BF868697, AW604726, BF868687, BE705905, AL119496, AL134533, U46345, AK023281, AK001170, AF113218, AB042827, AR066494, AX030435, AB026436, AJ251859, AR060234, AX046357, AJ279014, AR054110, A81671, and AR069079.</p>
HWBAK71	2564	853581	1 - 358	15 - 372	<p>AV705433, BE877146, AV654282, BE881230, AI557222, AV725709, AI064816, AV756682, AV702947, AA467922, AA469321, BE876183, BE878467, AV715748, AV727472, AV757055, AA467864, AV726503, AV717185, BG222560, BG222322, AV662257, AV705443, AV729255, AV726938, AV701879, AV724819, AL047841, AV725529, BF942332, AV706584, AV759547, AV692176, AV707611, AV759063, AV758197, AA533928, AA467862, AV738071, AI698669, BE877083, AA467872, AV653804, AA467983, BE877078, AW243938, AV729339, BE875275, BE880733, BE873792, AV691890, AV722499, AA657843, BG231240, BE874492, AV728243, AV762317,</p>

					<p>BE879882, BE874475, AW128905, BE878027, BE876813, AV709039, AV721822, AV721050, AA467967, AI951338, AA468250, BG231135, AA468184, AA467920, AV742491, AV759381, AW129315, AI207666, AI287963, AV742528, BG223149, AA468920, AA468143, AA467928, AW070665, AA468361, AI634214, BF680264, AW277108, BF942071, AA468002, AV711061, AW265193, AI374100, BE876648, AA468434, AV695014, AI065052, AV744874, AI535959, C17433, AA602107, AA468467, BE877183, AI720411, AA468095, BG230797, AI022799, BE875306, BE714703, C17062, AI418640, AV705090, AV653812, BE878818, AV712864, BE874917, AI207671, AI557473, AA226446, AW872525, BG231142, C17215, AI749178, BF942136, AI114862, AI718674, AA550877, BE044950, AA729610, AA468511, AA468996, BE153318, AA652499, AI535873, AI078862, AA654934, AA468176, BG223154, AV729805, AI626072, AV754426, AI460289, AA535933, AA574306, AI081273, AA094361, AA508188, AA420618, BG231064, BE879318, AA541818, AI065071, AA468395, BE879020, AA934542, AI799863, BG231144, AA513120, C18736, AA579481, AA724186, AI581768, AA230233, AA541815, AA978151, AI926958, AI535874, AI202491, AA781850, AI749551, AA569429, AA467968, AA550867, C16929, AI796602, AI735781, AI374099, AA978154, AA724187, AI033651, AW235024, AA535848, AV739800, BE907009, AA468004, BE896904, AA527093, AW440526, AA494138, BE878632, AI817810, C18396, AA652954, AA658970, AA224951, AI524381, AA978149, AA658047, AA480424, AA527020, AA548217, AA658218, AW237889, BE168579, AA564192, AA613771, BE874118, AA573677, AI245991, AA652909, AA687521, AA654470, AI092988, AA513599, AA468310, AA876498, AA572851, AI346033, BF003082, AV693913, AA641600, AA247586, AA533509, AV745072, AA551840, AA938056, AA494234, AA548923, BE878136, BG164804, AA641087, AA094820, AA535658, AA468981, AA484530, AA664671, AA938041, AI708944, AA558338, AW264966, AW265056, AI557600, AI581596, AA558381, BE174413, AA854696, AV701496, AW235484, AF004342, X62996, AX039612, V00662, X93334, D38112, D38113, X93335, AB017116, and L27601.</p>
HWBBU75	2565	780360	1 - 2717	15 - 2731	<p>BG254332, BG258993, AW189216, AW189167, BG110226, AA621650, AI524238, AW027302, AI085248, AI187104, AJ403112, AW129561, AA287508,</p>

					AA287381, AI569606, AW137239, AI814239, BE246677, BF356218, H87267, AI831754, H87322, W15561, AA831701, AA808000, BE086121, AW137386, AA932127, H92853, BF916818, R84247, BE672348, BF869463, AI380578, BE696648, N80792, AA811717, AA922507, BF945407, AK024474, and AF159356.
HWBCN81	2566	853580	1 - 2769	15 - 2783	BF836038, AW473927, AW966643, AV761676, AU151751, N70293, AA568311, W04238, AA630476, H15679, AL121039, BG180320, AI702049, AW264548, BE328286, BG059139, AA568303, AI200051, AV761486, AW265468, AA720582, BE043339, BF530611, AW151247, AI801563, BF876261, AV759089, BE875478, AU160445, AL041924, AW957502, H60912, AA493546, BE161469, AV683406, AA845690, AI590404, AW410844, AI760835, BE794230, BE045167, BF804385, AA810158, BE154381, AI572680, AI174703, AA489390, AW192930, AW571716, AW820105, AI792623, AI732720, BF034649, AW819483, AI927275, AA298365, AI267285, AW631267, AI078409, AA557945, AI753567, AI744963, AI049643, BF740656, AU159614, AI734119, AW976008, AA584360, R23873, H90845, BE042511, AA595661, BF970107, BG231195, AL048969, BE747923, AI791856, AW389929, BF913232, AV706458, AA653513, AA568433, AI247128, AI733523, AV760466, AW072963, AA610381, AA747234, BG122782, AW963489, AL109658, AC011495, AC008014, AL162573, AC009731, AC009123, AB003151, AL050335, AC013434, AC004499, AL391097, AC020750, AC010150, AC002558, AP000692, AL031733, Z73967, AC004025, AC016637, AL138919, AL121886, AC005519, AL133548, AF061032, AC027319, AL121753, AC007381, AL109984, AC005696, AL034429, AC002054, AL356244, AC004883, AC005484, AC011510, AL022322, AL080243, AC005015, AL445248, AC004858, AC018639, AL162615, AL139095, AP000689, AL109804, AC002366, AC004840, AL078581, AC004554, AC008753, AC007664, AL133215, AL135749, AC009953, AC005907, AL138832, AC005551, AC002115, AL139099, AP001721, AF001550, AC022407, AC009194, AC008018, AL354889, AL133245, AC005218, AC009116, AC004824, AC008651, AL109797, AL157951, AL445435, AL121601, AF134726, AL121899, AC000353, AC016830, AC087089, AC006137, Z84484, AC020552, AC006141, AC011450, AC005391, AC005828, AL138756, AC006205, AL358975, AF109907, AP002015,

09550083.091201

					AL035659, AL133244, AC005736, AF241734, AL356115, AC034246, AC005365, AL117381, D83989, AF124523, AP000503, AL035458, AL021546, AD000671, AC007055, AP001724, AL133445, AL109921, AC008038, AC007390, AL163209, AC007383, AJ400877, AC004813, AL391595, AL049694, AC006538, AC007954, AF146367, AC008372, AC004134, AC009756, AL390039, AL049872, AL117348, AC007298, AC011526, AC011479, AB009422, AC004933, AC006028, AC004257, AP000036, AC010271, AF001549, AC004000, AL022577, AL035400, AC006026, AC027644, AP001435, AC011475, AC025436, AL352979, AC010077, AC027279, AL023803, AL133163, AC009721, AC004797, AC005690, AL445143, AL117380, AC007216, AC002091, AL136979, AC006285, AC025594, AL096701, AC016652, AC007899, AC011811, Z93015, Z98884, AC004477, AC008569, AC008812, AC002350, AL035413, AC005207, AC002565, AC002094, AC006070, AB001523, Z98941, AL118501, AL022318, AC005409, AC003037, AC011362, AL132987, AC008599, AC020913, AC015550, AJ009610, AC006115, AL356299, AP000704, AL034430, Z86090, AL079304, AC008088, AC003109, AC007637, AC006511, AC018719, AC002401, AP000688, AP000502, AL133230, Z77249, AC005914, AK027152, AL049794, AC006329, AC008403, AC023880, AL080317, AF003626, AL139343, AJ295844, AC005800, AC007263, AC005695, AP001060, AL356575, AC008784, AL353748, U95742, AC011497, AL121585, AF042090, AC022149, AP000517, AL135839, AL133465, AL139100, AC009362, AC004031, AC005776, AC009475, AL024498, AC018720, AC004921, AL132794, AL021154, AC005482, AC010092, AL445532, AC006211, AC009510, AC007676, AP000350, AC008033, AL137060, AC005209, and AC010328.
HWBCP16	2567	647378	1 - 611	15 - 625	AW243699, U73479, AL135909, and AC069270.
HWBCX93	2568	853418	1 - 817	15 - 831	AW957665, AW075995, AW576203, BF339246, BF308177, BE881903, AI051657, BG059809, AW058408, BF689190, BE868083, W56301, BE301190, R05745, AA102223, AI091799, D61582, R01123, AA102222, AA375163, AI752483, AA376452, AW084324, AW275432, AI439525, AA661583, AW238137, BF812696, AW151541, AI828721, BG258103, BF309372, AI039257, AA631915, AW162314, AI049999, BF679568, AI702049, BG059139, AW243817,

09950087-09101

				<p> AI538404, BE244308, AW473160, AW265468, AW162332, AI860423, AW021399, AW020094, BF214695, AA112864, BG180320, AI753131, BE676856, AA728954, AI355246, BE273825, BE176819, AI554399, BF940118, AI308529, AA807704, AL121039, AI270280, AI559304, AA507499, AW148821, AA729004, BG027735, AI819419, BG152746, AA569220, AA297776, AW510403, AV653126, AI884404, BF901147, AI338575, AI860648, AW008217, AW410844, AW327673, T34066, AI031759, AV732057, AV753446, AW151848, AA197089, BE206375, AA601674, BE140949, AI636422, AI064968, AW576388, AV742390, AC006483, AK024787, AC009756, AC004659, AC005519, AL078611, AC005531, AK000385, AC010271, AL354836, AK027150, AC002996, AC011444, AC005358, AC005052, AL158040, AC016995, AP000280, AP000039, AP000107, AL139824, AL023879, AC004966, AC004552, AC010422, AL121675, AC011479, AL109923, AL133458, AC002543, AL139184, AL035086, AC018663, AC011526, AC003108, AL117382, AF195658, AL121897, AC007421, AC005280, Z97183, AL031680, AB017653, AL049843, AC008738, AC007240, AC004985, AP000256, AF207550, AC000353, AL035659, AL391259, U62317, AL035683, AL358777, AC010789, AC006449, Z83851, AL357992, AL050335, AL391114, AP001715, AP000098, AL096701, AC004851, AC008392, AC004974, AC003007, AC018633, AC004673, AC005911, AL133229, AC024561, AC006151, AL353748, AL138733, Z85986, AC005041, AL117337, AP000557, AL159168, AC008736, AL096791, AC004066, AF258545, AC008372, AL049759, AC009086, AL355838, AC009469, AC004752, AL035249, AC008848, AL049780, AC005695, AX039602, AC008044, AC020913, AC003957, AL136362, Z81364, AF196779, AL136228, AP000555, AC020904, AC009362, AF293359, AC005923, AC011480, AC008521, AC006530, AL109799, AL138717, AC005940, AC004381, AL109935, AC016025, AJ011930, AC011495, AL162505, AL163300, AL031058, AC011895, AL121890, Z95114, AC007383, AL034369, Z97632, AC002546, AC018673, AC005696, AL008582, Z98047, AL117334, AL138756, AL022316, AC008784, AF107885, AC008482, AC009399, AP001712, AC003100, AC005484, Z49236, AL080250, AF217413, AL157789, AL139120, AC010458, AC008635, U07561, AL133545, AL355385, AC006064, AC011287, AL121586, </p>
--	--	--	--	---

					AC002110, AL353678, AL034548, AC004762, AL158830, AL034417, AC005570, AC010326, AL050404, AC006345, AC006324, AL389886, AC005907, AC007957, AC004821, AC002105, AP001711, AL020993, AC007845, AL031282, AC004263, Z98742, AC009408, AL049761, Z83840, L78810, AL022165, AC004967, AC005841, AC002404, AC006543, AC018719, AC069248, U91327, AC004847, AC015651, AL035705, AC005091, AL133163, AC007676, AC004975, AC002563, AL138976, AC005365, AC005288, AL353812, AL008710, AC005412, AC005399, AC004233, AL121872, AC007536, AC022311, AC005182, AL138787, AC018797, AL133448, AC004643, AC002060, AL137230, AF317635, AC008688, AC005527, AL121967, Z98752, AC003963, AC011453, AL021453, AC005602, AL109758, AC002314, AL136304, AL162615, AC004900, AC000379, AC002310, AL353802, AL132713, AL035460, AL161670, Z98948, AP000346, AC006315, AL353807, AC006483, AC006483, AC006483, and AC073481.
HWBEF34	2569	799430	1 - 1454	15 - 1468	AV717583, AW337428, BF871047, AC010191, AC010191, and AC010191.
HWFB23	2570	853319	1 - 1397	15 - 1411	AV719367, AV719824, AA809261, BE671612, AA526520, AW514791, AI299369, AI653980, AI803977, AI377662, AW058300, AI281374, N64512, AI355041, AI038559, AW118603, AI246272, H02873, BE870177, AI970788, AI884393, AI702548, AA302790, AA633580, BE966154, AI245763, N22380, AA452590, AA502655, D59329, AV740627, H50615, AV744231, AV740287, AA302791, BF094920, N75621, AI205785, AV736353, BF222604, BF340004, BF111206, BF672836, BF102947, and H03768.
HWFB140	2571	826791	1 - 861	15 - 875	AW004885, AI765130, BF924471, AI381820, AA302642, AI824365, AP000785, AP000785, AP002340, and AP002340.
HWHGV77	2572	788555	1 - 1012	15 - 1026	AF005081, and AL353779.
HWHGW0 9	2573	702032	1 - 582	15 - 596	AW189948, N58823, N40196, AI806808, H78365, AI419770, BE874627, AI580662, AV713790, AA809439, AI186697, AI310126, AI261311, AA809455, AA019739, AA833673, AA971980, AA532665, AW264561, AA838000, AI800634, AA610052, BF000801, AI361312, AA904689, AI189791, AA970480, AW969490, AA088862, AA887119, AA575958, AI784506, BE222956, AI188971, AA906201, AI684587, AW009062, AA018103, AA059058, AI538082, AI751250, AI654089, AV741671, AW954079, AI083552, AA054140, AI933491, T47520, H86494, BE621190, H66353, AA935534, AA469072, AA634291, T50287, BF335535, AW750631,

					AW514695, AA887381, AW750632, H86061, AI268976, R24685, AV715976, AW614415, AV713864, AV715267, AV763925, BF968896, AW440198, BF971170, H66403, BF920846, BE765444, AA935511, BF128924, AA886637, BF036411, BF376603, AI971218, H78462, BG170547, AW883576, BE410537, Z19788, W01156, AV758546, AA016261, AF182041, AX017986, AF182414, AF113127, AL117550, AF151877, AF161526, A74434, AC012264, and AC012264.
HWHHA21	2574	638966	1 - 681	15 - 695	AW300079, W69650, AW291010, AF086289, and AF005080.
HWHPU44	2575	778090	1 - 857	15 - 871	AW276035, AI052543, AI269844, F26678, and BE705973.
HWHRC51	2576	853421	1 - 829	15 - 843	AV699657, AI672389, AU146627, BF475506, AI767322, AW953696, AI828672, AW953697, AU155457, AA553619, AI866742, AA808077, AI367812, AW022647, AW022875, AW957432, AA602347, BE182281, AW265509, AA640875, AW026601, C18790, AV721217, AA224926, AA651901, AA224835, AI581427, AU155347, BF883894, AA503350, AW612118, AW020289, C06444, AI023252, AW020357, R52941, BG223355, R53723, AA631489, AA353335, AI873340, H86877, AA632732, R79085, AA825730, AI784551, AA432003, AA936640, R78782, AI610979, AI832137, AA618185, AA808914, AA640349, AA947260, D63288, H86813, AA731390, AA494280, AA349142, BE676991, BE000573, BF061245, AA613341, AA302655, AV659756, R31243, AW265313, AV652132, AA563943, AV652191, AA381288, BF980434, AW265736, BF576873, BF928764, AI936025, AA551180, AV735034, BE972238, N99348, AA876590, BE092445, BF843805, BF693632, AA366513, AW026600, BF325835, BF325829, AV689224, BF325822, BF325820, BF325838, AC000353, AF001893, AC004000, L78810, Z75889, AB014084, AC004207, AP000515, AL161415, Z98255, U95741, Z93403, AL138914, AC000353, AC000353, AC000353, AP000441, and AP000441.
HWLAT50	2577	840753	1 - 2959	15 - 2973	AU124961, AU125676, AU117401, AU124066, AU142794, AU120366, AU124113, AU120456, BE877005, BE740295, AU141338, BG167721, AV714367, AV714673, AV686436, AU142175, BE304981, BE881100, BE250110, AU134181, BF025751, BE895076, BE622248, BE409732, BF530437, BE620787, BE745726, BE907256, BE733942, BG177013, BF683520, BG260709, BE272337, BF791820, BE740837, BE871431, BG249189, BE513215, BG104424, BE273992, AV717006, BG181090, BE747751, BE882158, BG117384, BE562622, AV705499, BE274707, BE742907, BE410659, BF338712,

				BE390178, AA618435, BE252543, BE304944, BF849703, BE387435, BE263772, BG252449, AU140315, AW393155, BG248941, BE271904, BE302912, AW401763, BF340560, BE271435, AA218934, BE867601, AI302708, BE390559, BE249815, BE391942, AA135472, BE697398, BE620245, AU149287, AI620213, AU131831, BF345543, AW361950, AU150881, BF733021, BF724789, BE543574, BF326587, AI719608, BE272764, AA488941, AA809494, BF513135, AV707470, BG170291, BE410631, AU158923, BE872556, AA772223, BE293766, AA868284, AU148718, H38068, N22759, BE293723, BE293570, BE747763, AA772307, AA863295, BG231986, BE870639, AA463604, AU155275, AU152149, AU159524, AI089946, AU158387, AA524148, N56943, BE709561, AU158420, AI800921, AI188323, AA524950, AW302264, BG119202, AI696916, W79668, AI093254, AU144238, AW002036, AA847756, AA224477, AW383920, AU158506, AU152524, N20184, R12134, BE697395, AA595472, AI857582, AA831011, AA135543, AI829858, AA130928, AA196636, BE389048, AU146453, AA565671, AA428061, AA777674, AA149667, AA461608, R18682, BE617254, AA219705, AW170341, AI313271, H50233, AW469372, AA216524, AW753234, AI672766, AW886897, BF758190, BF680915, AI288758, BE387891, BF766631, BE675067, N42998, BE789704, AV708618, BF765172, AA527313, AA171564, T59935, AW937851, AI557334, AA171885, AI051796, AI220979, BF740181, AI087122, W75944, AI199174, AI912535, AA639597, AI290537, AI299852, AL042170, N29086, N72103, AA861852, AA470662, AA599225, AI246626, AI356062, AA938598, BE390486, AW516050, AV744699, AI689428, BF894348, AA226950, AU153894, AA693475, R15067, AA244053, AA831013, W74130, H49287, AA130873, BF801711, AA714302, AW050958, AA829809, AW364166, H80481, AV702735, AA470541, BF846600, AI032131, AI128671, AA643976, AW576483, AA448218, H24857, AA229560, H43549, AA723067, R13592, AA513255, AA219772, H49286, AI038225, AA974538, T94807, H24856, AI933143, AI262723, H38069, AW079258, BE706867, AA935861, AI092200, AI302474, AA868497, M78751, R39888, AA011461, AV742747, D63486, X07997, AK027136, AX020124, AL137523, AK026541, I00734, AK027144, AK025113, AJ001388, AK025798, AK025084, E00617, E00717, E00778, AF287051, AB009690, AK026649, AF177401, AF076464, AK025860, AF116644, AF183393, AF113694,
--	--	--	--	--

				<p> AB047930, AJ003118, AY004290, U76419, U89295, X72889, I89947, AK025092, U31501, S61953, I48978, AF017437, AF140224, A91160, AL110269, A08910, A91162, AL110159, A08909, A12297, U42766, AK000653, AL137627, A08908, AL162008, AF116609, AL110225, AF022813, E07108, AL137480, AF090900, AK025857, I32738, AK026600, AL035458, Y10080, AF067728, AX042059, AF026124, AL389935, AF139986, AR038854, AK024546, AK026462, AL049426, AP001434, AB016226, AF111112, Y10655, AK027116, X66871, AK026057, AJ299431, AB025103, S82852, AL122093, AX015425, AK025517, AF201468, AC007458, AF100781, AL109672, A08913, AF262032, AL389982, AL133075, AF188698, AF044221, AK026038, AL359596, AF061573, AL137479, AX025493, A08456, AF119336, AK027121, X70685, I46765, A49139, S77771, AK024944, AL442082, AL117394, AP000020, AL080129, AF119899, AK026648, AF109683, U02475, AL355719, E12579, AK025209, AL137459, AK000421, M27260, AK027188, AK026927, AL080156, AL096720, A18777, AK025119, I18355, AL050277, AR085690, AL122045, I30339, I30334, L13297, AK024533, A77033, A77035, AK026642, AF069506, AB047941, AK027103, AF031147, AF111849, AX015416, AL080127, AL137533, AL353956, AK024524, AR083266, AF097996, AF079763, AK000655, AK027204, AB048975, AL122098, AL137705, AL080148, AX019230, AF008439, AL157464, AL110222, AF116631, AC006197, AK026626, AR020905, AF130110, AF130066, AL110196, AF217998, AB048913, I33392, AF118090, AY007100, AF113222, AF017790, AB007812, AB050431, I09499, AF130077, AL133014, U35846, AL355701, AF207829, S54890, AF155148, AK026647, AL162083, S36676, AF130055, AF159141, AF087943, AK000432, AB050510, AL137271, Z72491, AK027173, AK000074, AL355713, AF153205, AK027164, AC002464, AF000301, AF061795, AK025015, AL117457, AL050149, AF151685, AF111847, AL133568, AX019229, AL137292, Z97214, A17115, A18079, AX040958, AL121828, AL080154, AB044547, AF130059, AL157433, AL161802, AK026593, AL137267, AF314091, AK000418, AF142672, AX015403, A08912, AL137574, AF119857, AF130075, AL133608, AL137273, AF130102, A08911, I48979, AF119875, AK026534, I89931, AK000614, X76228, AB047248, AL080159, AL110296, AK024747, AK000618, AK000250, AK000197, AK027260, AL137485, J05277, AF106862, T60005, T94060, R13984, R40044, </p>
--	--	--	--	---

					R42065, R42065, R40044, H26768, H26878, H27071, H27083, H27280, H52356, H81400, N35033, N45042, W00326, W70202, W74251, W96207, W96300, AA011404, AA149732, AA165372, AA196918, AA224416, AA226902, AA227012, AA227198, AA460317, AA470998, AA481296, AA548195, AA592919, AA807641, AA827604, AA831396, AA862089, AA876627, AA876966, AA879161, AA927418, AA970339, AA991176, N55983, AA650180, AA211249, AA219650, AA448361, AA487921, AA488036, AA678876, AA778272, Z25362, Z42751, F02226, F00374, AC063942, AC069214, AC069214, AC069234, and Z97197.
HWLBO67	2578	834315	1 - 522	15 - 536	AC011739, AC011739, and AC011739.
HWLGP26	2579	834770	1 - 1884	15 - 1898	AL529987, AV699741, BF530871, AW207742, AI638032, BE540656, AI469103, AW575118, BE503442, AI652512, AA928729, AW592532, AI419960, AI057140, BF102699, AI365238, AI392687, N49757, AA832314, BF819685, AI025113, AI767982, AA766124, AU151363, AW402473, AI654109, AW070683, AI738949, AA815032, BG231904, AW578625, AI365240, AV688777, AA769270, AW613566, AA605050, AI459543, AI885560, AW361738, AV689923, AA046663, BE242932, BF895261, AA507121, AA046576, AI627311, AW296010, AA814924, BG253328, BE789823, BE243489, AA298793, AI458707, AI208523, AA694217, BE247085, AW137352, AV689313, AA077726, BE241524, AL523598, AV685517, BE241950, AV697781, AV691130, AV688018, AV697631, N52685, AW182301, AA910584, AA126044, AW390687, AL529988, AF176097, AK023002, AJ131891, and I25644.
HWLHO31	2580	837480	1 - 1687	15 - 1701	BF727430, BE379599, AW238333, BE380022, BF091252, BE832321, BF091254, AI033829, BF217856, W86466, W73267, BE728037, BF380085, AA346592, W86465, AW810953, AF279865, and AB001430.
HWLIL31	2581	793698	1 - 773	15 - 787	AI139990, AI961465, AA724414, AA582139, AI922007, AI635251, AI675404, AA626226, AI081214, BF084927, BF084935, AL134524, AW861944, AW858525, AW973214, AW858455, AW858526, AW970927, AW970107, AW972849, AW968196, AW979151, AW975212, AW979264, AW969785, AW970959, AW975037, AW975961, AW970101, AW969825, AW973185, AW973166, AW976023, AW971403, AW972292, AW975032, AW973957, AW975434, AW975959, AW969655, AW969626, AW975031, AW969780, AW969848, AW970025, AW979212, AW973219, AW969631,

				<p> AW975019, AW972845, AW969911, AW979204, AW975025, AW969673, AW974101, AW975015, AW975027, AW975154, AW979176, AW972377, AW974971, AW969748, AW969791, AW975876, AW975632, AW969861, AW970540, AW979238, AW974338, AW969988, AW969852, AW969778, AW979098, AW970936, AW973978, AW970969, AW975692, AW970084, AW975921, AW968212, AW970050, AW979127, AW969885, AW976024, AW975971, AW975244, AW974658, AW971968, AW969643, AW975132, AW970053, AW976517, AW979294, AW979169, AW969740, AW972880, AW979090, AW979220, AW973717, AW968207, AW979219, AW969839, AW604723, AW969625, AW970942, AW979113, AW971732, AW968200, AW973242, AW979134, AW970079, AW975105, AW969628, AW975952, AW975020, AW974975, AW972296, AW969680, AW976511, AW975981, AW974786, AW979129, AW972649, AW976031, AW970889, AW979002, AW974801, AW972670, AW974823, AW969816, AW975965, AW979232, AW877209, AW804686, AW968204, AW970070, AW969793, AW969634, AW979142, AW974785, AW974964, AW979037, AW975231, AW976000, AW970094, AW969857, AW975585, AW975221, AW969633, AW975635, AW973819, AW975254, AW979211, AW970108, AW979053, AW975628, AW971305, AW142134, AW973146, AW975225, AW979133, AW974660, AW970113, AW975084, AW975596, AW971378, AW973397, AW972939, AW972884, AW972695, AW973805, AW975002, AW971091, AW975990, AW972719, AW975942, AW969653, AW976515, AW968347, AW974089, AW979165, AW972380, AW975927, AW976510, AW970076, AW973230, AW971954, AW972680, AW975975, AW971326, AW969884, AW968213, AW972943, AW970105, AW969759, AW979083, AW976982, AW970587, AW979175, AW975646, AW973986, AW979064, AW969892, AW975938, AW975016, AW969823, AW975966, AW975643, AW969734, AW970921, AW969864, AW969766, AW970073, AW975982, AW979116, AW972806, AW969645, AW969664, AW973987, AW972706, AW973164, AW972821, AW970097, AW973824, AW971964, AW975904, AW973726, AW974379, </p>
--	--	--	--	---

					AW969862, AW969752, AW972702, AW975227, AW969809, AW969827, AW973718, AW973967, AW975933, AW969826, AW973734, AW973207, AW972882, AW973104, AW975028, AW969782, AW970868, AW972868, AW972803, AW969798, AW973985, AR066494, AX030435, AR023813, AR096545, A93931, A93923, AR069079, AR060234, D17247, AX046357, A93916, A81671, AJ276251, A85203, AJ276253, AJ276255, AJ276256, AJ279014, AJ251859, AL133053, AB026436, AR054110, AJ276254, and AL122101.
HWLJN08	2582	800626	1 - 1016	15 - 1030	BE300001, BF309344, BE300104, BF111874, BF968540, BE790918, AI076943, BE905154, BG166354, AA447420, AW675708, AW732847, BE295220, AI800955, AI204057, AI083659, AI248992, AI184659, BF204818, AI866483, AV752565, AA677077, AI149422, AI160393, AW339819, AI362801, AA442134, AA918843, AA035216, AA865237, AI082597, AI591234, BG164319, AW074181, BE048273, AA983292, AA443434, AA148610, AI497928, AA971924, AW103005, AI017651, AA807948, AW732861, AA909975, W68840, AA443301, AW025275, AI299836, F24136, N80272, AI056188, AI743407, AA442133, AI817457, AA563798, BF305961, F33099, AA491148, F30260, N94975, AA599349, AA062929, T78326, AA447419, R28029, AI936185, R96685, AI913023, AA854576, R28145, F18276, AW664849, W67311, AA312862, F25535, BF767154, T78708, AA148609, AW879578, W07400, AL119324, AW877209, AL134524, AL119457, AL119511, AL119399, AL042544, AL043152, AL043168, AI431323, AW858522, AL042382, AW970048, AL135012, AL043089, AW858525, AW858526, AL043321, AL042488, AL045891, AW861944, AI432644, AL042898, AI431307, AI431316, AL046356, AL042515, AL043091, AL079794, AL045327, AL047611, AI432666, AL041862, AL047163, AL042853, AL042729, AL042787, AL043239, BE883591, AL040207, BF525578, AI431238, AL047675, BG166654, AL038878, AL042655, AI432653, BG110517, BG113493, BG029667, AI567971, BG252929, AI371243, BE897632, BG164558, BE891834, AL042533, AW858455, AI866786, AL045328, AI433157, AI648567, AI690946, AI432656, AW151136, AI366910, AI539771, BG257535, AI537677, AI494201, AI500659, AI866465, AI815232, AI801325, AI500523, AI538850, BF796402, AI887775, AI582932, BG027628, BF338002, AI590043, AI923989, AI872423, AI284517, AI500706, AI491776, AI445237, AI289791, AW151138, AI440238, AI889189, AI521560, AI500662, AW172723, AI582912, AI539800,

09500566 002160

				<p> AI284509, AI538885, AI889168, AI440263, AI927233, AI866573, AI633493, AI434256, AL039390, BF795712, AI866469, BE886728, AL037081, AI434242, AI805769, AI888661, AI500714, AI284513, AI888118, AI285439, AI859991, AI436429, AI355779, AI889147, AI623736, AI581033, AI371228, AI491710, AI440252, AI890907, AI860003, AI610557, AI242736, AI828574, AI887499, AI539781, AI539707, AI702065, AI885949, AI285419, AI559957, AW089557, AI521571, AI469775, AI866581, AR014386, AR014390, AR014391, AR014389, AB033884, AR014388, AB026675, AR014387, U77594, AX030435, AL133049, AR066494, L10353, AL133053, AL122101, AJ251859, AX046357, Y11587, AX040974, AL133074, AX040958, X66871, AF113222, AK026613, AF044323, AF056191, I17544, AK025375, AB048975, AL137665, AF032666, AJ005690, M92439, E06743, AL137574, AK025465, I09499, AL389935, AK026600, X98066, I89947, I48978, AF047716, AF116688, AR034821, AK024747, AB048913, AF132730, AF151109, AK026642, AF019249, A08913, Z97214, AF002985, AL122049, AL122098, A08912, AK000690, AL162003, AK026164, AL137711, AF026008, AF274348, AF274347, AF106657, X66975, AL137480, AB049848, AF155148, AL122110, AL049324, AF130055, AF116609, Z13966, AK025431, AB050431, AL389947, AL353940, AF130110, AL137521, AL157433, S76508, AR060234, AF065135, AF079763, AK025350, AK026630, S69510, A08910, A08909, X76228, AK027081, I33392, AL122100, AJ003118, AF090901, AL133560, A57389, A08908, AL117626, AF218006, AF200464, S68736, AF139986, A18777, AR070212, A08916, AF026124, S83440, X93495, AK026532, AR038854, AK027104, AC010128, AF078844, AF118094, AK026057, AL133070, AX019230, AK026784, AK025541, AK025524, I89931, Y16258, Y16257, E02756, Y16256, AF114818, AK026746, E01314, AL080127, S77771, AL133665, AL050092, AL110218, I00734, A08911, AF111112, AF130104, AR087170, A86558, AK027096, AK026086, AF028823, X79812, AF089818, AK000137, AL137271, AF111851, AL080110, AF116650, E01812, AF111849, AK027164, AF094480, AF116602, U30290, AF090886, AK000310, E00617, E00717, E00778, AL117440, AL133014, AF185576, AL359620, S79832, AL355701, AJ006417, AX019229, AF102578, AF022363, AF008439, A08907, A65341, AF217966, AB052191, AB049880, AL162062, AK026556, I68732, I89934, AK026534, AK026504, U42031, AL137478, AK026571, AF130092, AF017790, AB044390, U87620, </p>
--	--	--	--	---

					AK025015, AL050393, AF106862, AF114168, AF271350, AL359941, X66862, AK026480, U92068, AL117587, U96683, AF260566, AL137533, AL096751, U35846, E02221, AK027136, AF116631, AL389939, E12747, AC024247, AK026865, AF119883, AF113694, AL162083, AK024533, A65340, AF067728, AX042059, AL389978, AK026528, AB048919, AL359600, E12806, AK025092, X52128, AK000250, AF188698, AL133606, AF000145, U49908, AF116682, AK026462, AK000450, AB019565, I48979, AF119337, AL049339, AB047248, AK027173, AF119909, AK025209, AL359601, AF205861, AL137256, AL133410, AL133410, and AL133410.
HWLRE03	2583	831198	1 - 2756	15 - 2770	AL534216, AU124415, AL534215, AU124249, BE793898, AU132166, BE792345, BG111555, AV688850, BG171107, BG028492, BF969425, AI801432, BG108221, BF338327, BF663592, BE619881, BF746890, AW965641, BG122860, BF665525, BE788680, BE003409, BE905238, AU137278, AW151135, AU128897, BE884319, AW262132, BE384410, BF742797, AV710041, AW769142, AW517359, AW338005, AW373588, AW468740, AW024843, BE619266, AU128495, AU151118, BE619730, BF690080, AI937477, BF331576, AI660064, BE963856, BG114091, AA536151, AU153751, AU152127, BE716089, BE716103, BE905313, AI624934, BG122469, BF591733, AA614338, BE716050, BF212934, BF436427, BE091534, AW374908, BF690005, BE716084, BE619170, AW024853, AI056781, AW579072, AI018651, AA316213, AI885775, AA701904, BF769669, BE273759, AW072631, AW438832, BF090279, AA718962, AW068084, AA693953, BF997218, AI139494, AI660421, BF997182, BF888379, BF770032, AA664290, BE716120, BF770040, BF918375, AA522683, T66125, BF695272, AA862983, AA428398, AL046516, AA972193, AA748129, AA504220, BF381340, AW136037, AI348174, BF769655, AA442961, AV684332, AA459911, AI873944, BF930962, AA497104, BF923552, BE716094, W87301, AA161216, AI372983, BF770037, AI828291, AA045068, AI056779, AA100277, AW068172, AW516076, AI244381, AI286345, AU124139, BF770006, N63176, BF985698, BF923559, AA350408, BF888406, BF769993, AA081089, AI352530, BF091975, BF769671, R52005, AA431627, AU158647, AA100326, AA368310, AI125481, BE004008, BF349060, AA496940, AA578753, BE812515, AA350409, AW410891, AW190017, BF768918, BF155861, AA398870, BF767919,

					<p>BF764594, BF771908, AA923654, R52006, BF357204, AA435899, R71767, BE871607, AA393437, BE693936, AI032221, R71806, BE076619, AA563608, AA081247, BF155867, T35217, AI357799, AW884241, R69972, BF764597, BE167977, AA985512, AW749731, AA431341, BE812695, BF769721, BF903475, T53979, AI379785, AI445967, BG024346, T80019, AV688441, AA356434, AA922349, AW965639, BE731858, BF763097, R38747, AA578094, R48216, AI355212, BE693932, AA159119, AW375234, BE735158, AA045450, AA367245, T05136, BF088681, AW375279, C21399, AW241404, BE964246, AA143624, BF907203, BF830391, AA625473, AW796532, T80321, AI452574, AA353270, BF377656, R15479, AA983169, BF889342, R48217, BE148770, R69973, BF475528, AI024745, AA969907, BE841884, AA383452, AA962116, BF091778, AW351601, AA292992, W87438, T09121, AI907921, BE019213, AW600910, T07116, BF368764, AA370080, AW403245, AW263361, BF998371, BE810503, N59506, BE836987, BF971195, BE715793, AK025149, AF055010, AL049972, AB007931, AK022924, AK023826, AK022573, AK001233, AK026543, AL137127, A75151, T53282, R38966, AA421365, AA426283, AL137127, AL137127, and AL137127.</p>
HWTAM38	2584	602274	1 - 584	15 - 598	<p>BF516514, N94326, AA827238, AA921725, AI671598, AA766820, AW188348, AI692659, AA664123, BE219817, AI824114, AI346940, AA743541, AW469089, AA858171, AA507413, AA857721, AW594160, AI765481, BE720623, AI269812, Z38498, T39272, AW959868, AV701728, AW960977, AV703989, AW951562, AW954235, AW950395, AW954006, AV655280, AV661704, AW952168, AW950443, AW954001, AW952323, AW952196, AV660258, AV726590, AW959823, AV661286, AW965720, AW963446, AW952167, AW951536, AV697880, AW953619, AW962978, AW958295, AW966055, AW952046, AW959830, AV706076, AW959986, AW955161, AW960629, AV705416, AW949731, AW958796, AV705684, AW950241, AW949517, AW949441, AV728652, AW960456, AW962392, AW963949, AW954211, AV709356, AW954237, AW965557, AW966539, AW961597, AW952419, AW965839, AV725393, AW951772, AV729473, AW957997, AV725255, AW965759, AV659322, AV727618, AV704234, AV709880, AW964860, AV725386, AV706223, AW952301, AW952592, AW960143,</p>

				<p> AV728270, AV660096, AV709580, AW959521, AW963354, AW962924, AW954386, AV658784, AW962934, AW966650, AW950635, AV647654, AW950276, AV652001, AV654035, AV707933, AV702164, AV725618, AW960579, AW964074, AV706279, AV707401, AV653353, AW951875, AV707171, AW952005, AW951768, AW954344, AW955139, AV701774, AV708786, AW961240, AV707510, AW952011, AW950006, AV707654, AV692972, AV704771, AW956291, AV725617, AW953992, AV702728, AV691615, AW955704, AV704376, AV709886, AV694789, AW955749, AW963601, AW951445, AV727822, AW957779, AW958896, AV651694, AV706183, AW949330, AV705550, AW960685, AV707822, AV701873, AW960276, AW952031, AV727126, AW951526, AW963700, AV727459, AV651897, AV728715, AW950211, AV705147, AW958620, AW954003, AW954310, AV653845, AW967329, AW953809, AW964112, AV658362, AW960655, AW954427, AV709555, AW952582, AW958849, AW950632, AV709935, AW952368, AW959895, AV729220, AW959870, AV702071, AW958163, AW949454, AW950254, AV706165, AV705504, AW962397, AV725582, AV725577, AW964994, AW956640, AW951728, AW949327, AV706527, AV684604, AW956773, AW956762, AW964800, AV645545, AV685688, AV689800, AV727238, AV725709, AV704097, AW960406, AW960067, AV701914, AW959806, AW949734, AW950256, AV703515, AV728404, AV726665, AV727807, AV728355, AV655577, AV726948, AW960545, AV705014, AV702790, AV702954, AV709256, AW959858, AV706827, AW955397, AV706162, AV709596, AW949942, AW954011, AW950520, AV658797, AV702574, AV707155, AV707770, AV654686, AW963234, AV684762, AV725441, AW967340, AV701874, AW967341, AV701880, AV728436, AV706501, AW960600, AW951549, AV703035, AV705655, AV725031, AL022067, AX008042, U08185, AX020190, AR062871, A25909, AR088705, AR038855, AX047062, AR070327, Y09813, AX047063, AX009343, AX009341, AX009337, AX009335, AX009345, AX009342, AX009344, AX009340, AX009339, AX009336, AX009338, AX009334, Z32836, AR079804, </p>
--	--	--	--	---

0995083 09201

					AR091393, AX009487, AX026824, AX026823, A85395, A70872, A85476, X81969, AJ244005, Y16359, AX001322, D50010, AJ244004, AX028305, AX021518, A20702, AR062872, AR062873, A20700, AX006825, AX006826, AX006822, D78345, AX001082, AF144029, A43189, A43188, AX006816, AR037157, D13509, AR017907, AJ244003, A70869, AX042372, AX008555, AX047064, A98420, A98423, A98432, A98436, A98417, A98427, U94592, AX043922, A98767, A93963, A93964, I63120, AR038762, AJ244007, AR091537, I19525, AR008429, AX027636, I44681, X83865, A84772, A84776, A84773, A84775, A84774, AR054109, A86792, AX006821, AR067731, AR067732, A58522, A91750, AX035462, AF006072, AR054723, X55486, I13349, I05558, A18053, AX011024, A93016, AJ276254, A60212, A60209, A60210, A60211, AJ276256, X82786, AB025273, A58524, AR096545, AX009712, AX003207, A58523, A35537, A35536, A02136, A04664, A02135, A04663, I84553, A82595, A82593, I84554, AX020191, Z30183, AR069650, E03627, I48927, A90655, D13316, E17098, AR093392, AX032992, A02712, AR073846, A77094, A77095, A22738, AR080470, AR077142, A81878, AX033488, AX033489, AX033490, A95051, I06859, AR095492, A18050, AX012337, A23334, A75888, I70384, AX033474, AX033486, AX033487, A64973, A60111, A23633, AR007512, AX032993, AX006823, AR031566, I00682, A11245, A11624, A11623, E00609, E13740, A11178, E01007, AX035980, A10361, I62368, A62298, AR043601, AR050070, AX032758, I15718, M28262, AJ270780, AF149828, AB043473, AR063812, I15717, I03331, S60422, I01995, E12615, A02710, AX018504, AR035193, A92133, E14304, A07700, AX027811, AX027809, A13393, A13392, AR031488, I13521, I52048, A27396, AX027813, AX030369, AX030368, AR069374, I25027, AR027100, I49890, I44531, I28266, AR069426, AR069375, AX042377, AR083151, AX027812, AX027810, AR091518, AX027816, AR072535, AX042375, AX027817, AX027818, AX027814, AR085082, AR085089, AR085091, I21869, I26929, AX042373, I44515, I26928, I26930, I26927, A91965, I44516, A70040, AR069417, AR085079, E16678, A82653, AR085083, I08051, AR093385, E16636, AL022067, AL358952, and AL358952.
HWTAW58	2585	793424	1 - 2292	15 - 2306	AU121246, AW965213, BE888880, BF969472, BE018770, BE538301, AW298151, BF439504, N63911, AI150685, AW276258, BG166820, AW084531,

					AU159590, AI798123, AA420972, BF108922, AA278448, AW440506, AL523771, AA397741, BF447725, AU151306, AW029430, AA844714, BF593900, AA309019, AI814715, AI375602, AI671082, BE675308, AW662265, AL121559, AI129271, AI264636, BE302333, AU147100, AU153824, AA279064, BF111183, BF338102, AI800930, BF437612, BE244434, AI128524, AA770328, AA604175, BE931113, W57888, AI459763, AI567988, AW610188, AA767365, AA741015, BE047468, AI300454, AI097191, AU159401, AL523770, AI268772, N98283, BF062217, AI634054, AI962365, AV764131, AI808419, AI242503, AV740486, W57889, W02788, AI282961, BE540122, AW878373, AA694446, AW449948, AW007119, W32688, BF240008, AA192930, BF438790, BF436246, W20437, AA527344, BE773309, AI002864, BG222067, H65457, AI279193, AW878386, BF740328, AI701745, AU156804, AI810271, AI005623, W30928, BE150887, H64859, AI932793, T06456, AW072955, BF692847, AW591747, T59893, H38505, BF438920, AI208375, H38460, BE246312, AA632819, R53834, AA771712, AA865975, BF692195, AI200641, T63806, T90944, R92802, AA830683, BE748241, BE243855, AI611170, AA508455, AI423743, AA336568, AI471407, T59984, BF222578, AW079714, BE246311, BF886717, BF818527, AI674229, AA345918, AW371086, AW381556, AA035196, AW380927, BE888710, BF369615, BF369640, AA035195, H26754, H26853, T85830, BE889189, N88827, AW795942, BE177371, N55947, AW795932, AW438919, AA813133, AW795939, AW795936, AW795919, AA723772, N55247, T63770, AW795831, AW795894, AW795920, BF825813, BG057876, BE774892, BF085157, AW798495, BF378604, BE175244, BE174881, BE175315, AL389982, AK001920, AF264779, AK023950, AJ236925, AF264780, and AP001788.
HWTBB42	2586	603089	1 - 77	15 - 91	AC004985, AC004985, and AC004985.
HWTBC75	2587	855978	1 - 685	15 - 699	AI972322, AA193309, AA528241, AA527684, AW235706, BF439445, AI436191, AA290870, AA890512, AA767452, AW299850, AW137415, AW089851, AI056055, AI023887, AI090024, AA041239, AI418205, AI808670, AI580941, AA283713, AI400431, BF110940, AI130923, AA040801, AI056706, AW295892, BF002797, AA917434, AI078176, AA218890, BF063099, AA683232, AA706411, AA479888, AA960854, AA693619, AA846649, AA960792, AI222234, AI084465, BF446937, AI367258, BE045520, N52967, AI269900, W15394, BE766239, AI632800, AA766379, R50797, BF915791, AI826582, BE765484, AI955173, AA216603,

					W58520, T03894, H51737, H09347, AA760918, BG109345, AI669191, AW137802, Z28597, Z39660, AI932942, AI910510, AA412174, H90518, F26651, AI349313, Z28661, AW082288, AW752307, H73642, R40788, AW235792, AA954763, BF475305, AW576920, AW351827, AI301633, Z24941, Z28596, AI800039, AA194263, AI953061, AA621533, BE787793, BE669753, AA425331, AV660952, BE166229, AW888630, BF589088, Z20686, N45966, N44279, AI742337, AI696486, AW947005, AW947119, BF575228, BF436010, BG106521, BE827833, H09407, AW888585, D20547, AW139334, AA931678, AW950094, AL157499, AL050211, AC004148, A75307, AC004148, and AC004148.
HWTBI25	2588	637736	1 - 324	15 - 338	AI079544, BG057124, T16588, AA513571, AI659917, AA054120, AW971303, BF095888, AC066603, and AC066603.
HWTBL86	2589	791720	1 - 2775	15 - 2789	BG026313, BE867570, BE379059, BE252956, BE895281, BG025115, BG118965, BG117744, BE894516, BE867853, AV655961, AV655065, BE463833, AI743344, AI936434, AA479845, AW069262, AW957415, AI632803, BF435420, AA743099, AI653387, AW139960, BF435344, AW263545, AI986184, AI494572, AW069847, AI356382, BF872252, N64780, AI343604, AI524683, BG254358, AA677460, AA311118, R76972, AV714216, AA485878, BE906880, AI034335, BE836962, AA824645, BG170378, AA421697, AI801400, AV653434, AA324167, AW405847, R17990, AV656465, AI017487, AI283437, AA742903, BF959586, AI432715, H90632, AI523653, AW009118, AI222058, R18249, AA487781, AI935830, N76192, AA333269, H78901, T50567, R20418, AW802235, BF927563, AA610567, H26543, AI040600, AI474629, AI802589, BF926304, AI827833, BF752219, AW292734, N35443, R61739, Z43983, R43537, AA300760, AW118828, AI698966, AA324168, AI915572, BG121443, BE706316, AW235187, BF446401, AA346807, BE168110, AA479721, AA971255, AA776840, AW664573, H39210, AV700303, AI216580, BF964480, BE315009, AW024732, BE206022, AI858448, AW576576, AI133000, AA935149, BG009390, BF884320, AI701554, BE673843, R29282, AW304068, AI244166, BF001810, AW592386, AW170695, C01378, AA683126, H39211, AA857017, AW953126, BE168122, AI351674, BF222238, BF803700, AI769446, BE327201, AI567778, AL034550, AB023195, AK025756, AR072721, AL117647, AL117518, AR072722, and AL034550.
HWTBX66	2590	732187	1 - 1131	15 - 1145	BF206625, BE562605, AI422195, BG111540, BE409545, BF374972, BE899627, AA434145,

					and AC007239.
HYAAC74	2591	762843	1 - 578	15 - 592	AI791896, AW935251, AW935250, AA367713, BF908630, AI821701, AC013276, and AC013276.
HYAAD61	2592	897736	1 - 2216	15 - 2230	AV705135, BE879798, AV655758, BE881037, AV755230, BE875022, BE879931, BE880405, BE879486, BG153219, BE880558, BE877897, AV707414, AV710871, BE878969, AV734388, BE877534, BE876227, AI207423, BE881383, BE877380, BE880003, BE875549, AV712842, BE878544, BE878868, BE880171, BE874129, BE874269, AV709769, AI305275, AI286358, AL039081, AV716518, AI268040, BG223306, AV723801, AI619745, BE874183, BE874959, AW277198, AW277172, AA458950, AW265739, AI557254, AV710411, AI224374, BE880000, BE877313, AV695843, BE873910, AV702721, AI524536, BE898244, BE876255, AI880197, BE877438, AL037048, AA552550, AV689111, AI653760, AV706425, BE877980, AV725088, AI206694, AW151854, AV729555, BE877360, AA244460, BE873776, AV725697, AA554077, AV729458, BE881406, AA565898, BE876129, AA523066, AA640693, AA226180, AV702339, AA654759, AV712023, AA594572, AA081407, BE220724, AA987694, AV704567, BE879161, AV702289, BE873721, AV728530, AV646639, AV715880, AI581199, AV704743, AA460448, AV703548, BE878189, BE875452, AA152380, AA487595, AV702362, BE876209, AV745360, AA192171, AV756865, AV745474, AW080716, AI281861, AV704410, BE873856, AV706544, AV706871, AV704891, AA635387, AV701911, AV705273, AV695888, AV727353, AV727222, BE875679, BE877724, AV703275, AV725220, AA653781, AV704989, AV646609, BE873887, AV703453, BE879920, AV713937, AI282481, AV717895, AW188463, AA523743, AV710865, AV646653, AA854071, AV707576, AA216405, AV706391, AA844117, AV726271, AV708364, AA579042, AA578602, AI630887, AA506176, AL048429, AV726269, AA177050, AA669077, AV646649, AV725312, AV727727, AV711596, AV704136, AA467980, AW419290, AV704821, AA876733, AV715527, BE881017, AV646504, AI133314, AV702418, AV744456, AV701103, AI133420, BE878144, AI065079, AV723380, BE877089, AV701289, AV646554, AV721093, BE876191, AW276852, AV660666, BE876145, AV728726, AI064797, AA631302, AA879253, BE878843, BE875363,

					AV702097, AI884494, BE880616, AV690345, AW073881, AI535915, AI926608, AA873420, AA655025, BE875048, AA642859, AI133218, AV682904, AA602266, AA188353, AV716924, BE881087, AI432215, AV705900, AV722579, BE881605, AV722125, AV751989, AA650124, BE877490, AI798362, AV723748, AI061660, AI832523, AW518375, BE677570, AA569249, AA503933, BE877021, AA069462, AA635953, BE876526, AI673024, AW129234, AV695102, AV693793, AA557411, AI989411, AV693806, AI535901, AW270021, AV686634, AV649901, AV662132, AA621867, AV727103, AI917042, BF061237, AV646616, BE880379, AA887174, BE874933, BE877550, AI636965, BE877849, AI566184, AI566210, AI863894, AI174749, AF161800, AK026530, AK026930, AX023293, AX023297, X62996, AX039612, V00662, X93334, D38112, AF004340, AF134583, X93335, D38113, D38116, X97707, X01775, I27367, AR083056, AK026993, I00734, AF130059, AF110329, AL389951, AK025541, AB050411, AL133645, E12580, E00617, E00717, E00778, S77771, AF125949, AX027129, AB047615, AL137640, I48978, AB007812, AL390167, AF162270, AF012536, AB025103, AF113676, AR038867, AL049466, AL137711, AF274348, AF274347, AL389935, AK026591, AB050534, AF119857, AK025772, AL122106, A65336, AL133098, AF104032, U72620, AC010128, A18777, AK025958, AL359620, AB047966, A08913, AF210052, AL161953, AB029065, AR068466, A08910, AF130100, AC004943, AF242525, A08907, I89931, A08909, AK027113, AK027081, AB047887, AK026613, A08912, AF261883, AK027136, I89934, AF035161, AF113013, AR087170, AF130104, A08908, AP000130, AP000208, and AR038854.
HYACC21	2593	696654	1 - 1209	15 - 1223	AF179633.
HYBAP75	2594	853561	1 - 1154	15 - 1168	AA348133.
HYBAQ24	2595	898367	1 - 1179	15 - 1193	BE672513, AW962455, AW468228, AA593830, BE831720, AW339591, AW440986, AA367270, AW440877, AW592568, AI807052, AI431239, AF111851, AF192522, AC018552, AC018552, and AC018552.
HYBAW56	2596	658736	1 - 570	15 - 584	
HYBBD81	2597	533784	1 - 882	15 - 896	AC004933, AC006028, AK022482, AC013603, and AC015779.

TABLE 4

Code	Description	Tissue	Organ	Cell Line	Disease	Vector
AR022	a_Heart	a_Heart				
AR023	a_Liver	a_Liver				
AR024	a_mammary gland	a_mammary gland				
AR025	a_Prostate	a_Prostate				
AR026	a_small intestine	a_small intestine				
AR027	a_Stomach	a_Stomach				
AR028	Blood B cells	Blood B cells				
AR029	Blood B cells activated	Blood B cells activated				
AR030	Blood B cells resting	Blood B cells resting				
AR031	Blood T cells activated	Blood T cells activated				
AR032	Blood T cells resting	Blood T cells resting				
AR033	brain	brain				
AR034	breast	breast				
AR035	breast cancer	breast cancer				
AR036	Cell Line CAOV3	Cell Line CAOV3				
AR037	cell line PA-1	cell line PA-1				
AR038	cell line transformed	cell line transformed				
AR039	colon	colon				
AR040	colon (9808co65R)	colon (9808co65R)				
AR041	colon (9809co15)	colon (9809co15)				
AR042	colon cancer	colon cancer				
AR043	colon cancer (9808co64R)	colon cancer (9808co64R)				
AR044	colon cancer 9809co14	colon cancer 9809co14				
AR045	corn clone 5	corn clone 5				
AR046	corn clone 6	corn clone 6				
AR047	corn clone2	corn clone2				
AR048	corn clone3	corn clone3				
AR049	Corn Clone4	Corn Clone4				
AR050	Donor II B Cells 24hrs	Donor II B Cells 24hrs				
AR051	Donor II B Cells 72hrs	Donor II B Cells 72hrs				
AR052	Donor II B-Cells 24 hrs.	Donor II B-Cells 24 hrs.				
AR053	Donor II B-Cells 72hrs	Donor II B-Cells 72hrs				
AR054	Donor II Resting B Cells	Donor II Resting B Cells				
AR055	Heart	Heart				
AR056	Human Lung (clonotech)	Human Lung (clonotech)				
AR057	Human Mammary (clontech)	Human Mammary (clontech)				
AR058	Human Thymus (clonotech)	Human Thymus (clonotech)				
AR059	Jurkat (unstimulated)	Jurkat (unstimulated)				
AR060	Kidney	Kidney				
AR061	Liver	Liver				
AR062	Liver (Clontech)	Liver (Clontech)				

AR063	Lymphocytes chronic lymphocytic leukaemia	Lymphocytes chronic lymphocytic leukaemia				
AR064	Lymphocytes diffuse large B cell lymphoma	Lymphocytes diffuse large B cell lymphoma				
AR065	Lymphocytes follicular lymphoma	Lymphocytes follicular lymphoma				
AR066	normal breast	normal breast				
AR067	Normal Ovarian (4004901)	Normal Ovarian (4004901)				
AR068	Normal Ovary 9508G045	Normal Ovary 9508G045				
AR069	Normal Ovary 9701G208	Normal Ovary 9701G208				
AR070	Normal Ovary 9806G005	Normal Ovary 9806G005				
AR071	Ovarian Cancer	Ovarian Cancer				
AR072	Ovarian Cancer (9702G001)	Ovarian Cancer (9702G001)				
AR073	Ovarian Cancer (9707G029)	Ovarian Cancer (9707G029)				
AR074	Ovarian Cancer (9804G011)	Ovarian Cancer (9804G011)				
AR075	Ovarian Cancer (9806G019)	Ovarian Cancer (9806G019)				
AR076	Ovarian Cancer (9807G017)	Ovarian Cancer (9807G017)				
AR077	Ovarian Cancer (9809G001)	Ovarian Cancer (9809G001)				
AR078	ovarian cancer 15799	ovarian cancer 15799				
AR079	Ovarian Cancer 17717AID	Ovarian Cancer 17717AID				
AR080	Ovarian Cancer 4004664B1	Ovarian Cancer 4004664B1				
AR081	Ovarian Cancer 4005315A1	Ovarian Cancer 4005315A1				
AR082	ovarian cancer 94127303	ovarian cancer 94127303				
AR083	Ovarian Cancer 96069304	Ovarian Cancer 96069304				
AR084	Ovarian Cancer 9707G029	Ovarian Cancer 9707G029				
AR085	Ovarian Cancer 9807G045	Ovarian Cancer 9807G045				
AR086	ovarian cancer 9809G001	ovarian cancer 9809G001				
AR087	Ovarian Cancer 9905C032RC	Ovarian Cancer 9905C032RC				
AR088	Ovarian cancer 9907 C00 3rd	Ovarian cancer 9907 C00 3rd				
AR089	Prostate	Prostate				
AR090	Prostate (clonotech)	Prostate (clonotech)				
AR091	prostate cancer	prostate cancer				
AR092	prostate cancer #15176	prostate cancer #15176				
AR093	prostate cancer #15509	prostate cancer #15509				
AR094	prostate cancer #15673	prostate cancer #15673				
AR095	Small Intestine (Clontech)	Small Intestine (Clontech)				
AR096	Spleen	Spleen				
AR097	Thymus T cells activated	Thymus T cells				

		activated				
AR098	Thymus T cells resting	Thymus T cells resting				
AR099	Tonsil	Tonsil				
AR100	Tonsil germinal center centroblast	Tonsil germinal center centroblast				
AR101	Tonsil germinal center B cell	Tonsil germinal center B cell				
AR102	Tonsil lymph node	Tonsil lymph node				
AR103	Tonsil memory B cell	Tonsil memory B cell				
AR104	Whole Brain	Whole Brain				
AR105	Xenograft ES-2	Xenograft ES-2				
AR106	Xenograft SW626	Xenograft SW626				
AR119	001: IL-2	001: IL-2				
AR120	001: IL-2.1	001: IL-2.1				
AR121	001: IL-2_b	001: IL-2_b				
AR124	002 : Monocytes untreated (1hr)	002 : Monocytes untreated (1hr)				
AR125	002 : Monocytes untreated (5hrs)	002 : Monocytes untreated (5hrs)				
AR126	002: Control.1C	002: Control.1C				
AR127	002: IL2.1C	002: IL2.1C				
AR130	003 : Placebo-treated Rat Lacrimal Gland	003 : Placebo-treated Rat Lacrimal Gland				
AR131	003 : Placebo-treated Rat Submandibular Gland	003 : Placebo-treated Rat Submandibular Gland				
AR135	004 : Monocytes untreated (5hrs)	004 : Monocytes untreated (5hrs)				
AR136	004 : Monocytes untreated 1hr	004 : Monocytes untreated 1hr				
AR139	005: Placebo (48hrs)	005: Placebo (48hrs)				
AR140	006: pC4 (24hrs)	006: pC4 (24hrs)				
AR141	006: pC4 (48hrs)	006: pC4 (48hrs)				
AR152	007: PHA(1hr)	007: PHA(1hr)				
AR153	007: PHA(6HRS)	007: PHA(6HRS)				
AR154	007: PMA(6hrs)	007: PMA(6hrs)				
AR155	008: 1449_#2	008: 1449_#2				
AR161	01: A - max 24	01: A - max 24				
AR162	01: A - max 26	01: A - max 26				
AR163	01: A - max 30	01: A - max 30				
AR164	01: B - max 24	01: B - max 24				
AR165	01: B - max 26	01: B - max 26				
AR166	01: B - max 30	01: B - max 30				
AR167	1449 Sample	1449 Sample				
AR168	3T3P10 1.0uM insulin	3T3P10 1.0uM insulin				
AR169	3T3P10 10nM Insulin	3T3P10 10nM Insulin				
AR170	3T3P10 10uM insulin	3T3P10 10uM insulin				
AR171	3T3P10 No Insulin	3T3P10 No Insulin				
AR172	3T3P4	3T3P4				
AR173	Adipose (41892)	Adipose (41892)				
AR174	Adipose Diabetic (41611)	Adipose Diabetic (41611)				
AR175	Adipose Diabetic (41661)	Adipose Diabetic (41661)				
AR176	Adipose Diabetic (41689)	Adipose Diabetic (41689)				
AR177	Adipose Diabetic (41706)	Adipose Diabetic				

		(41706)				
AR178	Adipose Diabetic (42352)	Adipose Diabetic (42352)				
AR179	Adipose Diabetic (42366)	Adipose Diabetic (42366)				
AR180	Adipose Diabetic (42452)	Adipose Diabetic (42452)				
AR181	Adipose Diabetic (42491)	Adipose Diabetic (42491)				
AR182	Adipose Normal (41843)	Adipose Normal (41843)				
AR183	Adipose Normal (41893)	Adipose Normal (41893)				
AR184	Adipose Normal (42452)	Adipose Normal (42452)				
AR185	Adrenal Gland	Adrenal Gland				
AR186	Adrenal Gland + Whole Brain	Adrenal Gland + Whole Brain				
AR187	B7(1hr)+ (inverted)	B7(1hr)+ (inverted)				
AR188	Breast (18275A2B)	Breast (18275A2B)				
AR189	Breast (4004199)	Breast (4004199)				
AR190	Breast (4004399)	Breast (4004399)				
AR191	Breast (4004943B7)	Breast (4004943B7)				
AR192	Breast (4005570B1)	Breast (4005570B1)				
AR193	Breast Cancer (4004127A30)	Breast Cancer (4004127A30)				
AR194	Breast Cancer (400443A21)	Breast Cancer (400443A21)				
AR195	Breast Cancer (4004643A2)	Breast Cancer (4004643A2)				
AR196	Breast Cancer (4004710A7)	Breast Cancer (4004710A7)				
AR197	Breast Cancer (4004943A21)	Breast Cancer (4004943A21)				
AR198	Breast Cancer (400553A2)	Breast Cancer (400553A2)				
AR199	Breast Cancer (9805C046R)	Breast Cancer (9805C046R)				
AR200	Breast Cancer (9806C012R)	Breast Cancer (9806C012R)				
AR201	Breast Cancer (ODQ 45913)	Breast Cancer (ODQ 45913)				
AR202	Breast Cancer (ODQ45913)	Breast Cancer (ODQ45913)				
AR203	Breast Cancer (ODQ4591B)	Breast Cancer (ODQ4591B)				
AR204	Colon Cancer (15663)	Colon Cancer (15663)				
AR205	Colon Cancer (4005144A4)	Colon Cancer (4005144A4)				
AR206	Colon Cancer (4005413A4)	Colon Cancer (4005413A4)				
AR207	Colon Cancer (4005570B1)	Colon Cancer (4005570B1)				
AR208	Control RNA #1	Control RNA #1				
AR209	Control RNA #2	Control RNA #2				
AR210	Cultured Preadipocyte (blue)	Cultured Preadipocyte (blue)				
AR211	Cultured Preadipocyte (Red)	Cultured Preadipocyte (Red)				
AR212	Donor II B-Cells 24hrs	Donor II B-Cells 24hrs				
AR213	Donor II Resting B-Cells	Donor II Resting B-Cells				

AR214	H114EP12 10nM Insulin	H114EP12 10nM Insulin				
AR215	H114EP12 (10nM insulin)	H114EP12 (10nM insulin)				
AR216	H114EP12 (2.6ug/ul)	H114EP12 (2.6ug/ul)				
AR217	H114EP12 (3.6ug/ul)	H114EP12 (3.6ug/ul)				
AR218	HUVEC #1	HUVEC #1				
AR219	HUVEC #2	HUVEC #2				
AR221	L6 undiff.	L6 undiff.				
AR222	L6 Undifferentiated	L6 Undifferentiated				
AR223	L6P8 + 10nM Insulin	L6P8 + 10nM Insulin				
AR224	L6P8 + HS	L6P8 + HS				
AR225	L6P8 10nM Insulin	L6P8 10nM Insulin				
AR226	Liver (00-06-A007B)	Liver (00-06-A007B)				
AR227	Liver (96-02-A075)	Liver (96-02-A075)				
AR228	Liver (96-03-A144)	Liver (96-03-A144)				
AR229	Liver (96-04-A138)	Liver (96-04-A138)				
AR230	Liver (97-10-A074B)	Liver (97-10-A074B)				
AR231	Liver (98-09-A242A)	Liver (98-09-A242A)				
AR232	Liver Diabetic (1042)	Liver Diabetic (1042)				
AR233	Liver Diabetic (41616)	Liver Diabetic (41616)				
AR234	Liver Diabetic (41955)	Liver Diabetic (41955)				
AR235	Liver Diabetic (42352R)	Liver Diabetic (42352R)				
AR236	Liver Diabetic (42366)	Liver Diabetic (42366)				
AR237	Liver Diabetic (42483)	Liver Diabetic (42483)				
AR238	Liver Diabetic (42491)	Liver Diabetic (42491)				
AR239	Liver Diabetic (99-09-A281A)	Liver Diabetic (99-09-A281A)				
AR240	Lung	Lung				
AR241	Lung (27270)	Lung (27270)				
AR242	Lung (2727Q)	Lung (2727Q)				
AR243	Lung Cancer (4005116A1)	Lung Cancer (4005116A1)				
AR244	Lung Cancer (4005121A5)	Lung Cancer (4005121A5)				
AR245	Lung Cancer (4005121A5))	Lung Cancer (4005121A5))				
AR246	Lung Cancer (4005340A4)	Lung Cancer (4005340A4)				
AR247	Mammary Gland	Mammary Gland				
AR248	Monocyte (CT)	Monocyte (CT)				
AR249	Monocyte (OCT)	Monocyte (OCT)				
AR250	Monocytes (CT)	Monocytes (CT)				
AR251	Monocytes (INFG 18 hr)	Monocytes (INFG 18 hr)				
AR252	Monocytes (INFG 18hr)	Monocytes (INFG 18hr)				
AR253	Monocytes (INFG 8-11)	Monocytes (INFG 8-11)				
AR254	Monocytes (O CT)	Monocytes (O CT)				
AR255	Muscle (91-01-A105)	Muscle (91-01-				

		A105)				
AR256	Muscle (92-04-A059)	Muscle (92-04-A059)				
AR257	Muscle (97-11-A056d)	Muscle (97-11-A056d)				
AR258	Muscle (99-06-A210A)	Muscle (99-06-A210A)				
AR259	Muscle (99-07-A203B)	Muscle (99-07-A203B)				
AR260	Muscle (99-7-A203B)	Muscle (99-7-A203B)				
AR261	Muscle Diabetic (42352R)	Muscle Diabetic (42352R)				
AR262	Muscle Diabetic (42366)	Muscle Diabetic (42366)				
AR263	NK-19 Control	NK-19 Control				
AR264	NK-19 IL Treated 72hrs	NK-19 IL Treated 72hrs				
AR265	NK-19 UK Treated 72 hrs.	NK-19 UK Treated 72 hrs.				
AR266	Omentum Normal (94-08-B009)	Omentum Normal (94-08-B009)				
AR267	Omentum Normal (97-01-A039A)	Omentum Normal (97-01-A039A)				
AR268	Omentum Normal (97-04-A114C)	Omentum Normal (97-04-A114C)				
AR269	Omentum Normal (97-06-A117C)	Omentum Normal (97-06-A117C)				
AR270	Omentum Normal (97-09-B004C)	Omentum Normal (97-09-B004C)				
AR271	Ovarian Cancer (17717AID)	Ovarian Cancer (17717AID)				
AR272	Ovarian Cancer (9905C023RC)	Ovarian Cancer (9905C023RC)				
AR273	Ovarian Cancer (9905C032RC)	Ovarian Cancer (9905C032RC)				
AR274	Ovary (9508G045)	Ovary (9508G045)				
AR275	Ovary (9701G208)	Ovary (9701G208)				
AR276	Ovary 9806G005	Ovary 9806G005				
AR277	Pancreas	Pancreas				
AR278	Placebo	Placebo				
AR279	rIL2 Control	rIL2 Control				
AR280	RSS288L	RSS288L				
AR281	RSS288LC	RSS288LC				
AR282	Salivary Gland	Salivary Gland				
AR283	Skeletal Muscle	Skeletal Muscle				
AR284	Skeletal Muscle (91-01-A105)	Skeletal Muscle (91-01-A105)				
AR285	Skeletal Muscle (42180)	Skeletal Muscle (42180)				
AR286	Skeletal Muscle (42386)	Skeletal Muscle (42386)				
AR287	Skeletal Muscle (42461)	Skeletal Muscle (42461)				
AR288	Skeletal Muscle (91-01-A105)	Skeletal Muscle (91-01-A105)				
AR289	Skeletal Muscle (92-04-A059)	Skeletal Muscle (92-04-A059)				
AR290	Skeletal Muscle (96-08-A171)	Skeletal Muscle (96-08-A171)				
AR291	Skeletal Muscle (97-07-A190A)	Skeletal Muscle (97-07-A190A)				
AR292	Skeletal Muscle Diabetic (42352)	Skeletal Muscle Diabetic (42352)				
AR293	Skeletal Muscle Diabetic	Skeletal Muscle				

	(42366)	Diabetic (42366)				
AR294	Skeletal Muscle Diabetic (42395)	Skeletal Muscle Diabetic (42395)				
AR295	Skeletal Muscle Diabetic (42483)	Skeletal Muscle Diabetic (42483)				
AR296	Skeletal Muscle Diabetic (42491)	Skeletal Muscle Diabetic (42491)				
AR297	Skeletal Muscle Diabetic 42352	Skeletal Muscle Diabetic 42352				
AR298	Skeletal Musle (42461)	Skeletal Musle (42461)				
AR299	Small Intestine	Small Intestine				
AR300	Stomach	Stomach				
AR301	T-Cell + HDPBQ71.fc 1449 16hrs	T-Cell + HDPBQ71.fc 1449 16hrs				
AR302	T-Cell + HDPBQ71.fc 1449 6hrs	T-Cell + HDPBQ71.fc 1449 6hrs				
AR303	T-Cell + IL2 16hrs	T-Cell + IL2 16hrs				
AR304	T-Cell + IL2 6hrs	T-Cell + IL2 6hrs				
AR306	T-Cell Untreated 16hrs	T-Cell Untreated 16hrs				
AR307	T-Cell Untreated 6hrs	T-Cell Untreated 6hrs				
AR308	T-Cells 24 hours	T-Cells 24 hours				
AR309	T-Cells 24 hrs	T-Cells 24 hrs				
AR310	T-Cells 24 hrs.	T-Cells 24 hrs.				
AR311	T-Cells 24hrs	T-Cells 24hrs				
AR312	T-Cells 4 days	T-Cells 4 days				
AR313	Thymus	Thymus				
AR314	TRE	TRE				
AR315	TREC	TREC				
AR316	Virtual Mixture	Virtual Mixture				
H0002	Human Adult Heart	Human Adult Heart	Heart			Uni-ZAP XR
H0003	Human Adult Liver	Human Adult Liver	Liver			Uni-ZAP XR
H0004	Human Adult Spleen	Human Adult Spleen	Spleen			Uni-ZAP XR
H0007	Human Cerebellum	Human Cerebellum	Brain			Uni-ZAP XR
H0008	Whole 6 Week Old Embryo					Uni-ZAP XR
H0009	Human Fetal Brain					Uni-ZAP XR
H0010	Human Fetal Hepatic	Human Fetal Liver	Liver			Uni-ZAP XR
H0011	Human Fetal Kidney	Human Fetal Kidney	Kidney			Uni-ZAP XR
H0012	Human Fetal Kidney	Human Fetal Kidney	Kidney			Uni-ZAP XR
H0013	Human 8 Week Whole Embryo	Human 8 Week Old Embryo	Embryo			Uni-ZAP XR
H0014	Human Gall Bladder	Human Gall Bladder	Gall Bladder			Uni-ZAP XR
H0015	Human Gall Bladder, fraction II	Human Gall Bladder	Gall Bladder			Uni-ZAP XR
H0018	Human Greater Omentum, fII remake	Human Greater Omentum	peritoneum			Uni-ZAP XR
H0019	Human Fetal Heart	Human Fetal Heart	Heart			pBluescript
H0020	Human Hippocampus	Human Hippocampus	Brain			Uni-ZAP XR
H0021	Human Infant Adrenal Gland	Human Infant Adrenal Gland	Adrenal gland			Uni-ZAP XR
H0022	Jurkat Cells	Jurkat T-Cell Line				Lambda ZAP II
H0023	Human Fetal Lung					Uni-ZAP XR
H0024	Human Fetal Lung III	Human Fetal Lung	Lung			Uni-ZAP XR
H0025	Human Adult Lymph Node	Human Adult Lymph Node	Lymph Node			Lambda ZAP II
H0026	Namalwa Cells	Namalwa B-Cell Line, EBV				Lambda ZAP II

		immortalized				
H0028	Human Old Ovary	Human Old Ovary	Ovary			pBluescript
H0029	Human Pancreas	Human Pancreas	Pancreas			Uni-ZAP XR
H0030	Human Placenta					Uni-ZAP XR
H0031	Human Placenta	Human Placenta	Placenta			Uni-ZAP XR
H0032	Human Prostate	Human Prostate	Prostate			Uni-ZAP XR
H0033	Human Pituitary	Human Pituitary				Uni-ZAP XR
H0036	Human Adult Small Intestine	Human Adult Small Intestine	Small Int.			Uni-ZAP XR
H0037	Human Adult Small Intestine	Human Adult Small Intestine	Small Int.			pBluescript
H0038	Human Testes	Human Testes	Testis			Uni-ZAP XR
H0039	Human Pancreas Tumor	Human Pancreas Tumor	Pancreas		disease	Uni-ZAP XR
H0040	Human Testes Tumor	Human Testes Tumor	Testis		disease	Uni-ZAP XR
H0041	Human Fetal Bone	Human Fetal Bone	Bone			Uni-ZAP XR
H0042	Human Adult Pulmonary	Human Adult Pulmonary	Lung			Uni-ZAP XR
H0044	Human Cornea	Human Cornea	eye			Uni-ZAP XR
H0045	Human Esophagus, Cancer	Human Esophagus, cancer	Esophagus		disease	Uni-ZAP XR
H0046	Human Endometrial Tumor	Human Endometrial Tumor	Uterus		disease	Uni-ZAP XR
H0047	Human Fetal Liver	Human Fetal Liver	Liver			Uni-ZAP XR
H0048	Human Pineal Gland	Human Pineal Gland				Uni-ZAP XR
H0050	Human Fetal Heart	Human Fetal Heart	Heart			Uni-ZAP XR
H0051	Human Hippocampus	Human Hippocampus	Brain			Uni-ZAP XR
H0052	Human Cerebellum	Human Cerebellum	Brain			Uni-ZAP XR
H0053	Human Adult Kidney	Human Adult Kidney	Kidney			Uni-ZAP XR
H0056	Human Umbilical Vein, Endo. remake	Human Umbilical Vein Endothelial Cells	Umbilical vein			Uni-ZAP XR
H0057	Human Fetal Spleen					Uni-ZAP XR
H0058	Human Thymus Tumor	Human Thymus Tumor	Thymus		disease	Lambda ZAP II
H0059	Human Uterine Cancer	Human Uterine Cancer	Uterus		disease	Lambda ZAP II
H0060	Human Macrophage	Human Macrophage	Blood	Cell Line		pBluescript
H0061	Human Macrophage	Human Macrophage	Blood	Cell Line		pBluescript
H0062	Human Thymus	Human Thymus	Thymus			Uni-ZAP XR
H0063	Human Thymus	Human Thymus	Thymus			Uni-ZAP XR
H0064	Human Right Hemisphere of Brain	Human Brain, right hemisphere	Brain			Uni-ZAP XR
H0067	Human left hemisphere, adult	Human Left Hemisphere, Adult	Brain			Lambda ZAP II
H0068	Human Skin Tumor	Human Skin Tumor	Skin		disease	Uni-ZAP XR
H0069	Human Activated T-Cells	Activated T-Cells	Blood	Cell Line		Uni-ZAP XR
H0070	Human Pancreas	Human Pancreas	Pancreas			Uni-ZAP XR
H0071	Human Infant Adrenal Gland	Human Infant Adrenal Gland	Adrenal gland			Uni-ZAP XR
H0073	Human Leiomyeloid Carcinoma	Human Leiomyeloid Carcinoma	Muscle		disease	Uni-ZAP XR
H0074	Human Platelets	Human Platelets	Blood	Cell Line		Uni-ZAP XR
H0075	Human Activated T-Cells (II)	Activated T-Cells	Blood	Cell Line		Uni-ZAP XR
H0077	Human Thymus Tumor	Human Thymus Tumor	Thymus		disease	Lambda ZAP II
H0078	Human Lung Cancer	Human Lung Cancer	Lung		disease	Lambda ZAP II
H0079	Human Whole 7 Week Old Embryo (II)	Human Whole 7 Week Old Embryo	Embryo			Uni-ZAP XR
H0080	Human Whole 6 Week	Human Whole Six	Embryo			Lambda ZAP II

	Old Embryo (II)	Week Old Embryo				
H0081	Human Fetal Epithelium (Skin)	Human Fetal Skin	Skin			Uni-ZAP XR
H0082	Human Fetal Muscle	Human Fetal Muscle	Sk Muscle			Uni-ZAP XR
H0083	HUMAN JURKAT MEMBRANE BOUND POLYSOMES	Jurkat Cells				Uni-ZAP XR
H0085	Human Colon	Human Colon				Lambda ZAP II
H0086	Human epithelioid sarcoma	Epithelioid Sarcoma, muscle	Sk Muscle		disease	Uni-ZAP XR
H0087	Human Thymus	Human Thymus				pBluescript
H0090	Human T-Cell Lymphoma	T-Cell Lymphoma	T-Cell		disease	Uni-ZAP XR
H0092	Human Pancreas Tumor	Human Pancreas Tumor	Pancreas		disease	Uni-ZAP XR
H0095	Human Greater Omentum, RNA Remake	Human Greater Omentum	peritoneum			Uni-ZAP XR
H0097	Human Adult Heart, subtracted	Human Adult Heart	Heart			pBluescript
H0098	Human Adult Liver, subtracted	Human Adult Liver	Liver			Uni-ZAP XR
H0099	Human Lung Cancer, subtracted	Human Lung Cancer	Lung			pBluescript
H0100	Human Whole Six Week Old Embryo	Human Whole Six Week Old Embryo	Embryo			Uni-ZAP XR
H0101	Human 7 Weeks Old Embryo, subtracted	Human Whole 7 Week Old Embryo	Embryo			Lambda ZAP II
H0102	Human Whole 6 Week Old Embryo (II), subt	Human Whole Six Week Old Embryo	Embryo			pBluescript
H0103	Human Fetal Brain, subtracted	Human Fetal Brain	Brain			Uni-ZAP XR
H0105	Human Fetal Heart, subtracted	Human Fetal Heart	Heart			pBluescript
H0107	Human Infant Adrenal Gland, subtracted	Human Infant Adrenal Gland	Adrenal gland			pBluescript
H0108	Human Adult Lymph Node, subtracted	Human Adult Lymph Node	Lymph Node			Uni-ZAP XR
H0109	Human Macrophage, subtracted	Macrophage	Blood	Cell Line		pBluescript
H0110	Human Old Ovary, subtracted	Human Old Ovary	Ovary			pBluescript
H0111	Human Placenta, subtracted	Human Placenta	Placenta			pBluescript
H0113	Human skin Tumor, subtracted	Human Skin Tumor	Skin			Uni-ZAP XR
H0116	Human Thymus Tumor, subtracted	Human Thymus Tumor	Thymus			pBluescript
H0117	Human Uterine Cancer, subtracted	Human Uterine Cancer	Uterus			pBluescript
H0119	Human Pediatric Kidney	Human Pediatric Kidney	Kidney			Uni-ZAP XR
H0120	Human Adult Spleen, subtracted	Human Adult Spleen	Spleen			Uni-ZAP XR
H0121	Human Cornea, subtracted	Human Cornea	eye			Uni-ZAP XR
H0122	Human Adult Skeletal Muscle	Human Skeletal Muscle	Sk Muscle			Uni-ZAP XR
H0123	Human Fetal Dura Mater	Human Fetal Dura Mater	Brain			Uni-ZAP XR
H0124	Human Rhabdomyosarcoma	Human Rhabdomyosarcoma	Sk Muscle		disease	Uni-ZAP XR
H0125	Cem cells cyclohexamide treated	Cyclohexamide Treated Cem, Jurkat, Raji, and Supt	Blood	Cell Line		Uni-ZAP XR
H0128	Jurkat cells, thiouridine activated	Jurkat Cells				Uni-ZAP XR
H0130	LNCAP untreated	LNCAP Cell Line	Prostate	Cell Line		Uni-ZAP XR

H0131	LNCAP + 0.3nM R1881	LNCAP Cell Line	Prostate	Cell Line		Uni-ZAP XR
H0132	LNCAP + 30nM R1881	LNCAP Cell Line	Prostate	Cell Line		Uni-ZAP XR
H0134	Raji Cells, cyclohexamide treated	Cyclohexamide Treated Cem, Jurkat, Raji, and Supt	Blood	Cell Line		Uni-ZAP XR
H0135	Human Synovial Sarcoma	Human Synovial Sarcoma	Synovium			Uni-ZAP XR
H0136	Supt Cells, cyclohexamide treated	Cyclohexamide Treated Cem, Jurkat, Raji, and Supt	Blood	Cell Line		Uni-ZAP XR
H0140	Activated T-Cells, 8 hrs.	Activated T-Cells	Blood	Cell Line		Uni-ZAP XR
H0141	Activated T-Cells, 12 hrs.	Activated T-Cells	Blood	Cell Line		Uni-ZAP XR
H0144	Nine Week Old Early Stage Human	9 Wk Old Early Stage Human	Embryo			Uni-ZAP XR
H0147	Human Adult Liver	Human Adult Liver	Liver			Uni-ZAP XR
H0149	7 Week Old Early Stage Human, subtracted	Human Whole 7 Week Old Embryo	Embryo			Uni-ZAP XR
H0150	Human Epididymus	Epididymis	Testis			Uni-ZAP XR
H0151	Early Stage Human Liver	Human Fetal Liver	Liver			Uni-ZAP XR
H0152	Early Stage Human Liver, fract (II)	Human Fetal Liver	Liver			Uni-ZAP XR
H0154	Human Fibrosarcoma	Human Skin Fibrosarcoma	Skin		disease	Uni-ZAP XR
H0155	Human Thymus, subtracted	Human Thymus Tumor	Thymus			pBluescript
H0156	Human Adrenal Gland Tumor	Human Adrenal Gland Tumor	Adrenal Gland		disease	Uni-ZAP XR
H0157	Activated T-Cells, 0 hrs, ligation 2	Activated T-Cells	Blood	Cell Line		Uni-ZAP XR
H0158	Activated T-Cells, 4 hrs., ligation 2	Activated T-Cells	Blood	Cell Line		Uni-ZAP XR
H0159	Activated T-Cells, 8 hrs., ligation 2	Activated T-Cells	Blood	Cell Line		Uni-ZAP XR
H0163	Human Synovium	Human Synovium	Synovium			Uni-ZAP XR
H0164	Human Trachea Tumor	Human Trachea Tumor	Trachea		disease	Uni-ZAP XR
H0165	Human Prostate Cancer, Stage B2	Human Prostate Cancer, stage B2	Prostate		disease	Uni-ZAP XR
H0166	Human Prostate Cancer, Stage B2 fraction	Human Prostate Cancer, stage B2	Prostate		disease	Uni-ZAP XR
H0167	Activated T-Cells, 24 hrs.	Activated T-Cells	Blood	Cell Line		Uni-ZAP XR
H0168	Human Prostate Cancer, Stage C	Human Prostate Cancer, stage C	Prostate		disease	Uni-ZAP XR
H0169	Human Prostate Cancer, Stage C fraction	Human Prostate Cancer, stage C	Prostate		disease	Uni-ZAP XR
H0170	12 Week Old Early Stage Human	Twelve Week Old Early Stage Human	Embryo			Uni-ZAP XR
H0171	12 Week Old Early Stage Human, II	Twelve Week Old Early Stage Human	Embryo			Uni-ZAP XR
H0172	Human Fetal Brain, random primed	Human Fetal Brain	Brain			Lambda ZAP II
H0173	Human Cardiomyopathy, RNA remake	Human Cardiomyopathy	Heart		disease	Uni-ZAP XR
H0175	H. Adult Spleen, ziplox					pSport1
H0176	CAMA1Ee Cell Line	CAMA1Ee Cell Line	Breast	Cell Line		Uni-ZAP XR
H0177	CAMA1Ee Cell Line	CAMA1Ee Cell Line	Breast	Cell Line		Uni-ZAP XR
H0178	Human Fetal Brain	Human Fetal Brain	Brain			Uni-ZAP XR
H0179	Human Neutrophil	Human Neutrophil	Blood	Cell Line		Uni-ZAP XR
H0181	Human Primary Breast Cancer	Human Primary Breast Cancer	Breast		disease	Uni-ZAP XR
H0182	Human Primary Breast Cancer	Human Primary Breast Cancer	Breast		disease	Uni-ZAP XR
H0183	Human Colon Cancer	Human Colon	Colon		disease	Uni-ZAP XR

		Cancer				
H0184	Human Colon Cancer, metasticized to live	Human Colon Cancer, metasticized to liver	Liver		disease	Lambda ZAP II
H0185	Activated T-Cell labeled with 4-thioluri	T-Cells	Blood	Cell Line		Lambda ZAP II
H0186	Activated T-Cell	T-Cells	Blood	Cell Line		Lambda ZAP II
H0187	Resting T-Cell	T-Cells	Blood	Cell Line		Lambda ZAP II
H0188	Human Normal Breast	Human Normal Breast	Breast			Uni-ZAP XR
H0189	Human Resting Macrophage	Human Macrophage/Monocytes	Blood	Cell Line		Uni-ZAP XR
H0190	Human Activated Macrophage (LPS)	Human Macrophage/Monocytes	Blood	Cell Line		Uni-ZAP XR
H0192	Cem Cells, cyclohexamide treated, subtra	Cyclohexamide Treated Cem, Jurkat, Raji, and Supt	Blood	Cell Line		Uni-ZAP XR
H0193	Cem Cells, cyclohexamide treated, differ	Cyclohexamide Treated Cem, Jurkat, Raji, and Supt	Blood	Cell Line		Uni-ZAP XR
H0194	Human Cerebellum, subtracted	Human Cerebellum	Brain			pBluescript
H0196	Human Cardiomyopathy, subtracted	Human Cardiomyopathy	Heart			Uni-ZAP XR
H0197	Human Fetal Liver, subtracted	Human Fetal Liver	Liver			Uni-ZAP XR
H0198	Human Fetal Liver, subtracted, pos. clon	Human Fetal Liver	Liver			Uni-ZAP XR
H0199	Human Fetal Liver, subtracted, neg clone	Human Fetal Liver	Liver			Uni-ZAP XR
H0200	Human Greater Omentum, fract II remake,	Human Greater Omentum	peritoneum			Uni-ZAP XR
H0201	Human Hippocampus, subtracted	Human Hippocampus	Brain			pBluescript
H0202	Jurkat Cells, cyclohexamide treated, subtraction	Cyclohexamide Treated Cem, Jurkat, Raji, and Supt	Blood	Cell Line		Uni-ZAP XR
H0203	Jurkat Cells, cyclohexamide treated, dif	Cyclohexamide Treated Cem, Jurkat, Raji, and Supt	Blood	Cell Line		Uni-ZAP XR
H0204	Human Colon Cancer, subtracted	Human Colon Cancer	Colon			pBluescript
H0205	Human Colon Cancer, differential	Human Colon Cancer	Colon			pBluescript
H0207	LNCAP, differential expression	LNCAP Cell Line	Prostate	Cell Line		pBluescript
H0208	Early Stage Human Lung, subtracted	Human Fetal Lung	Lung			pBluescript
H0209	Human Cerebellum, differentially expressed	Human Cerebellum	Brain			Uni-ZAP XR
H0211	Human Prostate, differential expression	Human Prostate	Prostate			pBluescript
H0212	Human Prostate, subtracted	Human Prostate	Prostate			pBluescript
H0213	Human Pituitary, subtracted	Human Pituitary				Uni-ZAP XR
H0214	Raji cells, cyclohexamide treated, subtracted	Cyclohexamide Treated Cem, Jurkat, Raji, and Supt	Blood	Cell Line		pBluescript
H0216	Supt cells, cyclohexamide treated, subtracted	Cyclohexamide Treated Cem, Jurkat, Raji, and Supt	Blood	Cell Line		pBluescript

H0217	Supt cells, cyclohexamide treated, differentially expressed	Cyclohexamide Treated Cem, Jurkat, Raji, and Supt	Blood	Cell Line		pBluescript
H0218	Activated T-Cells, 0hrs, subtracted	Activated T-Cells	Blood	Cell Line		Uni-ZAP XR
H0219	Activated T-Cells, 0hrs, differentially expressed	Activated T-Cells	Blood	Cell Line		Uni-ZAP XR
H0220	Activated T-Cells, 4 hrs, subtracted	Activated T-Cells	Blood	Cell Line		Uni-ZAP XR
H0222	Activated T-Cells, 8 hrs, subtracted	Activated T-Cells	Blood	Cell Line		Uni-ZAP XR
H0224	Activated T-Cells, 12 hrs, subtracted	Activated T-Cells	Blood	Cell Line		Uni-ZAP XR
H0225	Activated T-Cells, 12hrs, differentially expressed	Activated T-Cells	Blood	Cell Line		Uni-ZAP XR
H0228	C7MCF7 cell line, estrogen treated	C7MCF7 Cell Line, estrogen treated	Breast	Cell Line		Uni-ZAP XR
H0229	Early Stage Human Brain, random primed	Early Stage Human Brain	Brain			Lambda ZAP II
H0230	Human Cardiomyopathy, diff exp	Human Cardiomyopathy	Heart		disease	Uni-ZAP XR
H0231	Human Colon, subtraction	Human Colon				pBluescript
H0232	Human Colon, differential expression	Human Colon				pBluescript
H0233	Human Fetal Heart, Differential (Adult-Specific)	Human Fetal Heart	Heart			pBluescript
H0234	human colon cancer, metastatic to liver, differentially expressed	Human Colon Cancer, metastaticized to liver	Liver			pBluescript
H0235	Human colon cancer, metaticized to liver, subtraction	Human Colon Cancer, metastaticized to liver	Liver			pBluescript
H0239	Human Kidney Tumor	Human Kidney Tumor	Kidney		disease	Uni-ZAP XR
H0241	C7MCF7 cell line, estrogen treated, subtraction	C7MCF7 Cell Line, estrogen treated	Breast	Cell Line		Uni-ZAP XR
H0242	Human Fetal Heart, Differential (Fetal-Specific)	Human Fetal Heart	Heart			pBluescript
H0244	Human 8 Week Whole Embryo, subtracted	Human 8 Week Old Embryo	Embryo			Uni-ZAP XR
H0246	Human Fetal Liver-Enzyme subtraction	Human Fetal Liver	Liver			Uni-ZAP XR
H0247	Human Membrane Bound Polysomes- Enzyme Subtraction	Human Membrane Bound Polysomes	Blood	Cell Line		Uni-ZAP XR
H0249	HE7, subtracted by hybridization with E7 cDNA	Human Whole 7 Week Old Embryo	Embryo			Uni-ZAP XR
H0250	Human Activated Monocytes	Human Monocytes				Uni-ZAP XR
H0251	Human Chondrosarcoma	Human Chondrosarcoma	Cartilage		disease	Uni-ZAP XR
H0252	Human Osteosarcoma	Human Osteosarcoma	Bone		disease	Uni-ZAP XR
H0253	Human adult testis, large inserts	Human Adult Testis	Testis			Uni-ZAP XR
H0254	Breast Lymph node cDNA library	Breast Lymph Node	Lymph Node			Uni-ZAP XR
H0255	breast lymph node CDNA library	Breast Lymph Node	Lymph Node			Lambda ZAP II
H0256	HL-60, unstimulated	Human HL-60 Cells, unstimulated	Blood	Cell Line		Uni-ZAP XR

H0257	HL-60, PMA 4H	HL-60 Cells, PMA stimulated 4H	Blood	Cell Line		Uni-ZAP XR
H0261	H. cerebellum, Enzyme subtracted	Human Cerebellum	Brain			Uni-ZAP XR
H0263	human colon cancer	Human Colon Cancer	Colon		disease	Lambda ZAP II
H0264	human tonsils	Human Tonsil	Tonsil			Uni-ZAP XR
H0265	Activated T-Cell (12hs)/Thiouridine labelledEco	T-Cells	Blood	Cell Line		Uni-ZAP XR
H0266	Human Microvascular Endothelial Cells, fract. A	HMEC	Vein	Cell Line		Lambda ZAP II
H0267	Human Microvascular Endothelial Cells, fract. B	HMEC	Vein	Cell Line		Lambda ZAP II
H0268	Human Umbilical Vein Endothelial Cells, fract. A	HUVE Cells	Umbilical vein	Cell Line		Lambda ZAP II
H0269	Human Umbilical Vein Endothelial Cells, fract. B	HUVE Cells	Umbilical vein	Cell Line		Lambda ZAP II
H0270	HPAS (human pancreas, subtracted)	Human Pancreas	Pancreas			Uni-ZAP XR
H0271	Human Neutrophil, Activated	Human Neutrophil - Activated	Blood	Cell Line		Uni-ZAP XR
H0272	HUMAN TONSILS, FRACTION 2	Human Tonsil	Tonsil			Uni-ZAP XR
H0274	Human Adult Spleen, fractionII	Human Adult Spleen	Spleen			Uni-ZAP XR
H0275	Human Infant Adrenal Gland, Subtracted	Human Infant Adrenal Gland	Adrenal gland			pBluescript
H0279	K562 cells	K562 Cell line	cell line	Cell Line		ZAP Express
H0280	K562 + PMA (36 hrs)	K562 Cell line	cell line	Cell Line		ZAP Express
H0281	Lymph node, abnorm. cell line (ATCC #7225)	Lymph Node, abnormal cell line	Lymph Node	Cell Line		ZAP Express
H0282	HBGB's differential consolidation	Human Primary Breast Cancer	Breast			Uni-ZAP XR
H0284	Human OB MG63 control fraction I	Human Osteoblastoma MG63 cell line	Bone	Cell Line		Uni-ZAP XR
H0286	Human OB MG63 treated (10 nM E2) fraction I	Human Osteoblastoma MG63 cell line	Bone	Cell Line		Uni-ZAP XR
H0288	Human OB HOS control fraction I	Human Osteoblastoma HOS cell line	Bone	Cell Line		Uni-ZAP XR
H0290	Human OB HOS treated (1 nM E2) fraction I	Human Osteoblastoma HOS cell line	Bone	Cell Line		Uni-ZAP XR
H0292	Human OB HOS treated (10 nM E2) fraction I	Human Osteoblastoma HOS cell line	Bone	Cell Line		Uni-ZAP XR
H0293	WI 38 cells					Uni-ZAP XR
H0294	Amniotic Cells - TNF induced	Amniotic Cells - TNF induced	Placenta	Cell Line		Uni-ZAP XR
H0295	Amniotic Cells - Primary Culture	Amniotic Cells - Primary Culture	Placenta	Cell Line		Uni-ZAP XR
H0298	HCBB's differential consolidation	CAMA1Ee Cell Line	Breast	Cell Line		Uni-ZAP XR
H0299	HCBA's differential consolidation	CAMA1Ee Cell Line	Breast	Cell Line		Uni-ZAP XR
H0300	CD34 positive cells (Cord Blood)	CD34 Positive Cells	Cord Blood			ZAP Express
H0305	CD34 positive cells (Cord Blood)	CD34 Positive Cells	Cord Blood			ZAP Express
H0306	CD34 depleted Buffy Coat (Cord Blood)	CD34 Depleted Buffy Coat (Cord Blood)	Cord Blood			ZAP Express

H0309	Human Chronic Synovitis	Synovium, Chronic Synovitis/ Osteoarthritis	Synovium		disease	Uni-ZAP XR
H0310	human caudate nucleus	Brain	Brain			Uni-ZAP XR
H0313	human pleural cancer	pleural cancer			disease	pBluescript
H0316	HUMAN STOMACH	Human Stomach	Stomach			Uni-ZAP XR
H0318	HUMAN B CELL LYMPHOMA	Human B Cell Lymphoma	Lymph Node		disease	Uni-ZAP XR
H0320	Human frontal cortex	Human Frontal Cortex	Brain			Uni-ZAP XR
H0321	HUMAN SCHWANOMA	Schwanoma	Nerve		disease	Uni-ZAP XR
H0327	human corpus colosum	Human Corpus Callosum	Brain			Uni-ZAP XR
H0328	human ovarian cancer	Ovarian Cancer	Ovary		disease	Uni-ZAP XR
H0329	Dermatofibrosarcoma Protuberance	Dermatofibrosarcoma Protuberans	Skin		disease	Uni-ZAP XR
H0330	HCBB's Subtractive (- mito genes)	CAMAIEe Cell Line	Breast	Cell Line		Uni-ZAP XR
H0331	Hepatocellular Tumor	Hepatocellular Tumor	Liver		disease	Lambda ZAP II
H0333	Hemangiopericytoma	Hemangiopericytoma	Blood vessel		disease	Lambda ZAP II
H0334	Kidney cancer	Kidney Cancer	Kidney		disease	Uni-ZAP XR
H0339	Duodenum	Duodenum				Uni-ZAP XR
H0340	Corpus Callosum	Corpus Collosum-93052				Uni-ZAP XR
H0341	Bone Marrow Cell Line (RS4;11)	Bone Marrow Cell Line RS4;11	Bone Marrow	Cell Line		Uni-ZAP XR
H0342	Lingual Gyrus	Lingual Gyrus	Brain			Uni-Zap XR
H0343	stomach cancer (human)	Stomach Cancer - 5383A (human)			disease	Uni-ZAP XR
H0344	Adipose tissue (human)	Adipose - 6825A (human)				Uni-ZAP XR
H0345	SKIN	Skin - 4000868H	Skin			Uni-ZAP XR
H0346	Brain-medulloblastoma	Brain (Medulloblastoma)-9405C006R	Brain		disease	Uni-ZAP XR
H0349	human adult liver cDNA library	Human Adult Liver	Liver			pCMVSPORT 1
H0350	Human Fetal Liver, mixed 10 & 14 week	Human Fetal Liver, mixed 10&14 Week	Liver			Uni-ZAP XR
H0351	Glioblastoma	Glioblastoma	Brain		disease	Uni-ZAP XR
H0352	wilm's tumor	Wilm's Tumor			disease	Uni-ZAP XR
H0354	Human Leukocytes	Human Leukocytes	Blood	Cell Line		pCMVSPORT 1
H0355	Human Liver	Human Liver, normal Adult				pCMVSPORT 1
H0356	Human Kidney	Human Kidney	Kidney			pCMVSPORT 1
H0357	H. Normalized Fetal Liver, II	Human Fetal Liver	Liver			Uni-ZAP XR
H0359	KMH2 cell line	KMH2				ZAP Express
H0360	Hemangiopericytoma	Hemangiopericytoma			disease	
H0361	Human rejected kidney	Human Rejected Kidney			disease	pBluescript
H0362	HeLa cell line	HELA CELL LINE				pSport1
H0364	Human Osteoclastoma, excised	Human Osteoclastoma			disease	pBluescript
H0365	Osteoclastoma-normalized B	Human Osteoclastoma			disease	Uni-ZAP XR
H0366	L428 cell line	L428				ZAP Express
H0369	H. Atrophic Endometrium	Atrophic Endometrium and myometrium				Uni-ZAP XR
H0370	H. Lymph node breast Cancer	Lymph node with Met. Breast Cancer			disease	Uni-ZAP XR

H0372	Human Testes	Human Testes	Testis			pCMVSPORT 1
H0373	Human Heart	Human Adult Heart	Heart			pCMVSPORT 1
H0374	Human Brain	Human Brain				pCMVSPORT 1
H0375	Human Lung	Human Lung				pCMVSPORT 1
H0376	Human Spleen	Human Adult Spleen	Spleen			pCMVSPORT 1
H0379	Human Tongue, frac 1	Human Tongue				pSport1
H0380	Human Tongue, frac 2	Human Tongue				pSport1
H0381	Bone Cancer	Bone Cancer			disease	Uni-ZAP XR
H0383	Human Prostate BPH, re-excision	Human Prostate BPH				Uni-ZAP XR
H0384	Brain, Kozak	Human Brain				pCMVSPORT 1
H0386	Leukocyte and Lung; 4 screens	Human Leukocytes	Blood	Cell Line		pCMVSPORT 1
H0388	Human Rejected Kidney, 704 re-excision	Human Rejected Kidney			disease	pBluescript
H0389	H. Brain, X-Chromosome hybridization	Human Brain				pCMVSPORT 1
H0390	Human Amygdala Depression, re-excision	Human Amygdala Depression			disease	pBluescript
H0391	H. Meningioma, M6	Human Meningioma	brain			pSport1
H0392	H. Meningioma, M1	Human Meningioma	brain			pSport1
H0393	Fetal Liver, subtraction II	Human Fetal Liver	Liver			pBluescript
H0394	A-14 cell line	Redd-Sternberg cell				ZAP Express
H0395	A1-CELL LINE	Redd-Sternberg cell				ZAP Express
H0396	L1 Cell line	Redd-Sternberg cell				ZAP Express
H0398	Human Newborn Bladder	Human Newborn Bladder				pBluescript
H0399	Human Kidney Cortex, re-rescue	Human Kidney Cortex				Lambda ZAP II
H0400	Human Striatum Depression, re-rescue	Human Brain, Striatum Depression	Brain			Lambda ZAP II
H0401	Human Pituitary, subtracted V	Human Pituitary				pBluescript
H0402	CD34 depleted Buffy Coat (Cord Blood), re-excision	CD34 Depleted Buffy Coat (Cord Blood)	Cord Blood			ZAP Express
H0404	H. Umbilical Vein endothelial cells, uninduced	HUVE Cells	Umbilical vein	Cell Line		Uni-ZAP XR
H0405	Human Pituitary, subtracted VI	Human Pituitary				pBluescript
H0406	H Amygdala Depression, subtracted	Human Amygdala Depression				Uni-ZAP XR
H0408	Human kidney Cortex, subtracted	Human Kidney Cortex				pBluescript
H0409	H. Striatum Depression, subtracted	Human Brain, Striatum Depression	Brain			pBluescript
H0410	H. Male bladder, adult	H Male Bladder, Adult	Bladder			pSport1
H0411	H Female Bladder, Adult	Human Female Adult Bladder	Bladder			pSport1
H0412	Human umbilical vein endothelial cells, IL-4 induced	HUVE Cells	Umbilical vein	Cell Line		pSport1
H0413	Human Umbilical Vein Endothelial Cells, uninduced	HUVE Cells	Umbilical vein	Cell Line		pSport1
H0414	Ovarian Tumor I, OV5232	Ovarian Tumor, OV5232	Ovary		disease	pSport1
H0415	H. Ovarian Tumor, II, OV5232	Ovarian Tumor, OV5232	Ovary		disease	pCMVSPORT 2.0
H0416	Human Neutrophils, Activated, re-excision	Human Neutrophil - Activated	Blood	Cell Line		pBluescript
H0417	Human Pituitary,	Human Pituitary				pBluescript

	subtracted VIII					
H0418	Human Pituitary, subtracted VII	Human Pituitary				pBluescript
H0419	Bone Cancer, re-excision	Bone Cancer				Uni-ZAP XR
H0421	Human Bone Marrow, re-excision	Bone Marrow				pBluescript
H0422	T-Cell PHA 16 hrs	T-Cells	Blood	Cell Line		pSport1
H0423	T-Cell PHA 24 hrs	T-Cells	Blood	Cell Line		pSport1
H0424	Human Pituitary, subt IX	Human Pituitary				pBluescript
H0427	Human Adipose	Human Adipose, left hiplipoma				pSport1
H0428	Human Ovary	Human Ovary Tumor	Ovary			pSport1
H0429	K562 + PMA (36 hrs), re-excision	K562 Cell line	cell line	Cell Line		ZAP Express
H0431	H. Kidney Medulla, re-excision	Kidney medulla	Kidney			pBluescript
H0432	H. Kidney Pyramid	Kidney pyramids	Kidney			pBluescript
H0433	Human Umbilical Vein Endothelial cells, frac B, re-excision	HUVE Cells	Umbilical vein	Cell Line		pBluescript
H0434	Human Brain, striatum, re-excision	Human Brain, Striatum				pBluescript
H0435	Ovarian Tumor 10-3-95	Ovarian Tumor, OV350721	Ovary			pCMVSPORT 2.0
H0436	Resting T-Cell Library, II	T-Cells	Blood	Cell Line		pSport1
H0437	H Umbilical Vein Endothelial Cells, frac A, re-excision	HUVE Cells	Umbilical vein	Cell Line		Lambda ZAP II
H0438	H. Whole Brain #2, re-excision	Human Whole Brain #2				ZAP Express
H0439	Human Eosinophils	Eosinophils				pBluescript
H0440	FGF enriched mixed library	Mixed libraries				pCMVSPORT 1
H0441	H. Kidney Cortex, subtracted	Kidney cortex	Kidney			pBluescript
H0442	H. Striatum Depression, subt II	Human Brain, Striatum Depression	Brain			pBluescript
H0443	H. Adipose, subtracted	Human Adipose, left hiplipoma				pSport1
H0444	Spleen metastatic melanoma	Spleen, Metastatic malignant melanoma	Spleen		disease	pSport1
H0445	Spleen, Chronic lymphocytic leukemia	Human Spleen, CLL	Spleen		disease	pSport1
H0447	Salivary gland, re-excision	Human Salivary Gland	Salivary gland			Uni-ZAP XR
H0448	Salivary gland, subtracted	Human Salivary Gland	Salivary gland			Lambda ZAP II
H0449	CD34+ cell, I	CD34 positive cells				pSport1
H0450	CD34+cells, II	CD34 positive cells				pCMVSPORT 2.0
H0453	H. Kidney Pyramid, subtracted	Kidney pyramids	Kidney			pBluescript
H0455	H. Striatum Depression, subt	Human Brain, Striatum Depression	Brain			pBluescript
H0456	H Kidney Cortex, subtracted III	Human Kidney Cortex				pBluescript
H0457	Human Eosinophils	Human Eosinophils				pSport1
H0458	CD34+ cell, I, frac II	CD34 positive cells				pSport1
H0459	CD34+cells, II, FRACTION 2	CD34 positive cells				pCMVSPORT 2.0
H0461	H. Kidney Medulla, subtracted	Kidney medulla	Kidney			pBluescript
H0462	H. Amygdala Depression, subtracted		Brain			pBluescript

H0477	Human Tonsil, Lib 3	Human Tonsil	Tonsil			pSport1
H0478	Salivary Gland, Lib 2	Human Salivary Gland	Salivary gland			pSport1
H0479	Salivary Gland, Lib 3	Human Salivary Gland	Salivary gland			pSport1
H0480	L8 cell line	L8 cell line				ZAP Express
H0483	Breast Cancer cell line, MDA 36	Breast Cancer Cell line, MDA 36				pSport1
H0484	Breast Cancer Cell line, angiogenic	Breast Cancer Cell line, Angiogenic, 36T3				pSport1
H0485	Hodgkin's Lymphoma I	Hodgkin's Lymphoma I			disease	pCMVSPORT 2.0
H0486	Hodgkin's Lymphoma II	Hodgkin's Lymphoma II			disease	pCMVSPORT 2.0
H0487	Human Tonsils, lib 1	Human Tonsils				pCMVSPORT 2.0
H0488	Human Tonsils, Lib 2	Human Tonsils				pCMVSPORT 2.0
H0489	Crohn's Disease	Ileum	Intestine		disease	pSport1
H0490	HL-60, untreated, subtracted	Human HL-60 Cells, unstimulated	Blood	Cell Line		Uni-ZAP XR
H0491	HL-60, PMA 4H, subtracted	HL-60 Cells, PMA stimulated 4H	Blood	Cell Line		Uni-ZAP XR
H0492	HL-60, RA 4h, Subtracted	HL-60 Cells, RA stimulated for 4H	Blood	Cell Line		Uni-ZAP XR
H0494	Keratinocyte	Keratinocyte				pCMVSPORT 2.0
H0497	HEL cell line	HEL cell line		HEL 92.1.7		pSport1
H0505	Human Astrocyte	Human Astrocyte				pSport1
H0506	Ulcerative Colitis	Colon	Colon			pSport1
H0509	Liver, Hepatoma	Human Liver, Hepatoma, patient 8	Liver		disease	pCMVSPORT 3.0
H0510	Human Liver, normal	Human Liver, normal, Patient # 8	Liver			pCMVSPORT 3.0
H0512	Keratinocyte, lib 3	Keratinocyte				pCMVSPORT 2.0
H0517	Nasal polyps	Nasal polyps				pCMVSPORT 2.0
H0518	pBMC stimulated w/ poly I/C	pBMC stimulated with poly I/C				pCMVSPORT 3.0
H0519	NTERA2, control	NTERA2, Teratocarcinoma cell line				pCMVSPORT 3.0
H0520	NTERA2 + retinoic acid, 14 days	NTERA2, Teratocarcinoma cell line				pSport1
H0521	Primary Dendritic Cells, lib 1	Primary Dendritic cells				pCMVSPORT 3.0
H0522	Primary Dendritic cells, frac 2	Primary Dendritic cells				pCMVSPORT 3.0
H0523	Primary Dendritic cells, CapFinder2, frac 1	Primary Dendritic cells				pSport1
H0524	Primary Dendritic Cells, CapFinder, frac 2	Primary Dendritic cells				pSport1
H0525	PCR, pBMC I/C treated	pBMC stimulated with poly I/C				PCR II
H0528	Poly[I]/Poly[C] Normal Lung Fibroblasts	Poly[I]/Poly[C] Normal Lung Fibroblasts				pCMVSPORT 3.0
H0529	Myeloid Progenitor Cell Line	TF-1 Cell Line; Myeloid progenitor cell line				pCMVSPORT 3.0
H0530	Human Dermal Endothelial Cells, untreated	Human Dermal Endothelial Cells; untreated				pSport1
H0535	Human ovary tumor cell OV350721	Ovarian Tumor, OV350721	Ovary		disease	pSport1
H0537	H. Primary Dendritic	Primary Dendritic				pCMVSPORT 2.0

	Cells,lib 3	cells				
H0538	Merkel Cells	Merkel cells	Lymph node			pSport1
H0539	Pancreas Islet Cell Tumor	Pancreas Islet Cell Tumour	Pancreas		disease	pSport1
H0540	Skin, burned	Skin, leg burned	Skin			pSport1
H0542	T Cell helper I	Helper T cell				pCMVSPORT 3.0
H0543	T cell helper II	Helper T cell				pCMVSPORT 3.0
H0544	Human endometrial stromal cells	Human endometrial stromal cells				pCMVSPORT 3.0
H0545	Human endometrial stromal cells-treated with progesterone	Human endometrial stromal cells-treated with proge				pCMVSPORT 3.0
H0546	Human endometrial stromal cells-treated with estradiol	Human endometrial stromal cells-treated with estra				pCMVSPORT 3.0
H0547	NTERA2 teratocarcinoma cell line+retinoic acid (14 days)	NTERA2, Teratocarcinoma cell line				pSport1
H0549	H. Epididymus, caput & corpus	Human Epididymus, caput and corpus				Uni-ZAP XR
H0550	H. Epididymus, cauda	Human Epididymus, cauda				Uni-ZAP XR
H0551	Human Thymus Stromal Cells	Human Thymus Stromal Cells				pCMVSPORT 3.0
H0552	Signal trap,Femur Bone Marrow,pooled	Femur Bone marrow, pooled from 8 male/female				Other
H0553	Human Placenta	Human Placenta				pCMVSPORT 3.0
H0555	Rejected Kidney, lib 4	Human Rejected Kidney	Kidney		disease	pCMVSPORT 3.0
H0556	Activated T-cell(12h)/Thiouridine-re-excision	T-Cells	Blood	Cell Line		Uni-ZAP XR
H0559	HL-60, PMA 4H, re-excision	HL-60 Cells, PMA stimulated 4H	Blood	Cell Line		Uni-ZAP XR
H0560	KMH2	KMH2				pCMVSPORT 3.0
H0561	L428	L428				pCMVSPORT 3.0
H0562	Human Fetal Brain, normalized c5-11-26	Human Fetal Brain				pCMVSPORT 2.0
H0563	Human Fetal Brain, normalized 50021F	Human Fetal Brain				pCMVSPORT 2.0
H0564	Human Fetal Brain, normalized C5001F	Human Fetal Brain				pCMVSPORT 2.0
H0565	HUMAN Fetal Brain, normalized 100024F	Human Fetal Brain				pCMVSPORT 2.0
H0566	Human Fetal Brain,normalized c50F	Human Fetal Brain				pCMVSPORT 2.0
H0567	Human Fetal Brain, normalized A5002F	Human Fetal Brain				pCMVSPORT 2.0
H0569	Human Fetal Brain, normalized CO	Human Fetal Brain				pCMVSPORT 2.0
H0570	Human Fetal Brain, normalized C500H	Human Fetal Brain				pCMVSPORT 2.0
H0571	Human Fetal Brain, normalized C500HE	Human Fetal Brain				pCMVSPORT 2.0
H0572	Human Fetal Brain, normalized AC5002	Human Fetal Brain				pCMVSPORT 2.0
H0574	Hepatocellular Tumor; re-excision	Hepatocellular Tumor	Liver		disease	Lambda ZAP II
H0575	Human Adult Pulmonary;re-excision	Human Adult Pulmonary	Lung			Uni-ZAP XR
H0576	Resting T-Cell; re-excision	T-Cells	Blood	Cell Line		Lambda ZAP II
H0578	Human Fetal Thymus	Fetal Thymus	Thymus			pSport1

H0579	Pericardium	Pericardium	Heart			pSport1
H0580	Dendritic cells, pooled	Pooled dendritic cells				pCMVSPORT 3.0
H0581	Human Bone Marrow, treated	Human Bone Marrow	Bone Marrow			pCMVSPORT 3.0
H0583	B Cell lymphoma	B Cell Lymphoma	B Cell		disease	pCMVSPORT 3.0
H0584	Activated T-cells, 24 hrs, re-excision	Activated T-Cells	Blood	Cell Line		Uni-ZAP XR
H0585	Activated T-Cells, 12 hrs, re-excision	Activated T-Cells	Blood	Cell Line		Uni-ZAP XR
H0586	Healing groin wound, 6.5 hours post incision	healing groin wound, 6.5 hours post incision - 2/	groin		disease	pCMVSPORT 3.0
H0587	Healing groin wound; 7.5 hours post incision	Groin-2/19/97	groin		disease	pCMVSPORT 3.0
H0589	CD34 positive cells (cord blood), re-ex	CD34 Positive Cells	Cord Blood			ZAP Express
H0590	Human adult small intestine, re-excision	Human Adult Small Intestine	Small Int.			Uni-ZAP XR
H0591	Human T-cell lymphoma; re-excision	T-Cell Lymphoma	T-Cell		disease	Uni-ZAP XR
H0592	Healing groin wound - zero hr post-incision (control)	HGS wound healing project; abdomen			disease	pCMVSPORT 3.0
H0593	Olfactory epithelium; nasal cavity	Olfactory epithelium from roof of left nasal cavity				pCMVSPORT 3.0
H0594	Human Lung Cancer; re-excision	Human Lung Cancer	Lung		disease	Lambda ZAP II
H0595	Stomach cancer (human); re-excision	Stomach Cancer - 5383A (human)			disease	Uni-ZAP XR
H0596	Human Colon Cancer; re-excision	Human Colon Cancer	Colon			Lambda ZAP II
H0597	Human Colon; re-excision	Human Colon				Lambda ZAP II
H0598	Human Stomach; re-excision	Human Stomach	Stomach			Uni-ZAP XR
H0599	Human Adult Heart; re-excision	Human Adult Heart	Heart			Uni-ZAP XR
H0600	Healing Abdomen wound; 70&90 min post incision	Abdomen			disease	pCMVSPORT 3.0
H0601	Healing Abdomen Wound; 15 days post incision	Abdomen			disease	pCMVSPORT 3.0
H0602	Healing Abdomen Wound; 21&29 days post incision	Abdomen			disease	pCMVSPORT 3.0
H0604	Human Pituitary, re-excision	Human Pituitary				pBluescript
H0606	Human Primary Breast Cancer; re-excision	Human Primary Breast Cancer	Breast		disease	Uni-ZAP XR
H0607	H. Leukocytes, normalized cot 50A3	H. Leukocytes				pCMVSPORT 1
H0608	H. Leukocytes, control	H. Leukocytes				pCMVSPORT 1
H0609	H. Leukocytes, normalized cot > 500A	H. Leukocytes				pCMVSPORT 1
H0610	H. Leukocytes, normalized cot 5A	H. Leukocytes				pCMVSPORT 1
H0611	H. Leukocytes, normalized cot 500 B	H. Leukocytes				pCMVSPORT 1
H0612	H. Leukocytes, normalized cot 50 B	H. Leukocytes				pCMVSPORT 1
H0613	H. Leukocytes, normalized cot 5B	H. Leukocytes				pCMVSPORT 1
H0614	H. Leukocytes,	H. Leukocytes				pCMVSPORT 1

	normalized cot 500 A					
H0615	Human Ovarian Cancer Reexcision	Ovarian Cancer	Ovary		disease	Uni-ZAP XR
H0616	Human Testes, Reexcision	Human Testes	Testis			Uni-ZAP XR
H0617	Human Primary Breast Cancer Reexcision	Human Primary Breast Cancer	Breast		disease	Uni-ZAP XR
H0618	Human Adult Testes, Large Inserts, Reexcision	Human Adult Testis	Testis			Uni-ZAP XR
H0619	Fetal Heart	Human Fetal Heart	Heart			Uni-ZAP XR
H0620	Human Fetal Kidney; Reexcision	Human Fetal Kidney	Kidney			Uni-ZAP XR
H0622	Human Pancreas Tumor; Reexcision	Human Pancreas Tumor	Pancreas		disease	Uni-ZAP XR
H0623	Human Umbilical Vein; Reexcision	Human Umbilical Vein Endothelial Cells	Umbilical vein			Uni-ZAP XR
H0624	12 Week Early Stage Human II; Reexcision	Twelve Week Old Early Stage Human	Embryo			Uni-ZAP XR
H0625	Ku 812F Basophils Line	Ku 812F Basophils				pSport1
H0626	Saos2 Cells; Untreated	Saos2 Cell Line; Untreated				pSport1
H0627	Saos2 Cells; Vitamin D3 Treated	Saos2 Cell Line; Vitamin D3 Treated				pSport1
H0628	Human Pre-Differentiated Adipocytes	Human Pre-Differentiated Adipocytes				Uni-ZAP XR
H0629	Human Leukocyte, control #2	Human Normalized leukocyte				pCMVSPORT 1
H0630	Human Leukocytes, normalized control #4	Human Normalized leukocyte				pCMVSPORT 1
H0631	Saos2, Dexamethosome Treated	Saos2 Cell Line; Dexamethosome Treated				pSport1
H0632	Hepatocellular Tumor; re-excision	Hepatocellular Tumor	Liver			Lambda ZAP II
H0633	Lung Carcinoma A549 TNFalpha activated	TNFalpha activated A549--Lung Carcinoma			disease	pSport1
H0634	Human Testes Tumor, re-excision	Human Testes Tumor	Testis		disease	Uni-ZAP XR
H0635	Human Activated T-Cells, re-excision	Activated T-Cells	Blood	Cell Line		Uni-ZAP XR
H0637	Dendritic Cells From CD34 Cells	Dendritic cells from CD34 cells				pSport1
H0638	CD40 activated monocyte dendritic cells	CD40 activated monocyte dendritic cells				pSport1
H0639	Ficoll Human Stromal Cells, 5Fu treated	Ficoll Human Stromal Cells, 5Fu treated				Other
H0640	Ficoll Human Stromal Cells, Untreated	Ficoll Human Stromal Cells, Untreated				Other
H0641	LPS activated derived dendritic cells	LPS activated monocyte derived dendritic cells				pSport1
H0642	Hep G2 Cells, lambda library	Hep G2 Cells				Other
H0643	Hep G2 Cells, PCR library	Hep G2 Cells				Other
H0644	Human Placenta (re-excision)	Human Placenta	Placenta			Uni-ZAP XR
H0645	Fetal Heart, re-excision	Human Fetal Heart	Heart			Uni-ZAP XR
H0646	Lung, Cancer (4005313 A3): Invasive Poorly	Metastatic squamous cell lung				pSport1

	Differentiated Lung Adenocarcinoma,	carcinoma, poorly di				
H0647	Lung, Cancer (4005163 B7): Invasive, Poorly Diff. Adenocarcinoma, Metastatic	Invasive poorly differentiated lung adenocarcinoma			disease	pSport1
H0648	Ovary, Cancer: (4004562 B6) Papillary Serous Cystic Neoplasm, Low Malignant Pot	Papillary Cstic neoplasm of low malignant potentia			disease	pSport1
H0649	Lung, Normal: (4005313 B1)	Normal Lung				pSport1
H0650	B-Cells	B-Cells				pCMVSPORT 3.0
H0651	Ovary, Normal: (9805C040R)	Normal Ovary				pSport1
H0652	Lung, Normal: (4005313 B1)	Normal Lung				pSport1
H0653	Stromal Cells	Stromal Cells				pSport1
H0654	Lung, Cancer: (4005313 A3) Invasive Poorly-differentiated Metastatic lung adenoc	Metastatic Squamous cell lung Carcinoma poorly dif				Other
H0656	B-cells (unstimulated)	B-cells (unstimulated)				pSport1
H0657	B-cells (stimulated)	B-cells (stimulated)				pSport1
H0658	Ovary, Cancer (9809C332): Poorly differentiated adenocarcinoma	9809C332- Poorly differentiate	Ovary & Fallopian Tubes		disease	pSport1
H0659	Ovary, Cancer (15395A1F): Grade II Papillary Carcinoma	Grade II Papillary Carcinoma, Ovary	Ovary		disease	pSport1
H0660	Ovary, Cancer: (15799A1F) Poorly differentiated carcinoma	Poorly differentiated carcinoma, ovary			disease	pSport1
H0661	Breast, Cancer: (4004943 A5)	Breast cancer			disease	pSport1
H0662	Breast, Normal: (4005522B2)	Normal Breast - #4005522(B2)	Breast			pSport1
H0663	Breast, Cancer: (4005522 A2)	Breast Cancer - #4005522(A2)	Breast		disease	pSport1
H0664	Breast, Cancer: (9806C012R)	Breast Cancer	Breast		disease	pSport1
H0665	Stromal cells 3.88	Stromal cells 3.88				pSport1
H0666	Ovary, Cancer: (4004332 A2)	Ovarian Cancer, Sample #4004332A2			disease	pSport1
H0667	Stromal cells(HBM3.18)	Stromal cell(HBM 3.18)				pSport1
H0668	stromal cell clone 2.5	stromal cell clone 2.5				pSport1
H0669	Breast, Cancer: (4005385 A2)	Breast Cancer (4005385A2)	Breast			pSport1
H0670	Ovary, Cancer(4004650 A3): Well-Differentiated Micropapillary Serous Carcinoma	Ovarian Cancer - 4004650A3				pSport1
H0671	Breast, Cancer: (9802C02OE)	Breast Cancer-Sample # 9802C02OE				pSport1
H0672	Ovary, Cancer: (4004576 A8)	Ovarian Cancer(4004576A8)	Ovary			pSport1
H0673	Human Prostate Cancer, Stage B2; re-excision	Human Prostate Cancer, stage B2	Prostate			Uni-ZAP XR
H0674	Human Prostate Cancer,	Human Prostate	Prostate			Uni-ZAP XR

	Stage C; re-excision	Cancer, stage C				
H0675	Colon, Cancer: (9808C064R)	Colon Cancer 9808C064R				pCMVSPORT 3.0
H0676	Colon, Cancer: (9808C064R)-total RNA	Colon Cancer 9808C064R				pCMVSPORT 3.0
H0677	TNFR degenerate oligo	B-Cells				PCR II
H0678	screened clones from placental library	Placenta	Placenta			Other
H0679	screened clones from Tonsil library	Human Tonsils				Other
H0682	Serous Papillary Adenocarcinoma	serous papillary adenocarcinoma (9606G304SPA3B)				pCMVSPORT 3.0
H0683	Ovarian Serous Papillary Adenocarcinoma	Serous papillary adenocarcinoma, stage 3C (9804G01)				pCMVSPORT 3.0
H0684	Serous Papillary Adenocarcinoma	Ovarian Cancer- 9810G606	Ovaries			pCMVSPORT 3.0
H0685	Adenocarcinoma of Ovary, Human Cell Line, # OVCAR-3	Adenocarcinoma of Ovary, Human Cell Line, # OVCAR-				pCMVSPORT 3.0
H0686	Adenocarcinoma of Ovary, Human Cell Line	Adenocarcinoma of Ovary, Human Cell Line, # SW-626				pCMVSPORT 3.0
H0687	Human normal ovary(#9610G215)	Human normal ovary(#9610G215)	Ovary			pCMVSPORT 3.0
H0688	Human Ovarian Cancer(#9807G017)	Human Ovarian cancer(#9807G017), mRNA from Maura Ru				pCMVSPORT 3.0
H0689	Ovarian Cancer	Ovarian Cancer, #9806G019				pCMVSPORT 3.0
H0690	Ovarian Cancer, # 9702G001	Ovarian Cancer, #9702G001				pCMVSPORT 3.0
H0691	Normal Ovary, #9710G208	normal ovary, #9710G208				pCMVSPORT 3.0
H0692	BLyS Receptor from Expression Cloning	B Cell Lymphoma	B Cell			pCMVSPORT 3.0
H0693	Normal Prostate #ODQ3958EN	Normal Prostate Tissue # ODQ3958EN				pCMVSPORT 3.0
H0694	Prostate gland adenocarcinoma	Prostate gland, adenocarcinoma, mod/diff, gleason	prostate gland			pCMVSPORT 3.0
H0695	mononucleocytes from patient	mononucleocytes from patient at Shady Grove Hospit				pCMVSPORT 3.0
N0006	Human Fetal Brain	Human Fetal Brain				
N0007	Human Hippocampus	Human Hippocampus				
N0009	Human Hippocampus, prescreened	Human Hippocampus				
N0011	Human Brain	Human Brain				
S0001	Brain frontal cortex	Brain frontal cortex	Brain			Lambda ZAP II
S0002	Monocyte activated	Monocyte-activated	blood	Cell Line		Uni-ZAP XR
S0003	Human Osteoclastoma	Osteoclastoma	bone		disease	Uni-ZAP XR
S0004	Prostate	Prostate BPH	Prostate			Lambda ZAP II
S0005	Heart	Heart-left ventricle	Heart			pCDNA
S0006	Neuroblastoma	Human Neural Blastoma			disease	pCDNA
S0007	Early Stage Human Brain	Human Fetal Brain				Uni-ZAP XR
S0010	Human Amygdala	Amygdala				Uni-ZAP XR
S0011	STROMAL - OSTEOCLASTOMA	Osteoclastoma	bone		disease	Uni-ZAP XR
S0013	Prostate	Prostate	prostate			Uni-ZAP XR

S0014	Kidney Cortex	Kidney cortex	Kidney			Uni-ZAP XR
S0015	Kidney medulla	Kidney medulla	Kidney			Uni-ZAP XR
S0016	Kidney Pyramids	Kidney pyramids	Kidney			Uni-ZAP XR
S0021	Whole brain	Whole brain	Brain			ZAP Express
S0022	Human Osteoclastoma Stromal Cells - unamplified	Osteoclastoma Stromal Cells				Uni-ZAP XR
S0024	Human Kidney Medulla - unamplified	Human Kidney Medulla				
S0025	Human Kidney Pyramids - unamplified	Human Kidney Pyramids				
S0026	Stromal cell TF274	stromal cell	Bone marrow	Cell Line		Uni-ZAP XR
S0027	Smooth muscle, serum treated	Smooth muscle	Pulmonary artery	Cell Line		Uni-ZAP XR
S0028	Smooth muscle, control	Smooth muscle	Pulmonary artery	Cell Line		Uni-ZAP XR
S0029	brain stem	Brain stem	brain			Uni-ZAP XR
S0030	Brain pons	Brain Pons	Brain			Uni-ZAP XR
S0031	Spinal cord	Spinal cord	spinal cord			Uni-ZAP XR
S0032	Smooth muscle-ILb induced	Smooth muscle	Pulmonary artery	Cell Line		Uni-ZAP XR
S0035	Brain medulla oblongata	Brain medulla oblongata	Brain			Uni-ZAP XR
S0036	Human Substantia Nigra	Human Substantia Nigra				Uni-ZAP XR
S0037	Smooth muscle, IL1b induced	Smooth muscle	Pulmonary artery	Cell Line		Uni-ZAP XR
S0038	Human Whole Brain #2 - Oligo dT > 1.5Kb	Human Whole Brain #2				ZAP Express
S0039	Hypothalamus	Hypothalamus	Brain			Uni-ZAP XR
S0040	Adipocytes	Human Adipocytes from Osteoclastoma				Uni-ZAP XR
S0041	Thalamus	Human Thalamus				Uni-ZAP XR
S0042	Testes	Human Testes				ZAP Express
S0044	Prostate BPH	prostate BPH	Prostate		disease	Uni-ZAP XR
S0045	Endothelial cells-control	Endothelial cell	endothelial cell-lung	Cell Line		Uni-ZAP XR
S0046	Endothelial-induced	Endothelial cell	endothelial cell-lung	Cell Line		Uni-ZAP XR
S0048	Human Hypothalamus, Alzheimer's	Human Hypothalamus, Alzheimer's			disease	Uni-ZAP XR
S0049	Human Brain, Striatum	Human Brain, Striatum				Uni-ZAP XR
S0050	Human Frontal Cortex, Schizophrenia	Human Frontal Cortex, Schizophrenia			disease	Uni-ZAP XR
S0051	Human Hypothalamus, Schizophrenia	Human Hypothalamus, Schizophrenia			disease	Uni-ZAP XR
S0052	neutrophils control	human neutrophils	blood	Cell Line		Uni-ZAP XR
S0053	Neutrophils IL-1 and LPS induced	human neutrophil induced	blood	Cell Line		Uni-ZAP XR
S0106	STRIATUM DEPRESSION		BRAIN		disease	Uni-ZAP XR
S0110	Brain Amygdala Depression		Brain		disease	Uni-ZAP XR
S0112	Hypothalamus		Brain			Uni-ZAP XR
S0114	Anergic T-cell	Anergic T-cell		Cell Line		Uni-ZAP XR
S0116	Bone marrow	Bone marrow	Bone marrow			Uni-ZAP XR
S0118	Smooth muscle control 2	Smooth muscle	Pulmonary artery	Cell Line		Uni-ZAP XR
S0122	Osteoclastoma-normalized	Osteoclastoma	bone		disease	pBluescript

	A					
S0124	Smooth muscle-edited A	Smooth muscle	Pulmonary artery	Cell Line		Uni-ZAP XR
S0126	Osteoblasts	Osteoblasts	Knee	Cell Line		Uni-ZAP XR
S0132	Epithelial-TNF α and INF induced	Airway Epithelial				Uni-ZAP XR
S0134	Apoptotic T-cell	apoptotic cells		Cell Line		Uni-ZAP XR
S0136	PERM TF274	stromal cell	Bone marrow	Cell Line		Lambda ZAP II
S0140	eosinophil-IL5 induced	eosinophil	lung	Cell Line		Uni-ZAP XR
S0142	Macrophage-oxLDL	macrophage-oxidized LDL treated	blood	Cell Line		Uni-ZAP XR
S0144	Macrophage (GM-CSF treated)	Macrophage (GM-CSF treated)				Uni-ZAP XR
S0146	prostate-edited	prostate BPH	Prostate			Uni-ZAP XR
S0148	Normal Prostate	Prostate	prostate			Uni-ZAP XR
S0150	LNCAP prostate cell line	LNCAP Cell Line	Prostate	Cell Line		Uni-ZAP XR
S0152	PC3 Prostate cell line	PC3 prostate cell line				Uni-ZAP XR
S0168	Prostate/LNCAP, subtraction I	PC3 prostate cell line				pBluescript
S0174	Prostate-BPH subtracted II	Human Prostate BPH				pBluescript
S0176	Prostate, normal, subtraction I	Prostate	prostate			Uni-ZAP XR
S0180	Bone Marrow Stroma, TNF&LPS ind	Bone Marrow Stroma, TNF & LPS induced			disease	Uni-ZAP XR
S0182	Human B Cell 8866	Human B- Cell 8866				Uni-ZAP XR
S0184	7TM Receptor enriched, lib II	PBLs, 7TM receptor enriched				Other
S0188	Prostate,BPH, Lib 2	Human Prostate BPH			disease	pSport1
S0190	Prostate BPH,Lib 2, subtracted	Human Prostate BPH				pSport1
S0192	Synovial Fibroblasts (control)	Synovial Fibroblasts				pSport1
S0194	Synovial hypoxia	Synovial Fibroblasts				pSport1
S0196	Synovial IL-1/TNF stimulated	Synovial Fibroblasts				pSport1
S0202	7TM-pbdd	PBLs, 7TM receptor enriched				PCRII
S0206	Smooth Muscle- HASTE normalized	Smooth muscle	Pulmonary artery	Cell Line		pBluescript
S0208	Mesangial cell, frac 1	Mesangial cell				pSport1
S0210	Mesangial cell, frac 2	Mesangial cell				pSport1
S0212	Bone Marrow Stromal Cell, untreated	Bone Marrow Stromal Cell,untreated				pSport1
S0214	Human Osteoclastoma, re-excision	Osteoclastoma	bone		disease	Uni-ZAP XR
S0216	Neutrophils IL-1 and LPS induced	human neutrophil induced	blood	Cell Line		Uni-ZAP XR
S0218	Apoptotic T-cell, re-excision	apoptotic cells		Cell Line		Uni-ZAP XR
S0220	H. hypothalamus, frac A;re-excision	Hypothalamus	Brain			ZAP Express
S0222	H. Frontal cortex,epileptic;re-excision	H. Brain, Frontal Cortex, Epileptic	Brain		disease	Uni-ZAP XR
S0242	Synovial Fibroblasts (II/TNF), subt	Synovial Fibroblasts				pSport1
S0250	Human Osteoblasts II	Human Osteoblasts	Femur		disease	pCMVSPORT 2.0
S0252	7TM-PIMIX	PBLs, 7TM				PCRII

		receptor enriched				
S0256	7TM-PHMIX	PBLS, 7TM receptor enriched				PCRII
S0260	Spinal Cord, re-excision	Spinal cord	spinal cord			Uni-ZAP XR
S0264	PPMIX	PPMIX (Human Pituitary)	Pituitary			PCRII
S0266	PLMIX	PLMIX (Human Lung)	Lung			PCRII
S0268	PRMIX	PRMIX (Human Prostate)	prostate			PCRII
S0270	PTMIX	PTMIX (Human Thymus)	Thymus			PCRII
S0276	Synovial hypoxia-RSF subtracted	Synovial fobroblasts (rheumatoid)	Synovial tissue			pSport1
S0278	H Macrophage (GM-CSF treated), re-excision	Macrophage (GM- CSF treated)				Uni-ZAP XR
S0280	Human Adipose Tissue, re-excision	Human Adipose Tissue				Uni-ZAP XR
S0282	Brain Frontal Cortex, re- excision	Brain frontal cortex	Brain			Lambda ZAP II
S0292	Osteoarthritis (OA-4)	Human Osteoarthritic Cartilage	Bone		disease	pSport1
S0294	Larynx tumor	Larynx tumor	Larynx,voca l cord		disease	pSport1
S0296	Normal lung	Normal lung	Lung			pSport1
S0298	Bone marrow stroma,treated	Bone marrow stroma,treatedSB	Bone marrow			pSport1
S0300	Frontal lobe,dementia;re- excision	Frontal Lobe dementia/Alzheimer' 's	Brain			Uni-ZAP XR
S0306	Larynx normal #10 261- 273	Larynx normal				pSport1
S0308	Spleen/normal	Spleen normal				pSport1
S0310	Normal trachea	Normal trachea				pSport1
S0312	Human osteoarthritic;fraction II	Human osteoarthritic cartilage			disease	pSport1
S0314	Human osteoarthritic;fraction I	Human osteoarthritic cartilage			disease	pSport1
S0316	Human Normal Cartilage,Fraction I	Human Normal Cartilage				pSport1
S0318	Human Normal Cartilage Fraction II	Human Normal Cartilage				pSport1
S0320	Human Larynx	Larynx	Epiglottis			pSport1
S0324	Human Brain	Brain	Cerebellum			pSport1
S0328	Palate carcinoma	Palate carcinoma	Uvula		disease	pSport1
S0330	Palate normal	Palate normal	Uvula			pSport1
S0332	Pharynx carcinoma	Pharynx carcinoma	Hypopharyn x			pSport1
S0334	Human Normal Cartilage Fraction III	Human Normal Cartilage				pSport1
S0336	Human Normal Cartilage Fraction IV	Human Normal Cartilage				pSport1
S0338	Human Osteoarthritic Cartilage Fraction III	Human osteoarthritic cartilage			disease	pSport1
S0340	Human Osteoarthritic Cartilage Fraction IV	Human osteoarthritic cartilage			disease	pSport1
S0342	Adipocytes;re-excision	Human Adipocytes from Osteoclastoma				Uni-ZAP XR
S0344	Macrophage-oxLDL; re- excision	macrophage- oxidized LDL	blood	Cell Line		Uni-ZAP XR

		treated				
S0346	Human Amygdala;re-excision	Amygdala				Uni-ZAP XR
S0348	Cheek Carcinoma	Cheek Carcinoma			disease	pSport1
S0350	Pharynx Carcinoma	Pharynx carcinoma	Hypopharynx		disease	pSport1
S0352	Larynx Carcinoma	Larynx carcinoma			disease	pSport1
S0354	Colon Normal II	Colon Normal	Colon			pSport1
S0356	Colon Carcinoma	Colon Carcinoma	Colon		disease	pSport1
S0358	Colon Normal III	Colon Normal	Colon			pSport1
S0360	Colon Tumor II	Colon Tumor	Colon		disease	pSport1
S0362	Human Gastrocnemius	Gastrocnemius muscle				pSport1
S0364	Human Quadriceps	Quadriceps muscle				pSport1
S0366	Human Soleus	Soleus Muscle				pSport1
S0368	Human Pancreatic Langerhans	Islets of Langerhans				pSport1
S0370	Larynx carcinoma II	Larynx carcinoma			disease	pSport1
S0372	Larynx carcinoma III	Larynx carcinoma			disease	pSport1
S0374	Normal colon	Normal colon				pSport1
S0376	Colon Tumor	Colon Tumor			disease	pSport1
S0378	Pancreas normal PCA4 No	Pancreas Normal PCA4 No				pSport1
S0380	Pancreas Tumor PCA4 Tu	Pancreas Tumor PCA4 Tu			disease	pSport1
S0382	Larynx carcinoma IV	Larynx carcinoma			disease	pSport1
S0384	Tongue carcinoma	Tongue carcinoma			disease	pSport1
S0386	Human Whole Brain, re-excision	Whole brain	Brain			ZAP Express
S0388	Human Hypothalamus,schizophrenia, re-excision	Human Hypothalamus, Schizophrenia			disease	Uni-ZAP XR
S0390	Smooth muscle, control; re-excision	Smooth muscle	Pulmonary artery	Cell Line		Uni-ZAP XR
S0392	Salivary Gland	Salivary gland; normal				pSport1
S0394	Stomach;normal	Stomach; normal				pSport1
S0396	Uterus; normal	Uterus; normal				pSport1
S0398	Testis; normal	Testis; normal				pSport1
S0400	Brain; normal	Brain; normal				pSport1
S0402	Adrenal Gland,normal	Adrenal gland; normal				pSport1
S0404	Rectum normal	Rectum, normal				pSport1
S0406	Rectum tumour	Rectum tumour				pSport1
S0408	Colon, normal	Colon, normal				pSport1
S0410	Colon, tumour	Colon, tumour				pSport1
S0412	Temporal cortex-Alzheimer; subtracted	Temporal cortex, alzheimer			disease	Other
S0414	Hippocampus, Alzheimer Subtracted	Hippocampus, Alzheimer Subtracted				Other
S0418	CHME Cell Line;treated 5 hrs	CHME Cell Line; treated				pCMVSPORT 3.0
S0420	CHME Cell Line,untreated	CHME Cell line, untreated				pSport1
S0422	Mo7e Cell Line GM-CSF treated (1ng/ml)	Mo7e Cell Line GM-CSF treated (1ng/ml)				pCMVSPORT 3.0
S0424	TF-1 Cell Line GM-CSF Treated	TF-1 Cell Line GM-CSF Treated				pSport1
S0426	Monocyte activated; re-excision	Monocyte-activated	blood	Cell Line		Uni-ZAP XR
S0428	Neutrophils control; re-excision	human neutrophils	blood	Cell Line		Uni-ZAP XR

S0430	Aryepiglottis Normal	Aryepiglottis Normal				pSport1
S0432	Sinus piniformis Tumour	Sinus piniformis Tumour				pSport1
S0434	Stomach Normal	Stomach Normal			disease	pSport1
S0436	Stomach Tumour	Stomach Tumour			disease	pSport1
S0438	Liver Normal Met5No	Liver Normal Met5No				pSport1
S0440	Liver Tumour Met 5 Tu	Liver Tumour				pSport1
S0442	Colon Normal	Colon Normal				pSport1
S0444	Colon Tumor	Colon Tumour			disease	pSport1
S0446	Tongue Tumour	Tongue Tumour				pSport1
S0448	Larynx Normal	Larynx Normal				pSport1
S0450	Larynx Tumour	Larynx Tumour				pSport1
S0452	Thymus	Thymus				pSport1
S0454	Placenta	Placenta	Placenta			pSport1
S0456	Tongue Normal	Tongue Normal				pSport1
S0458	Thyroid Normal (SDCA2 No)	Thyroid normal				pSport1
S0460	Thyroid Tumour	Thyroid Tumour				pSport1
S0462	Thyroid Thyroiditis	Thyroid Thyroiditis				pSport1
S0464	Larynx Normal	Larynx Normal				pSport1
S0466	Larynx Tumor	Larynx Tumor			disease	pSport1
S0468	Ea.hy.926 cell line	Ea.hy.926 cell line				pSport1
S0470	Adenocarcinoma	PYFD			disease	pSport1
S0472	Lung Mesothelium	PYBT				pSport1
S0474	Human blood platelets	Platelets	Blood platelets			Other
S0665	Human Amygdala; re-excision	Amygdala				Uni-ZAP XR
S3012	Smooth Muscle Serum Treated, Norm	Smooth muscle	Pulmonary artery	Cell Line		pBluescript
S3014	Smooth muscle, serum induced, re-exc	Smooth muscle	Pulmonary artery	Cell Line		pBluescript
S6014	H. hypothalamus, frac A	Hypothalamus	Brain			ZAP Express
S6016	H. Frontal Cortex, Epileptic	H. Brain, Frontal Cortex, Epileptic	Brain		disease	Uni-ZAP XR
S6022	H. Adipose Tissue	Human Adipose Tissue				Uni-ZAP XR
S6024	Alzheimers, spongy change	Alzheimer's/Spongy change	Brain		disease	Uni-ZAP XR
S6026	Frontal Lobe, Dementia	Frontal Lobe dementia/Alzheimer's	Brain			Uni-ZAP XR
S6028	Human Manic Depression Tissue	Human Manic depression tissue	Brain		disease	Uni-ZAP XR
T0001	Human Brown Fat	Brown Fat				pBluescript SK-
T0002	Activated T-cells	Activated T-Cell, PBL fraction	Blood	Cell Line		pBluescript SK-
T0003	Human Fetal Lung	Human Fetal Lung				pBluescript SK-
T0004	Human White Fat	Human White Fat				pBluescript SK-
T0006	Human Pineal Gland	Human Pineal Gland				pBluescript SK-
T0007	Colon Epithelium	Colon Epithelium				pBluescriptISK
T0008	Colorectal Tumor	Colorectal Tumor			disease	pBluescript SK-
T0010	Human Infant Brain	Human Infant Brain				Other
T0023	Human Pancreatic Carcinoma	Human Pancreatic Carcinoma			disease	pBluescript SK-
T0039	HSA 172 Cells	Human HSA172 cell line				pBluescript SK-
T0040	HSC172 cells	SA172 Cells				pBluescript SK-
T0041	Jurkat T-cell G1 phase	Jurkat T-cell				pBluescript SK-
T0042	Jurkat T-Cell, S phase	Jurkat T-Cell Line				pBluescript SK-

T0047	T lymphocytes >70	T lymphocytes > 70				pBluescript SK-
T0048	Human Aortic Endothelium	Human Aortic Endothilium				pBluescript SK-
T0049	Aorta endothelial cells + TNF-a	Aorta endothelial cells				pBluescript SK-
T0060	Human White Adipose	Human White Fat				pBluescript SK-
T0067	Human Thyroid	Human Thyroid				pBluescript SK-
T0068	Normal Ovary, Premenopausal	Normal Ovary, Premenopausal				pBluescript SK-
T0069	Human Uterus, normal	Human Uterus, normal				pBluescript SK-
T0071	Human Bone Marrow	Human Bone Marrow				pBluescript SK-
T0078	Human Liver, normal adult	Human Liver, normal Adult				pBluescript SK-
T0079	Human Kidney, normal Adult	Human Kidney, normal Adult				pBluescript SK-
T0082	Human Adult Retina	Human Adult Retina				pBluescript SK-
T0090	Liver, normal					pBluescript SK-
T0091	Liver, hepatocellular carcinoma					pBluescript SK-
T0104	HCC cell line metastasis to liver					pBluescript SK-
T0109	Human (HCC) cell line liver (mouse) metastasis, remake					pBluescript SK-
T0110	Human colon carcinoma (HCC) cell line, remake					pBluescript SK-
T0112	Human (Caco-2) cell line, adenocarcinoma, colon					pBluescript SK-
T0114	Human (Caco-2) cell line, adenocarcinoma, colon, remake					pBluescript SK-
T0115	Human Colon Carcinoma (HCC) cell line					pBluescript SK-
L0002	Atrium cDNA library Human heart					
L0004	ClonTech HL 1065a					
L0005	Clontech human aorta polyA+ mRNA (#6572)					
L0012	HDMEC cDNA library					
L0015	Human					
L0021	Human adult (K.Okubo)					
L0022	Human adult lung 3" directed MboI cDNA					
L0024	Human brain ARSanders					
L0032	Human chromosome 12p cDNAs					
L0034	Human chromosome 14					
L0038	Human chromosome 6					
L0040	Human colon mucosa					
L0041	Human epidermal keratinocyte					
L0051	Human mRNA (Tripodis and Ragoussis)					
L0052	Human normalized K562-cDNA					
L0055	Human promyelocyte					
L0060	Human thymus NSTH II					
L0065	Liver HepG2 cell line.					
L0070	Selected chromosome 21 cDNA library					
L0096	Subtracted human retina					
L0097	Subtracted human retinal					

	pigment epithelium (RPE)					
L0103	DKFZphamy1	amygdala				
L0105	Human aorta polyA+ (TFujiwara)	aorta				
L0109	Human brain cDNA	brain				
L0118	Human fetal brain S. Meier-Ewert	brain				
L0126	Human fibroblast cDNA	fibroblast				
L0138	Human normal gingiva	normal gingiva				
L0141	Human pancreatic islet cell	pancreatic islet				
L0142	Human placenta cDNA (TFujiwara)	placenta				
L0143	Human placenta polyA+ (TFujiwara)	placenta				
L0146	Human fovea cDNA	retinal fovea				
L0149	DKFZphsnu1	subthalamic nucleus				
L0157	Human fetal brain (TFujiwara)		brain			
L0162	Human brain frontal cortex	frontal cortex	brain			
L0163	Human heart cDNA (YNakamura)		heart			
L0171	Human lung adenocarcinoma A549	lung adenocarcinoma		A549		
L0175	Human retina cell line ARPE-19	retina		ARPE-19		
L0177	Human newborn melanocytes (T.Vogt)			Clonetics Corp. (San Diego, CA) strain #68 and 2486		
L0182	Human HeLa (Y.Wang)			HeLa		
L0186	Human salivary gland cell line HSG	salivary gland		HSG		
L0194	Human pancreatic cancer cell line Patu 8988t	pancreatic cancer		Patu 8988t		
L0351	Infant brain, Bento Soares					BA, M13-derived
L0352	Normalized infant brain, Bento Soares					BA, M13-derived
L0353	21q Placenta, F. Tassone and K. Gardiner					Bluescript
L0355	P, Human foetal Brain Whole tissue					Bluescript
L0356	S, Human foetal Adrenals tissue					Bluescript
L0361	Stratagene ovary (#937217)		ovary			Bluescript SK
L0362	Stratagene ovarian cancer (#937219)					Bluescript SK-
L0363	NCI_CGAP_GC2	germ cell tumor				Bluescript SK-
L0364	NCI_CGAP_GC5	germ cell tumor				Bluescript SK-
L0365	NCI_CGAP_Phe1	pheochromocytoma				Bluescript SK-
L0366	Stratagene schizo brain S11	schizophrenic brain S-11 frontal lobe				Bluescript SK-
L0367	NCI_CGAP_Sch1	Schwannoma tumor				Bluescript SK-
L0368	NCI_CGAP_SS1	synovial sarcoma				Bluescript SK-
L0369	NCI_CGAP_AA1	adrenal adenoma	adrenal gland			Bluescript SK-
L0370	Johnston frontal cortex	pooled frontal lobe	brain			Bluescript SK-
L0371	NCI_CGAP_Br3	breast tumor	breast			Bluescript SK-
L0372	NCI_CGAP_Co12	colon tumor	colon			Bluescript SK-

L0373	NCI_CGAP_Co11	tumor	colon			Bluescript SK-
L0374	NCI_CGAP_Co2	tumor	colon			Bluescript SK-
L0375	NCI_CGAP_Kid6	kidney tumor	kidney			Bluescript SK-
L0376	NCI_CGAP_Lar1	larynx	larynx			Bluescript SK-
L0377	NCI_CGAP_HN2	squamous cell carcinoma from vocal cord	larynx			Bluescript SK-
L0378	NCI_CGAP_Lu1	lung tumor	lung			Bluescript SK-
L0379	NCI_CGAP_Lym3	lymphoma	lymph node			Bluescript SK-
L0380	NCI_CGAP_HN1	squamous cell carcinoma	lymph node			Bluescript SK-
L0381	NCI_CGAP_HN4	squamous cell carcinoma	pharynx			Bluescript SK-
L0382	NCI_CGAP_Pr25	epithelium (cell line)	prostate			Bluescript SK-
L0383	NCI_CGAP_Pr24	invasive tumor (cell line)	prostate			Bluescript SK-
L0384	NCI_CGAP_Pr23	prostate tumor	prostate			Bluescript SK-
L0385	NCI_CGAP_Gas1	gastric tumor	stomach			Bluescript SK-
L0386	NCI_CGAP_HN3	squamous cell carcinoma from base of tongue	tongue			Bluescript SK-
L0387	NCI_CGAP_GCB0	germinal center B-cells	tonsil			Bluescript SK-
L0388	NCI_CGAP_HN6	normal gingiva (cell line from immortalized kerati				Bluescript SK-
L0389	NCI_CGAP_HN5	normal gingiva (cell line from primary keratinocyt				Bluescript SK-
L0393	B, Human Liver tissue					gt11
L0394	H, Human adult Brain Cortex tissue					gt11
L0404	b4HB3MA Cot109+103+85-Bio					Lafmid A
L0405	b4HB3MA Cot109+103-Bio					Lafmid A
L0411	1-NIB					Lafmid BA
L0414	b4HB3MA					Lafmid BA
L0415	b4HB3MA Cot8-HAP-Ft					Lafmid BA
L0416	b4HB3MA-Cot0.38-HAP-B					Lafmid BA
L0417	b4HB3MA-Cot0.38-HAP-Ft-6					Lafmid BA
L0424	b4HB3MA-Cot14.5					Lafmid BA
L0425	b4HB3MA-Cot18-Bio					Lafmid BA
L0426	b4HB3MA-Cot51.5-HAP-Ft					Lafmid BA
L0427	b4HB3MA-FT20%-Biotin					Lafmid BA
L0428	Cot1374Ft-4HB3MA					Lafmid BA
L0430	Cot250Ft-b4HB3MA					Lafmid BA
L0434	Infant brain library of Dr. M. Soares					lafmid BA
L0435	Infant brain, LLNL array of Dr. M. Soares 1NIB					lafmid BA
L0437	N-b4HB3MA-Cot109					Lafmid BA
L0438	normalized infant brain cDNA	total brain	brain			lafmid BA
L0439	Soares infant brain 1NIB		whole brain			Lafmid BA
L0441	2HB3MK					Lafmid BK
L0442	4HB3MK					Lafmid BK
L0443	b4HB3MK					Lafmid BK
L0446	N4HB3MK					Lafmid BK
L0448	3HFLSK20					Lafmid K
L0451	N3HFLSK20					Lafmid K

L0453	BATM1					lambda gt10
L0454	Clontech adult human fat cell library HL1108A					lambda gt10
L0455	Human retina cDNA randomly primed sublibrary	retina	eye			lambda gt10
L0456	Human retina cDNA Tsp509I-cleaved sublibrary	retina	eye			lambda gt10
L0459	Adult heart, Clontech					Lambda gt11
L0460	Adult heart, Lambda gt11					Lambda gt11
L0462	WATM1					lambda gt11
L0463	fetal brain cDNA	brain	brain			lambda gt11
L0465	TEST1, Human adult Testis tissue					lambda nm1149
L0468	HE6W					lambda zap
L0469	T, Human adult Rhabdomyosarcoma cell-line					Lambda Zap
L0470	BL29 Burkitt's lymphoma, Pascalis Sideras					lambda ZAP 2
L0471	Human fetal heart, Lambda ZAP Express					Lambda ZAP Express
L0475	KG1-a Lambda Zap Express cDNA library			KG1-a		Lambda Zap Express (Stratagene)
L0476	Fetal brain, Stratagene					Lambda ZAP II
L0477	HPLA CCLee	placenta				Lambda ZAP II
L0480	Stratagene cat#937212 (1992)					Lambda ZAP, pBluescript SK(-)
L0481	CD34+DIRECTIONAL					Lambda ZAPII
L0482	HT29M6					Lambda ZAPII
L0483	Human pancreatic islet					Lambda ZAPII
L0485	STRATAGENE Human skeletal muscle cDNA library, cat. #936215.	skeletal muscle	leg muscle			Lambda ZAPII
L0487	Human peripheral blood (Steve Elledge)	whole peripheral blood				Lambda-Yes
L0492	Human Genomic					pAMP
L0493	NCI_CGAP_Ov26	papillary serous carcinoma	ovary			pAMP1
L0497	NCI_CGAP_HSC4	CD34+, CD38- from normal bone marrow donor	bone marrow			pAMP1
L0498	NCI_CGAP_HSC3	CD34+, T negative, patient with chronic myelogenous	bone marrow			pAMP1
L0499	NCI_CGAP_HSC2	stem cell 34+/38+	bone marrow			pAMP1
L0500	NCI_CGAP_Brn20	oligodendroglioma	brain			pAMP1
L0501	NCI_CGAP_Brn21	oligodendroglioma	brain			pAMP1
L0502	NCI_CGAP_Br15	adenocarcinoma	breast			pAMP1
L0503	NCI_CGAP_Br17	adenocarcinoma	breast			pAMP1
L0504	NCI_CGAP_Br13	breast carcinoma in situ	breast			pAMP1
L0505	NCI_CGAP_Br12	invasive carcinoma	breast			pAMP1
L0506	NCI_CGAP_Br16	lobular carcinoma in situ	breast			pAMP1
L0507	NCI_CGAP_Br14	normal epithelium	breast			pAMP1
L0508	NCI_CGAP_Lu25	bronchioalveolar carcinoma	lung			pAMP1
L0509	NCI_CGAP_Lu26	invasive adenocarcinoma	lung			pAMP1

L0510	NCI_CGAP_Ov33	borderline ovarian carcinoma	ovary			pAMP1
L0511	NCI_CGAP_Ov34	borderline ovarian carcinoma	ovary			pAMP1
L0512	NCI_CGAP_Ov36	borderline ovarian carcinoma	ovary			pAMP1
L0513	NCI_CGAP_Ov37	early stage papillary serous carcinoma	ovary			pAMP1
L0514	NCI_CGAP_Ov31	papillary serous carcinoma	ovary			pAMP1
L0515	NCI_CGAP_Ov32	papillary serous carcinoma	ovary			pAMP1
L0517	NCI_CGAP_Pr1					pAMP10
L0518	NCI_CGAP_Pr2					pAMP10
L0519	NCI_CGAP_Pr3					pAMP10
L0520	NCI_CGAP_Alv1	alveolar rhabdomyosarcoma				pAMP10
L0521	NCI_CGAP_Ew1	Ewing's sarcoma				pAMP10
L0522	NCI_CGAP_Kid1	kidney				pAMP10
L0523	NCI_CGAP_Lip2	liposarcoma				pAMP10
L0524	NCI_CGAP_Li1	liver				pAMP10
L0525	NCI_CGAP_Li2	liver				pAMP10
L0526	NCI_CGAP_Pr12	metastatic prostate bone lesion				pAMP10
L0527	NCI_CGAP_Ov2	ovary				pAMP10
L0528	NCI_CGAP_Pr5	prostate				pAMP10
L0529	NCI_CGAP_Pr6	prostate				pAMP10
L0530	NCI_CGAP_Pr8	prostate				pAMP10
L0531	NCI_CGAP_Pr20	prostate metastasis, liver				pAMP10
L0532	NCI_CGAP_Thy1	thyroid				pAMP10
L0533	NCI_CGAP_HSC1	stem cells	bone marrow			pAMP10
L0534	Chromosome 7 Fetal Brain cDNA Library	brain	brain			pAMP10
L0535	NCI_CGAP_Br5	infiltrating ductal carcinoma	breast			pAMP10
L0536	NCI_CGAP_Br4	normal ductal tissue	breast			pAMP10
L0537	NCI_CGAP_Ov6	normal cortical stroma	ovary			pAMP10
L0539	Chromosome 7 Placental cDNA Library		placenta			pAMP10
L0540	NCI_CGAP_Pr10	invasive prostate tumor	prostate			pAMP10
L0541	NCI_CGAP_Pr7	low-grade prostatic neoplasia	prostate			pAMP10
L0542	NCI_CGAP_Pr11	normal prostatic epithelial cells	prostate			pAMP10
L0543	NCI_CGAP_Pr9	normal prostatic epithelial cells	prostate			pAMP10
L0544	NCI_CGAP_Pr4	prostatic intraepithelial neoplasia - high grade	prostate			pAMP10
L0545	NCI_CGAP_Pr4.1	prostatic intraepithelial neoplasia - high grade	prostate			pAMP10
L0546	NCI_CGAP_Pr18	stroma	prostate			pAMP10
L0547	NCI_CGAP_Pr16	tumor	prostate			pAMP10
L0548	Chromosome 7 Thymus cDNA Library	thymus	thymus			pAMP10
L0549	NCI_CGAP_HN10	carcinoma in situ from retromolar trigone				pAMP10

L0550	NCI_CGAP_HN9	normal squamous epithelium from retromolar trigone				pAMP10
L0551	NCI_CGAP_HN7	normal squamous epithelium, floor of mouth				pAMP10
L0552	NCI_CGAP_HN8	well-differentiated invasive carcinoma, floor of m				pAMP10
L0553	NCI_CGAP_Co22	colonic adenocarcinoma	colon			pAMP10
L0554	NCI_CGAP_Li8		liver			pAMP10
L0555	NCI_CGAP_Lu34	large cell carcinoma	lung			pAMP10
L0556	NCI_CGAP_Lu34.1	large cell carcinoma	lung			pAMP10
L0558	NCI_CGAP_Ov40	endometrioid ovarian metastasis	ovary			pAMP10
L0559	NCI_CGAP_Ov39	papillary serous ovarian metastasis	ovary			pAMP10
L0560	NCI_CGAP_HN12	moderate to poorly differentiated invasive carcino	tongue			pAMP10
L0561	NCI_CGAP_HN11	normal squamous epithelium	tongue			pAMP10
L0562	Chromosome 7 HeLa cDNA Library			HeLa cell line; ATCC		pAMP10
L0563	Human Bone Marrow Stromal Fibroblast	bone marrow				pBluescript
L0564	Jia bone marrow stroma	bone marrow stroma				pBluescript
L0565	Normal Human Trabecular Bone Cells	Bone	Hip			pBluescript
L0579	Human fetal brain QBoqin2	cerebrum and cerebellum				pBluescript SK
L0581	Stratagene liver (#937224)		liver			pBluescript SK
L0583	Stratagene cDNA library Human fibroblast, cat#937212					pBluescript SK(+)
L0584	Stratagene cDNA library Human heart, cat#936208					pBluescript SK(+)
L0586	HTCDL1					pBluescript SK(-)
L0587	Stratagene colon HT29 (#937221)					pBluescript SK-
L0588	Stratagene endothelial cell 937223					pBluescript SK-
L0589	Stratagene fetal retina 937202					pBluescript SK-
L0590	Stratagene fibroblast (#937212)					pBluescript SK-
L0591	Stratagene HeLa cell s3 937216					pBluescript SK-
L0592	Stratagene hNT neuron (#937233)					pBluescript SK-
L0593	Stratagene neuroepithelium (#937231)					pBluescript SK-
L0594	Stratagene neuroepithelium NT2RAMI 937234					pBluescript SK-
L0595	Stratagene NT2 neuronal precursor 937230	neuroepithelial cells	brain			pBluescript SK-
L0596	Stratagene colon (#937204)		colon			pBluescript SK-
L0597	Stratagene corneal stroma (#937222)		cornea			pBluescript SK-

L0598	Morton Fetal Cochlea	cochlea	ear			pBluescript SK-
L0599	Stratagene lung (#937210)		lung			pBluescript SK-
L0600	Weizmann Olfactory Epithelium	olfactory epithelium	nose			pBluescript SK-
L0601	Stratagene pancreas (#937208)		pancreas			pBluescript SK-
L0602	Pancreatic Islet	pancreatic islet	pancreas			pBluescript SK-
L0603	Stratagene placenta (#937225)		placenta			pBluescript SK-
L0604	Stratagene muscle 937209	muscle	skeletal muscle			pBluescript SK-
L0605	Stratagene fetal spleen (#937205)	fetal spleen	spleen			pBluescript SK-
L0606	NCI_CGAP_Lym5	follicular lymphoma	lymph node			pBluescript SK-
L0607	NCI_CGAP_Lym6	mantle cell lymphoma	lymph node			pBluescript SK-
L0608	Stratagene lung carcinoma 937218	lung carcinoma	lung	NCI-H69		pBluescript SK-
L0609	Schiller astrocytoma	astrocytoma	brain			pBluescript SK-(Stratagene)
L0611	Schiller meningioma	meningioma	brain			pBluescript SK-(Stratagene)
L0612	Schiller oligodendroglioma	oligodendroglioma	brain			pBluescript SK-(Stratagene)
L0615	22 week old human fetal liver cDNA library					pBluescriptII SK(-)
L0617	Chromosome 22 exon					pBluescriptIIK S+
L0619	Chromosome 9 exon II					pBluescriptIIK S+
L0622	HM1					pcDNAII (Invitrogen)
L0623	HM3	pectoral muscle (after mastectomy)				pcDNAII (Invitrogen)
L0625	NCI_CGAP_AR1	bulk alveolar tumor				pCMV-SPORT2
L0626	NCI_CGAP_GC1	bulk germ cell seminoma				pCMV-SPORT2
L0627	NCI_CGAP_Co1	bulk tumor	colon			pCMV-SPORT2
L0628	NCI_CGAP_Ov1	ovary bulk tumor	ovary			pCMV-SPORT2
L0629	NCI_CGAP_Mel3	metastatic melanoma to bowel	bowel (skin primary)			pCMV-SPORT4
L0630	NCI_CGAP_CNS1	substantia nigra	brain			pCMV-SPORT4
L0631	NCI_CGAP_Br7		breast			pCMV-SPORT4
L0632	NCI_CGAP_Li5	hepatic adenoma	liver			pCMV-SPORT4
L0633	NCI_CGAP_Lu6	small cell carcinoma	lung			pCMV-SPORT4
L0634	NCI_CGAP_Ov8	serous adenocarcinoma	ovary			pCMV-SPORT4
L0635	NCI_CGAP_PNS1	dorsal root ganglion	peripheral nervous system			pCMV-SPORT4
L0636	NCI_CGAP_Pit1	four pooled pituitary adenomas	brain			pCMV-SPORT6
L0637	NCI_CGAP_Brn53	three pooled meningiomas	brain			pCMV-SPORT6
L0638	NCI_CGAP_Brn35	tumor, 5 pooled (see description)	brain			pCMV-SPORT6
L0639	NCI_CGAP_Brn52	tumor, 5 pooled (see description)	brain			pCMV-SPORT6

L0640	NCI_CGAP_Br18	four pooled high-grade tumors, including two prima	breast			pCMV-SPORT6
L0641	NCI_CGAP_Co17	juvenile granulosa tumor	colon			pCMV-SPORT6
L0642	NCI_CGAP_Co18	moderately differentiated adenocarcinoma	colon			pCMV-SPORT6
L0643	NCI_CGAP_Co19	moderately differentiated adenocarcinoma	colon			pCMV-SPORT6
L0644	NCI_CGAP_Co20	moderately differentiated adenocarcinoma	colon			pCMV-SPORT6
L0645	NCI_CGAP_Co21	moderately differentiated adenocarcinoma	colon			pCMV-SPORT6
L0646	NCI_CGAP_Co14	moderately-differentiated adenocarcinoma	colon			pCMV-SPORT6
L0647	NCI_CGAP_Sar4	five pooled sarcomas, including myxoid liposarcoma	connective tissue			pCMV-SPORT6
L0648	NCI_CGAP_Eso2	squamous cell carcinoma	esophagus			pCMV-SPORT6
L0649	NCI_CGAP_GU1	2 pooled high-grade transitional cell tumors	genitourinary tract			pCMV-SPORT6
L0650	NCI_CGAP_Kid13	2 pooled Wilms' tumors, one primary and one metast	kidney			pCMV-SPORT6
L0651	NCI_CGAP_Kid8	renal cell tumor	kidney			pCMV-SPORT6
L0652	NCI_CGAP_Lu27	four pooled poorly-differentiated adenocarcinomas	lung			pCMV-SPORT6
L0653	NCI_CGAP_Lu28	two pooled squamous cell carcinomas	lung			pCMV-SPORT6
L0654	NCI_CGAP_Lu31		lung, cell line			pCMV-SPORT6
L0655	NCI_CGAP_Lym12	lymphoma, follicular mixed small and large cell	lymph node			pCMV-SPORT6
L0656	NCI_CGAP_Ov38	normal epithelium	ovary			pCMV-SPORT6
L0657	NCI_CGAP_Ov23	tumor, 5 pooled (see description)	ovary			pCMV-SPORT6
L0658	NCI_CGAP_Ov35	tumor, 5 pooled (see description)	ovary			pCMV-SPORT6
L0659	NCI_CGAP_Pan1	adenocarcinoma	pancreas			pCMV-SPORT6
L0661	NCI_CGAP_Mel15	malignant melanoma, metastatic to lymph node	skin			pCMV-SPORT6
L0662	NCI_CGAP_Gas4	poorly differentiated adenocarcinoma with signet r	stomach			pCMV-SPORT6
L0663	NCI_CGAP_Ut2	moderately-differentiated endometrial adenocarcino	uterus			pCMV-SPORT6
L0664	NCI_CGAP_Ut3	poorly-differentiated endometrial adenocarcinoma,	uterus			pCMV-SPORT6

L0665	NCI_CGAP_Ut4	serous papillary carcinoma, high grade, 2 pooled t	uterus			pCMV-SPORT6
L0666	NCI_CGAP_Ut1	well-differentiated endometrial adenocarcinoma, 7	uterus			pCMV-SPORT6
L0667	NCI_CGAP_CML1	myeloid cells, 18 pooled CML cases, BCR/ABL rearra	whole blood			pCMV-SPORT6
L0669	Human MCF7 cDNA subtracted with MDA-MB-231 cDNA	breast adenocarcinoma	breast	MCF7		pCR II [Invitrogen]
L0681	Stanley Frontal SN individual	frontal lobe (see description)	brain			pCR2.1 (Invitrogen)
L0682	Stanley Frontal NB pool 2	frontal lobe (see description)	brain			pCR2.1-TOPO (Invitrogen)
L0683	Stanley Frontal NS pool 2	frontal lobe (see description)	brain			pCR2.1-TOPO (Invitrogen)
L0684	Stanley Frontal SB pool 1	frontal lobe (see description)	brain			pCR2.1-TOPO (Invitrogen)
L0686	Stanley Frontal SN pool 2	frontal lobe (see description)	brain			pCR2.1-TOPO (Invitrogen)
L0687	Stanley Hippocampus NB pool 1	hippocampus (see description)	brain			pCR2.1-TOPO (Invitrogen)
L0688	Stanley Hippocampus SB pool 1	hippocampus (see description)	brain			pCR2.1-TOPO (Invitrogen)
L0689	Stanley Hippocampus SN pool 1	hippocampus (see description)	brain			pCR2.1-TOPO (Invitrogen)
L0695	Human Glioblastoma Cell		Brain	BT-325		PCRII, Invitrogen
L0697	Testis 1					PGEM 5zf(+)
L0698	Testis 2					PGEM 5zf(+)
L0716	PMA-induced HL60 cell subtraction library			PMA-induced HL60 human leukemic cell line		pSPORT 1
L0717	Gessler Wilms tumor					pSPORT1
L0718	Testis 5					pSPORT1
L0719	human embryo cDNA library	Whole embryo				pSPORT1
L0731	Soares_pregnant_uterus_NbHPU		uterus			pT7T3-Pac
L0738	Human colorectal cancer					pT7T3D
L0740	Soares melanocyte 2NbHM	melanocyte				pT7T3D (Pharmacia) with a modified polylinker
L0741	Soares adult brain N2b4HB55Y		brain			pT7T3D (Pharmacia) with a modified polylinker
L0742	Soares adult brain N2b5HB55Y		brain			pT7T3D (Pharmacia) with a modified polylinker
L0743	Soares breast 2NbHBst		breast			pT7T3D (Pharmacia) with a modified polylinker
L0744	Soares breast 3NbHBst		breast			pT7T3D (Pharmacia) with a modified polylinker

L0745	Soares retina N2b4HR	retina	eye			pT7T3D (Pharmacia) with a modified polylinker
L0746	Soares retina N2b5HR	retina	eye			pT7T3D (Pharmacia) with a modified polylinker
L0747	Soares_fetal_heart_NbHH 19W		heart			pT7T3D (Pharmacia) with a modified polylinker
L0748	Soares fetal liver spleen 1NFLS		Liver and Spleen			pT7T3D (Pharmacia) with a modified polylinker
L0749	Soares_fetal_liver_spleen _1NFLS_S1		Liver and Spleen			pT7T3D (Pharmacia) with a modified polylinker
L0750	Soares_fetal_lung_NbHL1 9W		lung			pT7T3D (Pharmacia) with a modified polylinker
L0751	Soares ovary tumor NbHOT	ovarian tumor	ovary			pT7T3D (Pharmacia) with a modified polylinker
L0752	Soares_parathyroid_tumor _NbHPA	parathyroid tumor	parathyroid gland			pT7T3D (Pharmacia) with a modified polylinker
L0753	Soares_pineal_gland_N3H PG		pineal gland			pT7T3D (Pharmacia) with a modified polylinker
L0754	Soares placenta Nb2HP		placenta			pT7T3D (Pharmacia) with a modified polylinker
L0755	Soares_placenta_8to9wee ks_2NbHP8to9W		placenta			pT7T3D (Pharmacia) with a modified polylinker
L0756	Soares_multiple_sclerosis _2NbHMSP	multiple sclerosis lesions				pT7T3D (Pharmacia) with a modified polylinker V_TYPE
L0757	Soares_senescent_fibrobla sts_NbHSF	senescent fibroblast				pT7T3D (Pharmacia) with a modified polylinker V_TYPE
L0758	Soares_testis_NHT					pT7T3D-Pac (Pharmacia) with a modified polylinker
L0759	Soares_total_fetus_Nb2H F8_9w					pT7T3D-Pac (Pharmacia) with a modified polylinker
L0760	Barstead aorta HPLRB3	aorta				pT7T3D-Pac (Pharmacia) with a modified polylinker

L0761	NCI_CGAP_CLL1	B-cell, chronic lymphocytic leukemia				pT7T3D-Pac (Pharmacia) with a modified polylinker
L0762	NCI_CGAP_Br1.1	breast				pT7T3D-Pac (Pharmacia) with a modified polylinker
L0763	NCI_CGAP_Br2	breast				pT7T3D-Pac (Pharmacia) with a modified polylinker
L0764	NCI_CGAP_Co3	colon				pT7T3D-Pac (Pharmacia) with a modified polylinker
L0765	NCI_CGAP_Co4	colon				pT7T3D-Pac (Pharmacia) with a modified polylinker
L0766	NCI_CGAP_GCB1	germinal center B cell				pT7T3D-Pac (Pharmacia) with a modified polylinker
L0767	NCI_CGAP_GC3	pooled germ cell tumors				pT7T3D-Pac (Pharmacia) with a modified polylinker
L0768	NCI_CGAP_GC4	pooled germ cell tumors				pT7T3D-Pac (Pharmacia) with a modified polylinker
L0769	NCI_CGAP_Brn25	anaplastic oligodendroglioma	brain			pT7T3D-Pac (Pharmacia) with a modified polylinker
L0770	NCI_CGAP_Brn23	glioblastoma (pooled)	brain			pT7T3D-Pac (Pharmacia) with a modified polylinker
L0771	NCI_CGAP_Co8	adenocarcinoma	colon			pT7T3D-Pac (Pharmacia) with a modified polylinker
L0772	NCI_CGAP_Co10	colon tumor RER+	colon			pT7T3D-Pac (Pharmacia) with a modified polylinker
L0773	NCI_CGAP_Co9	colon tumor RER+	colon			pT7T3D-Pac (Pharmacia) with a modified polylinker
L0774	NCI_CGAP_Kid3		kidney			pT7T3D-Pac (Pharmacia) with a modified polylinker
L0775	NCI_CGAP_Kid5	2 pooled tumors (clear cell type)	kidney			pT7T3D-Pac (Pharmacia) with a modified polylinker
L0776	NCI_CGAP_Lu5	carcinoid	lung			pT7T3D-Pac (Pharmacia) with a modified polylinker
L0777	Soares_NhHMPu_S1	Pooled human melanocyte, fetal	mixed (see below)			pT7T3D-Pac (Pharmacia)

		heart, and pregnant				with a modified polylinker
L0778	Barstead pancreas HPLRB1		pancreas			pT7T3D-Pac (Pharmacia) with a modified polylinker
L0779	Soares_NFL_T_GBC_S1		pooled			pT7T3D-Pac (Pharmacia) with a modified polylinker
L0780	Soares_NSF_F8_9W_OT_PA_P_S1		pooled			pT7T3D-Pac (Pharmacia) with a modified polylinker
L0782	NCI_CGAP_Pr21	normal prostate	prostate			pT7T3D-Pac (Pharmacia) with a modified polylinker
L0783	NCI_CGAP_Pr22	normal prostate	prostate			pT7T3D-Pac (Pharmacia) with a modified polylinker
L0784	NCI_CGAP_Lei2	leiomyosarcoma	soft tissue			pT7T3D-Pac (Pharmacia) with a modified polylinker
L0785	Barstead spleen HPLRB2		spleen			pT7T3D-Pac (Pharmacia) with a modified polylinker
L0786	Soares_NbHFB		whole brain			pT7T3D-Pac (Pharmacia) with a modified polylinker
L0787	NCI_CGAP_Sub1					pT7T3D-Pac (Pharmacia) with a modified polylinker
L0788	NCI_CGAP_Sub2					pT7T3D-Pac (Pharmacia) with a modified polylinker
L0789	NCI_CGAP_Sub3					pT7T3D-Pac (Pharmacia) with a modified polylinker
L0790	NCI_CGAP_Sub4					pT7T3D-Pac (Pharmacia) with a modified polylinker
L0791	NCI_CGAP_Sub5					pT7T3D-Pac (Pharmacia) with a modified polylinker
L0792	NCI_CGAP_Sub6					pT7T3D-Pac (Pharmacia) with a modified polylinker
L0793	NCI_CGAP_Sub7					pT7T3D-Pac (Pharmacia) with a modified polylinker
L0794	NCI_CGAP_GC6	pooled germ cell tumors				pT7T3D-Pac (Pharmacia) with a modified polylinker

L0796	NCI_CGAP_Brn50	medulloblastoma	brain			pT7T3D-Pac (Pharmacia) with a modified polylinker
L0800	NCI_CGAP_Co16	colon tumor, RER+	colon			pT7T3D-Pac (Pharmacia) with a modified polylinker
L0803	NCI_CGAP_Kid11		kidney			pT7T3D-Pac (Pharmacia) with a modified polylinker
L0804	NCI_CGAP_Kid12	2 pooled tumors (clear cell type)	kidney			pT7T3D-Pac (Pharmacia) with a modified polylinker
L0805	NCI_CGAP_Lu24	carcinoid	lung			pT7T3D-Pac (Pharmacia) with a modified polylinker
L0806	NCI_CGAP_Lu19	squamous cell carcinoma, poorly differentiated (4	lung			pT7T3D-Pac (Pharmacia) with a modified polylinker
L0807	NCI_CGAP_Ov18	fibrotheoma	ovary			pT7T3D-Pac (Pharmacia) with a modified polylinker
L0808	Barstead prostate BPH HPLRB4 1		prostate			pT7T3D-Pac (Pharmacia) with a modified polylinker
L0809	NCI_CGAP_Pr28		prostate			pT7T3D-Pac (Pharmacia) with a modified polylinker
L0811	BATM2					PTZ18
L2245	NEM subtracted human fetal kidney cDNA					pUEX1
L2250	Human cerebral cortex	cerebral cortex				
L2251	Human fetal lung	Fetal lung				
L2252	Human placenta	placenta				
L2269	NCI_CGAP_Thy11	follicular carcinoma	thyroid			pAMP10
L3872	NCI_CGAP_Skn1		skin, normal, 4 pooled sa			pCMV-SPORT6
L3904	NCI_CGAP_Brn64	glioblastoma with EGFR amplification	brain			pCMV-SPORT6
L3905	NCI_CGAP_Brn67	anaplastic oligodendroglioma with 1p/19q loss	brain			pCMV-SPORT6
L4497	NCI_CGAP_Br22	invasive ductal carcinoma, 3 pooled samples	breast			pCMV-SPORT6
L4501	NCI_CGAP_Sub8					pT7T3D-Pac (Pharmacia) with a modified polylinker
L4508	NCI_CGAP_Thy8	normal epithelium	thyroid			pAMP10
L4557	NCI_CGAP_Adr1	neuroblastoma	adrenal gland			pCMV-SPORT6
L4559	NCI_CGAP_Thy3	follicular carcinoma	thyroid			pCMV-SPORT6
L4669	NCI_CGAP_Ov41	serous papillary tumor	ovary			pCMV-SPORT6
L4747	NCI_CGAP_Brn41	oligodendroglioma	brain			pT7T3D-Pac

						(Pharmacia) with a modified polylinker
L4775	NCI_CGAP_Thy12	papillary carcinoma	thyroid			pAMP10
L5286	NCI_CGAP_Thy10	medullary carcinoma	thyroid			pAMP10
L5564	NCI_CGAP_HN20		normal head/neck tissue			pAMP1
L5565	NCI_CGAP_Brn66	glioblastoma with probably TP53 mutation and witho	brain			pCMV- SPORT6
L5566	NCI_CGAP_Brn70	anaplastic oligodendroglioma	brain			pCMV- SPORT6.ccdB
L5568	NCI_CGAP_HN21	nasopharyngeal carcinoma	head/neck			pAMP1
L5569	NCI_CGAP_HN17	normal epithelium	nasopharynx			pAMP10
L5574	NCI_CGAP_HN19	normal epithelium	nasopharynx			pAMP10
L5575	NCI_CGAP_Brn65	glioblastoma without EGFR amplification	brain			pCMV- SPORT6
L5622	NCI_CGAP_Skn3		skin			pCMV- SPORT6
L5623	NCI_CGAP_Skn4	squamous cell carcinoma	skin			pCMV- SPORT6

TABLE 5

OMIM Reference	Description
100678	ACAT2 deficiency
100690	Myasthenic syndrome, slow-channel congenital, 601462
100730	Myasthenia gravis, neonatal transient
101000	Meningioma, NF2-related, sporadic Schwannoma, sporadic
101000	Neurofibromatosis, type 2
101000	Neurolemmomatosis
101000	Malignant mesothelioma, sporadic
102200	Somatotrophinoma
102480	Male infertility due to acrosin deficiency
102540	Cardiomyopathy, idiopathic dilated
102578	Leukemia, acute promyelocytic, PML/RARA type
102700	Severe combined immunodeficiency due to ADA deficiency
102700	Hemolytic anemia due to ADA excess
102770	Myoadenylate deaminase deficiency
102772	[AMP deaminase deficiency, erythrocytic]
103000	Hemolytic anemia due to adenylate kinase deficiency
103050	Autism, succinylpurinemic
103050	Adenylosuccinase deficiency
103581	Albright hereditary osteodystrophy-2
103600	[Dysalbuminemic hyperthyroxinemia]

103600	[Dysalbuminemic hyperzincemia], 194470
103600	Analbuminemia
103720	Alcoholism, susceptibility to
103850	Aldolase A deficiency
103950	Emphysema due to alpha-2-macroglobulin deficiency
104150	[AFP deficiency, congenital]
104150	[Hereditary persistence of alpha-fetoprotein]
104170	NAGA deficiency, mild
104170	Schindler disease
104170	Kanzaki disease
104311	Alzheimer disease-3
104500	Amelogenesis imperfecta-2, hypoplastic local type
104770	Amyloidosis, secondary, susceptibility to
105580	Anal canal carcinoma
105600	Dyserythropoietic anemia, congenital, type III
106100	Angioedema, hereditary
106150	Hypertension, essential, susceptibility to
106150	Preeclampsia, susceptibility to
106165	Hypertension, essential, 145500
106210	Peters anomaly
106210	Cataract, congenital, with late-onset corneal dystrophy
106210	Foveal hypoplasia, isolated, 136520
106210	Aniridia
106300	Ankylosing spondylitis
107271	CD59 deficiency
107300	Antithrombin III deficiency
107470	Atypical mycobacterial infection, familial disseminated, 209950
107470	BCG infection, generalized familial
107470	Tuberculosis, susceptibility to
107670	Apolipoprotein A-II deficiency
107680	ApoA-I and apoC-III deficiency, combined
107680	Corneal clouding, autosomal recessive
107680	Amyloidosis, 3 or more types
107680	Hypertriglyceridemia, one form
107680	Hypoalphalipoproteinemia
107720	Hypertriglyceridemia
107741	Hyperlipoproteinemia, type III
107777	Diabetes insipidus, nephrogenic, autosomal recessive, 222000
107970	Arrhythmogenic right ventricular dysplasia-1
108120	Distal arthrogyposis-1
108725	Atherosclerosis, susceptibility to
108730	Brody myopathy, 601003
108800	Atrial septal defect, secundum type
108962	Hypertension, salt-resistant
109150	Machado-Joseph disease
109270	Renal tubular acidosis, distal, 179800

109270	Spherocytosis, hereditary
109270	[Acanthocytosis, one form]
109270	[Elliptocytosis, Malaysian-Melanesian type]
109270	Hemolytic anemia due to band 3 defect
109565	Lymphoma, B-cell
109565	Lymphoma, diffuse large cell
109690	Asthma, nocturnal, susceptibility to
109690	Obesity, susceptibility to
109700	Hemodialysis-related amyloidosis
110100	Blepharophimosis, epicanthus inversus, and ptosis, type 1
110700	Vivax malaria, susceptibility to
112250	Bone dysplasia with medullary fibrosarcoma
112262	Fibrodysplasia ossificans progressiva, 135100
113300	Brachydactyly type E
113705	Ovarian cancer
113705	Breast cancer-1
113721	Breast cancer
113811	Epidermolysis bullosa, generalized atrophic benign, 226650
113900	Heart block, progressive familial, type I
114208	Malignant hyperthermia susceptibility 5, 601887
114208	Hypokalemic periodic paralysis, 170400
114240	Muscular dystrophy, limb-girdle, type 2A, 253600
114290	Campomelic dysplasia with autosomal sex reversal
114350	Leukemia, acute myeloid
114400	Lynch cancer family syndrome II
114550	Hepatocellular carcinoma
114835	Monocyte carboxyesterase deficiency
115200	Cardiomyopathy, dilated, 1A
115470	Cat eye syndrome
115500	Acatlasemia
115650	Cataract, anterior polar-1
115660	Cataract, cerulean, type 1
115665	Cataract, congenital, Volkmann type
116800	Cataract, Marner type
116806	Colorectal cancer
116860	Cavernous angiomatic malformations
117700	[Hypoceruloplasminemia, hereditary]
117700	Hemosiderosis, systemic, due to aceruloplasminemia
118210	Charcot-Marie-Tooth neuropathy-2A
118425	Myotonia congenita, dominant, 160800
118425	Myotonia congenita, recessive, 255700
118425	Myotonia levior, recessive
118470	[CETP deficiency]
118485	Polycystic ovary syndrome with hyperandrogenemia
118504	Epilepsy, benign neonatal, type 1, 121200
118504	Epilepsy, nocturnal frontal lobe, 600513

118511	Schizophrenia, neurophysiologic defect in
118800	Choreoathetosis, familial paroxysmal
119300	van der Woude syndrome
120070	Alport syndrome, autosomal recessive, 203780
120110	Metaphyseal chondrodysplasia, Schmid type
120120	Epidermolysis bullosa dystrophica, dominant, 131750
120120	Epidermolysis bullosa dystrophica, recessive, 226600
120120	Epidermolysis bullosa, pretibial, 131850
120131	Alport syndrome, autosomal recessive, 203780
120131	Hematuria, familial benign
120140	Osteoarthritis, precocious
120140	SED congenita
120140	SMED Strudwick type
120140	Stickler syndrome, type I
120140	Wagner syndrome, type II
120140	Achondrogenesis-hypochondrogenesis, type II
120140	Kniest dysplasia
120150	Osteogenesis imperfecta, 4 clinical forms, 166200, 166210, 259420, 166220
120150	Osteoporosis, idiopathic, 166710
120150	Ehlers-Danlos syndrome, type VIIA1, 130060
120160	Osteogenesis imperfecta, 4 clinical forms, 166200, 166210, 259420, 166220
120160	Osteoporosis, idiopathic, 166710
120160	Ehlers-Danlos syndrome, type VIIA2, 130060
120160	Marfan syndrome, atypical
120180	Ehlers-Danlos syndrome, type III
120180	Ehlers-Danlos syndrome, type IV, 130050
120180	Fibromuscular dysplasia of arteries, 135580
120180	Aneurysm, familial, 100070
120190	Ehlers-Danlos syndrome, type I, 130000
120215	Ehlers-Danlos syndrome, type I, 130000
120215	Ehlers-Danlos syndrome, type II, 130010
120220	Bethlem myopathy, 158810
120240	Bethlem myopathy, 158810
120250	Bethlem myopathy, 158810
120260	Epiphyseal dysplasia, multiple, type 2, 600204
120280	Stickler syndrome, type III
120280	Marshall syndrome, 154780
120290	OSMED syndrome, 215150
120290	Stickler syndrome, type II, 184840
120435	Muir-Torre syndrome, 158320
120435	Colorectal cancer, hereditary, nonpolyposis, type 1 Ovarian cancer
120436	Muir-Torre family cancer syndrome, 158320
120436	Turcot syndrome with glioblastoma, 276300
120436	Colorectal cancer, hereditary nonpolyposis, type 2
120550	C1q deficiency, type A

120570	C1q deficiency, type B
120575	C1q deficiency, type C
120580	C1r/C1s deficiency, combined
120620	SLE susceptibility
120620	CR1 deficiency
120700	C3 deficiency
120810	C4 deficiency
120820	C4 deficiency
120900	C5 deficiency
120920	Measles, susceptibility to
120940	C9 deficiency
120950	C8 deficiency, type I
120960	C8 deficiency, type II
121014	Heterotaxia, visceroatrial, autosomal recessive
121050	Contractural arachnodactyly, congenital
121360	Myeloid leukemia, acute, M4Eo subtype
121800	Corneal dystrophy, crystalline, Schnyder
122720	Nicotine addiction, protection from
122720	Coumarin resistance, 122700
123000	Cranio metaphyseal dysplasia
123100	Craniosynostosis, type 1
123101	Craniosynostosis, type 2
123270	[Creatine kinase, brain type, ectopic expression of]
123580	Cataract, congenital, autosomal dominant
123620	Cataract, cerulean, type 2, 601547
123660	Cataract, Coppock-like
123829	Melanoma
123940	White sponge nevus, 193900
124020	Mephenytoin poor metabolizer
124030	Parkinsonism, susceptibility to
124030	Debrisoquine sensitivity
124200	Darier disease (keratosis follicularis)
125264	Leukemia, acute nonlymphocytic
125270	Porphyria, acute hepatic
125270	Lead poisoning, susceptibility to
125490	Dentinogenesis imperfecta-1
125660	Myopathy, desminopathic
125660	Cardiomyopathy
125852	Insulin-dependent diabetes mellitus-2
126060	Anemia, megaloblastic, due to DHFR deficiency
126337	Myxoid liposarcoma
126340	Xeroderma pigmentosum, group D, 278730
126391	DNA ligase I deficiency
126452	Autonomic nervous system dysfunction
126452	[Novelty seeking personality]
126600	Drusen, radial, autosomal dominant

126650	Chloride diarrhea, congenital, Finnish type, 214700
126650	Colon cancer
128100	Dystonia-1, torsion
129010	Neuropathy, congenital hypomyelinating, 1
129490	Ectodermal dysplasia-3, anhidrotic
129900	EEC syndrome-1
130410	Glutaricaciduria, type IIB
130500	Elliptocytosis-1
131100	Multiple endocrine neoplasia I
131100	Prolactinoma, hyperparathyroidism, carcinoid syndrome
131100	Carcinoid tumor of lung
131195	Hereditary hemorrhagic telangiectasia-1, 187300
131210	Atherosclerosis, susceptibility to
131242	Shah-Waardenburg syndrome, 277580
131400	Eosinophilia, familial
131440	Eosinophilic myeloproliferative disorder
131950	Epidermolysis bullosa, Ogna type
132700	Cylindromatosis
132810	Diphenylhydantoin toxicity
132810	Fetal hydantoin syndrome
133170	Erythremia
133171	[Erythrocytosis, familial], 133100
133200	Erythrokeratoderma variabilis
133430	Breast cancer
133430	Estrogen resistance
133450	Neuroepithelioma
133450	Ewing sarcoma
133510	Trichothiodystrophy
133510	Xeroderma pigmentosum, group B
133550	Dicarboxylicaminoaciduria, 222730
133701	Exostoses, multiple, type 2
133780	Vitreoretinopathy, exudative, familial
134370	Membroproliferative glomerulonephritis
134370	Factor H deficiency
134370	Hemolytic-uremic syndrome, 235400
134570	Factor XIII A deficiency
134580	Factor XIII B deficiency
134637	Autoimmune lymphoproliferative syndrome
134790	Hyperferritinemia-cataract syndrome, 600886
134934	Thanatophoric dysplasia, types I and II, 187600
134934	Achondroplasia, 100800
134934	Craniosynostosis, nonsyndromic
134934	Crouzon syndrome with acanthosis nigricans
134934	Hypochondroplasia, 146000
135300	Fibromatosis, gingival
135600	Ehlers-Danlos syndrome, type X

135940	Ichthyosis vulgaris, 146700
136132	[Fish-odor syndrome], 602079
136350	Pfeiffer syndrome, 101600
136435	Ovarian dysgenesis, hypergonadotropic, with normal karyotype, 233300
136530	Male infertility, familial
136836	Fucosyltransferase-6 deficiency
137350	Amyloidosis, Finnish type, 105120
138030	[Hyperproglucagonemia]
138033	Diabetes mellitus, type II
138040	Cortisol resistance
138079	Hyperinsulinism, familial, 602485
138079	MODY, type 2, 125851
138140	Glucose transport defect, blood-brain barrier
138160	Diabetes mellitus, noninsulin-dependent
138160	Fanconi-Bickel syndrome, 227810
138190	Diabetes mellitus, noninsulin-dependent
138250	P5CS deficiency
138320	Hemolytic anemia due to glutathione peroxidase deficiency
138491	Startle disease, autosomal recessive
138491	Startle disease/hyperekplexia, autosomal dominant, 149400
138491	Hyperekplexia and spastic paraparesis
138570	Non-insulin dependent diabetes mellitus, susceptibility to
138720	Bernard-Soulier syndrome, type B
138971	Kostmann neutropenia, 202700
138981	Pulmonary alveolar proteinosis, 265120
139130	Hypertension, essential, susceptibility to, 145500
139150	Basal cell carcinoma
139190	Gigantism due to GHRF hypersecretion
139190	Isolated growth hormone deficiency due to defect in GHRF
139250	Isolated growth hormone deficiency, Illig type with absent GH and Kowarski type with bioinactive GH
139320	Pituitary ACTH secreting adenoma
139320	Pseudohypoparathyroidism, type Ia, 103580
139320	Somatotrophinoma
139320	McCune-Albright polyostotic fibrous dysplasia, 174800
139330	Night blindness, congenital stationary
139350	Epidermolytic hyperkeratosis, 113800
139350	Keratoderma, palmoplantar, nonepidermolytic
139360	Pituitary ACTH-secreting adenoma
140100	[Anhaptoglobinemia]
140100	[Hypohaptoglobinemia]
141750	Alpha-thalassemia/mental retardation syndrome, type 1
141800	Methemoglobinemias, alpha-
141800	Thalassemias, alpha-
141800	Erythremias, alpha-
141800	Heinz body anemias, alpha-

141850	Thalassemia, alpha-
141850	Erythrocytosis
141850	Heinz body anemia
141850	Hemoglobin H disease
141850	Hypochromic microcytic anemia
141900	Methemoglobinemias, beta-
141900	Sickle cell anemia
141900	Thalassemias, beta-
141900	Erythremias, beta-
141900	HPFH, deletion type
141900	Heinz body anemias, beta-
142000	Thalassemia due to Hb Lepore
142000	Thalassemia, delta-
142200	HPFH, nondeletion type A
142250	HPFH, nondeletion type G
142270	Hereditary persistence of fetal hemoglobin
142335	Hereditary persistence of fetal hemoglobin, heterocellular, Indian type
142360	Thrombophilia due to heparin cofactor II deficiency
142470	[Hereditary persistence of fetal hemoglobin, heterocellular]
142640	Thrombophilia due to elevated HRG
142680	Periodic fever, familial
142857	Pemphigoid, susceptibility to
142858	Beryllium disease, chronic, susceptibility to
142946	Holoprosencephaly-4
142983	Tooth agenesis, selective, 106600
142989	Synpolydactyly, type II, 186000
143100	Huntington disease
143200	Wagner syndrome
143200	Erosive vitreoretinopathy
143890	Hypercholesterolemia, familial
144200	Epidermolytic palmoplantar keratoderma
145001	Hyperparathyroidism-jaw tumor syndrome
145260	Pseudohypoadosteronism, type II
145410	Opitz G syndrome, type II
145981	Hypocalciuric hypercalcemia, type II
146150	Hypomelanosis of Ito
146760	[IgG receptor I, phagocytic, familial deficiency of]
146790	Lupus nephritis, susceptibility to
147050	Atopy
147141	Leukemia, acute lymphoblastic
147200	[Kappa light chain deficiency]
147280	Hepatocellular carcinoma
147440	Growth retardation with deafness and mental retardation
147450	Amyotrophic lateral sclerosis, due to SOD1 deficiency, 105400
147570	Interferon, immune, deficiency
147670	Rabson-Mendenhall syndrome

147670	Diabetes mellitus, insulin-resistant, with acanthosis nigricans
147670	Leprechaunism
147680	Severe combined immunodeficiency due to IL2 deficiency
147781	Atopy, susceptibility to
147790	Leukemia, acute lymphocytic, with 4/11 translocation
147791	Jacobsen syndrome
148040	Epidermolysis bullosa simplex, Koebner, Dowling-Meara, and Weber-Cockayne types, 131900, 131760, 131800
148041	Pachyonychia congenita, Jadassohn-Lewandowsky type, 167200
148043	Meesmann corneal dystrophy, 122100
148065	White sponge nevus, 193900
148066	Epidermolysis bullosa simplex, Koebner, Dowling-Meara, and Weber-Cockayne types, 131900, 131760, 131800
148066	Epidermolysis bullosa simplex, recessive, 601001
148067	Nonepidermolytic palmoplantar keratoderma, 600962
148067	Pachyonychia congenita, Jadassohn-Lewandowsky type, 167200
148069	Pachyonychia congenita, Jackson-Lawler type, 167210
148070	Liver disease, susceptibility to, from hepatotoxins or viruses
148080	Epidermolytic hyperkeratosis, 113800
148370	Keratolytic winter erythema
148500	Tylosis with esophageal cancer
150200	[Placental lactogen deficiency]
150210	Lactoferrin-deficient neutrophils, 245480
150240	Cutis laxa, marfanoid neonatal type
150250	Larsen syndrome, autosomal dominant
150270	Laryngeal adductor paralysis
150292	Epidermolysis bullosa, Herlitz junctional type, 226700
150310	Epidermolysis bullosa, Herlitz junctional type, 226700
150310	Epidermolysis bullosa, generalized atrophic benign, 226650
151385	Leukemia, acute myeloid
151390	Leukemia, acute T-cell
151410	Leukemia, chronic myeloid
151440	Leukemia, T-cell acute lymphoblastoid
151670	Hepatic lipase deficiency
152427	Long QT syndrome-2
152445	Vohwinkel syndrome, 124500
152445	Erythrokeratoderma, progressive symmetric, 602036
152760	Hypogonadotropic hypogonadism due to GNRH deficiency, 227200
152790	Precocious puberty, male, 176410
152790	Leydig cell hypoplasia
153454	Ehlers-Danlos syndrome, type VI, 225400
153455	Cutis laxa, recessive, type I, 219100
153700	Macular dystrophy, vitelliform type
153840	Macular dystrophy, atypical vitelliform
153880	Macular dystrophy, dominant cystoid
153900	Stargardt disease-2
154275	Malignant hyperthermia susceptibility 2

154276	Malignant hyperthermia susceptibility 3
154500	Treacher Collins mandibulofacial dysostosis
154550	Carbohydrate-deficient glycoprotein syndrome, type Ib, 602579
154705	Marfan syndrome, type II
156225	Muscular dystrophy, congenital merosin-deficient
156232	Mesomelic dysplasia, Kantaputra type
156570	Methylcobalamin deficiency, cbl G type
156600	Microcoria, congenital
156845	Tietz syndrome, 103500
156845	Waardenburg syndrome, type IIA, 193510
156845	Waardenburg syndrome/ocular albinism, digenic, 103470
156850	Cataract, congenital, with microphthalmia
157147	Abetalipoproteinemia, 200100
157170	Holoprosencephaly-2
157640	PEO with mitochondrial DNA deletions, type 1
157655	Lactic acidosis due to defect in iron-sulfur cluster of complex I
157900	Moebius syndrome
159000	Muscular dystrophy, limb-girdle, type 1A
159001	Muscular dystrophy, limb-girdle, type 1B
159440	Charcot-Marie-Tooth neuropathy-1B, 118200
159440	Dejerine-Sottas disease, myelin P-related, 145900
159440	Hypomyelination, congenital
159555	Leukemia, myeloid/lymphoid or mixed-lineage
160777	Griscelli disease, 214450
160781	Cardiomyopathy, hypertrophic, mid-left ventricular chamber type
160900	Myotonic dystrophy
161015	Mitochondrial complex I deficiency, 252010
162100	Neuralgic amyotrophy with predilection for brachial plexus
162200	Neurofibromatosis, type 1
162200	Watson syndrome, 193520
162400	Neuropathy, hereditary sensory and autonomic, type 1
163729	Hypertension, pregnancy-induced
163890	Parkinson disease, type 1, 601508
164009	Leukemia, acute promyelocytic, NUMA/RARA type
164040	Leukemia, acute promyelocytic, NPM/RARA type
164160	Obesity, severe, due to leptin deficiency
164200	Oculodentodigital dysplasia
164200	Syndactyly, type III, 186100
164500	Spinocerebellar ataxia-7
164731	Ovarian carcinoma, 167000
164759	Ovarian carcinoma
164860	Renal cell carcinoma, papillary, familial and sporadic
164920	Piebaldism
164920	Mast cell leukemia
164920	Mastocytosis with associated hematologic disorder
164953	Liposarcoma

165240	Pallister-Hall syndrome, 146510
165240	Postaxial polydactyly type A1, 174200
165240	Greig cephalopolysyndactyly syndrome, 175700
165320	Hepatocellular carcinoma
165500	Optic atrophy 1
166600	Osteopetrosis, AD, type II
167000	Ovarian cancer, serous
167250	Paget disease of bone
167409	Optic nerve coloboma with renal disease, 120330
167410	Rhabdomyosarcoma, alveolar, 268220
167415	Hypothyroidism, congenital, due to thyroid dysgenesis or hypoplasia
168000	Paraganglioma, familial nonchromaffin, 1
168360	Paraneoplastic sensory neuropathy
168461	Multiple myeloma, 254250
168461	Parathyroid adenomatosis 1
168461	Centrocytic lymphoma
168468	Metaphyseal chondrodysplasia, Murk Jansen type, 156400
168500	Parietal foramina
168610	Parkinsonism-dementia with pallidopontonigral degeneration
169600	Hailey-Hailey disease
170261	Bare lymphocyte syndrome, type I, due to TAP2 deficiency
170500	Myotonia congenita, atypical acetazolamide-responsive
170500	Paramyotonia congenita, 168300
170500	Hyperkalemic periodic paralysis
170650	Periodontitis, juvenile
170995	Zellweger syndrome-2
171050	Colchicine resistance
171060	Cholestasis, progressive familial intrahepatic, type III, 602347
171190	Hypertension, essential, 145500
171650	Lysosomal acid phosphatase deficiency
171760	Hypophosphatasia, adult, 146300
171760	Hypophosphatasia, infantile, 241500
171860	Hemolytic anemia due to phosphofructokinase deficiency
172400	Hemolytic anemia due to glucosephosphate isomerase deficiency
172400	Hydrops fetalis, one form
172411	Colorectal cancer, resistance to
172430	Enolase deficiency
172471	Glycogenosis, hepatic, autosomal
172490	Phosphorylase kinase deficiency of liver and muscle, 261750
173350	Plasminogen Tochigi disease
173350	Plasminogen deficiency, types I and II
173350	Thrombophilia, dysplasminogenemic
173360	Thrombophilia due to excessive plasminogen activator inhibitor
173360	Hemorrhagic diathesis due to PAI1 deficiency
173370	Plasminogen activator deficiency
173610	Platelet alpha/delta storage pool deficiency

173850	Polio, susceptibility to
173870	Xeroderma pigmentosum
173870	Fanconi anemia
173910	Polycystic kidney disease, adult, type II
174000	Medullary cystic kidney disease, AD
174810	Osteolysis, familial expansile
174900	Polyposis, juvenile intestinal
175100	Turcot syndrome, 276300
175100	Adenomatous polyposis coli
175100	Adenomatous polyposis coli, attenuated
175100	Colorectal cancer
175100	Desmoid disease, hereditary, 135290
175100	Gardner syndrome
176000	Porphyria, acute intermittent
176100	Porphyria cutanea tarda
176100	Porphyria, hepatoerythropoietic
176260	Episodic ataxia/myokymia syndrome, 160120
176261	Jervell and Lange-Nielsen syndrome, 220400
176270	Prader-Willi syndrome
176300	[Dystransthyretinemic hyperthyroxinemia]
176300	Carpal tunnel syndrome, familial
176300	Amyloid neuropathy, familial, several allelic types
176300	Amyloidosis, senile systemic
176450	Sacral agenesis-1
176705	Breast cancer, sporadic
176730	Diabetes mellitus, rare form
176730	Hyperproinsulinemia, familial
176730	MODY, one form
176801	Metachromatic leukodystrophy due to deficiency of SAP-1
176801	Gaucher disease, variant form
176830	Obesity, adrenal insufficiency, and red hair
176830	ACTH deficiency
176860	Purpura fulminans, neonatal
176860	Thrombophilia due to protein C deficiency
176880	Protein S deficiency
176930	Dysprothrombinemia
176930	Hypoprothrombinemia
176947	Selective T-cell defect
177070	Spherocytosis, hereditary, Japanese type
177070	Hermansky-Pudlak syndrome, 203300
177400	Apnea, postanesthetic
177900	Psoriasis susceptibility-1
178300	Ptosis, hereditary congenital, 1
178600	Pulmonary hypertension, familial primary
178640	Pulmonary alveolar proteinosis, congenital, 265120
179095	Male infertility

179450	Ragweed sensitivity
179615	Reticulosis, familial histiocytic, 267700
179615	Severe combined immunodeficiency, B cell-negative, 601457
179616	Severe combined immunodeficiency, B cell-negative, 601457
179755	Renal cell carcinoma, papillary, 1
179820	[Hyperproreninemia]
180020	Retinal cone dystrophy-1
180069	Retinal dystrophy, autosomal recessive, childhood-onset
180069	Retinitis pigmentosa-20
180069	Leber congenital amaurosis-2, 204100
180071	Retinitis pigmentosa, autosomal recessive
180072	Night blindness, congenital stationary, type 3, 163500
180072	Retinitis pigmentosa, autosomal recessive
180100	Retinitis pigmentosa-1
180104	Retinitis pigmentosa-9
180105	Retinitis pigmentosa-10
180240	Leukemia, acute promyelocytic
180250	Retinol binding protein, deficiency of
180297	Anemia, hemolytic, Rh-null, suppressor type, 268150
180380	Night blindness, congenital stationery, rhodopsin-related
180380	Retinitis pigmentosa, autosomal recessive
180380	Retinitis pigmentosa-4, autosomal dominant
180381	Oguchi disease-2, 258100
180385	Leukemia, acute T-cell
180721	Retinitis pigmentosa, digenic
180840	Susceptibility to IDDM
180860	Russell-Silver syndrome
180901	Malignant hyperthermia susceptibility 1, 145600
180901	Central core disease, 117000
181030	Salivary gland pleomorphic adenoma
181031	Oguchi disease-1, 258100
181405	Scapuloperoneal spinal muscular atrophy, New England type
181430	Scapuloperoneal syndrome, myopathic type
181460	Schistosoma mansoni, susceptibility/resistance to
181510	Schizophrenia
182138	Anxiety-related personality traits
182280	Small-cell cancer of lung
182380	Glucose/galactose malabsorption
182381	Renal glucosuria, 253100
182452	Lung cancer, small cell
182500	Cataract, congenital
182600	Spastic paraplegia-3A
182601	Spastic paraplegia-4
182860	Pyropoikilocytosis
182860	Spherocytosis, recessive
182860	Elliptocytosis-2

182870	Spherocytosis-1
182870	Elliptocytosis-3
182870	Anemia, neonatal hemolytic, fatal and near-fatal
182900	Spherocytosis-2
183600	Split hand/foot malformation, type 1
185000	Stomatocytosis I
185430	Atherosclerosis, susceptibility to
185470	Myopathy due to succinate dehydrogenase deficiency
185800	Symphalangism, proximal
186580	Arthrocuteaneuveal granulomatosis
186740	Immunodeficiency due to defect in CD3-gamma
186770	Leukemia, T-cell acute lymphocytic
186780	CD3, zeta chain, deficiency
186830	Immunodeficiency, T-cell receptor/CD3 complex
186860	Leukemia/lymphoma, T-cell
186880	Leukemia/lymphoma, T-cell
187040	Leukemia-1, T-cell acute lymphoblastic
187680	6-mercaptopurine sensitivity
188025	Thrombocytopenia, Paris-Trousseau type
188070	Bleeding disorder due to defective thromboxane A2 receptor
188400	Velocardiofacial syndrome, 192430
188400	DiGeorge syndrome
188450	Goiter, adolescent multinodular
188450	Goiter, nonendemic, simple
188450	Hypothyroidism, hereditary congenital
188540	Hypothyroidism, nongoitrous
188826	Sorsby fundus dystrophy, 136900
189800	Preeclampsia/eclampsia
189980	Leukemia, chronic myeloid
190000	Atransferrinemia
190020	Bladder cancer, 109800
190040	Meningioma, SIS-related
190040	Dermatofibrosarcoma protuberans
190040	Giant-cell fibroblastoma
190100	Geniospasm
190160	Thyroid hormone resistance, 274300, 188570
190182	Colon cancer
190182	Colorectal cancer, familial nonpolyposis, type 6
190195	Ichthyosiform erythroderma, congenital, 242100
190195	Ichthyosis, lamellar, autosomal recessive, 242300
190198	Leukemia, T-cell acute lymphoblastic
190450	Hemolytic anemia due to triosephosphate isomerase deficiency
190605	Triphalangeal thumb-polysyndactyly syndrome
190685	Down syndrome
190900	Colorblindness, tritan
191030	Nemaline myopathy-1, 161800

191044	Cardiomyopathy, familial hypertrophic
191045	Cardiomyopathy, familial hypertrophic, 2, 115195
191092	Tuberous sclerosis-2
191100	Tuberous sclerosis-1
191181	Cervical carcinoma
191290	Segawa syndrome, recessive
191315	Insensitivity to pain, congenital, with anhidrosis, 256800
191540	[Urate oxidase deficiency]
192090	Ovarian carcinoma
192090	Breast cancer, lobular
192090	Endometrial carcinoma
192090	Gastric cancer, familial, 137215
192340	Diabetes insipidus, neurohypophyseal, 125700
192500	Jervell and Lange-Nielsen syndrome, 220400
192500	Long QT syndrome-1
192974	Neonatal alloimmune thrombocytopenia
192974	Glycoprotein Ia deficiency
193100	Hypophosphatemic rickets, autosomal dominant
193235	Vitreoretinopathy, neovascular inflammatory
193300	Renal cell carcinoma
193300	von Hippel-Lindau syndrome
193400	von Willebrand disease
193500	Rhabdomyosarcoma, alveolar, 268220
193500	Waardenburg syndrome, type I
193500	Waardenburg syndrome, type III, 148820
193500	Craniofacial-deafness-hand syndrome, 122880
194070	Wilms tumor, type 1
194070	Denys-Drash syndrome
194070	Frasier syndrome, 136680
194071	Wilms tumor, type 2
194071	Adrenocortical carcinoma, hereditary, 202300
194190	Wolf-Hirschhorn syndrome
200150	Choreoacanthocytosis
200350	Acetyl-CoA carboxylase deficiency
200990	Acrocallosal syndrome
201450	Acyl-CoA dehydrogenase, medium chain, deficiency of
201460	Acyl-CoA dehydrogenase, long chain, deficiency of
201470	Acyl-CoA dehydrogenase, short-chain, deficiency of
201910	Adrenal hyperplasia, congenital, due to 21-hydroxylase deficiency
202110	Adrenal hyperplasia, congenital, due to 17-alpha-hydroxylase deficiency
202200	Glucocorticoid deficiency, due to ACTH unresponsiveness
203100	Waardenburg syndrome/ocular albinism, digenic, 103470
203100	Albinism, oculocutaneous, type IA
203200	Albinism, ocular, autosomal recessive
203200	Albinism, oculocutaneous, type II
203310	Ocular albinism, autosomal recessive

203500	Alkaptonuria
203740	Alpha-ketoglutarate dehydrogenase deficiency
203750	3-ketothiolase deficiency
203800	Alstrom syndrome
204200	Ceroid-lipofuscinosis, neuronal-3, juvenile
204500	Ceroid-lipofuscinosis, neuronal 2, classic late infantile
205100	Amyotrophic lateral sclerosis, juvenile
207750	Hyperlipoproteinemia, type Ib
207800	Argininemia
208100	Arthrogryposis multiplex congenita, neurogenic
208250	Jacobs syndrome
209900	Bardet-Biedl syndrome 2
209901	Bardet-Biedl syndrome 1
214300	Klippel-Feil syndrome
215700	Citrullinemia
216550	Cohen syndrome
216900	Achromatopsia
216950	C1r/C1s deficiency, combined
217000	C2 deficiency
217050	C6 deficiency
217050	Combined C6/C7 deficiency
217070	C7 deficiency
217095	Conotruncal cardiac anomalies
217800	Macular corneal dystrophy
218000	Andermann syndrome
218030	Apparent mineralocorticoid excess, hypertension due to
219800	Cystinosis, nephropathic
221770	Polycystic lipomembranous osteodysplasia with sclerosing leukencephalopathy
221820	Gliososis, familial progressive subcortical
222100	Diabetes mellitus, insulin-dependent-1
222300	Wolfram syndrome
222600	Atelosteogenesis II, 256050
222600	Achondrogenesis Ib, 600972
222600	Diastrophic dysplasia
222700	Lysinuric protein intolerance
222800	Hemolytic anemia due to bisphosphoglycerate mutase deficiency
222900	Sucrose intolerance
223000	Lactase deficiency, adult, 223100
223000	Lactase deficiency, congenital
223360	Dopamine-beta-hydroxylase deficiency
224100	Congenital dyserythropoietic anemia II
225500	Ellis-van Creveld syndrome
227220	[Eye color, brown]
227500	Factor VII deficiency
227600	Factor X deficiency
227646	Fanconi anemia, type D

228960	[Kininogen deficiency]
229300	Friedreich ataxia
229300	Friedreich ataxia with retained reflexes
229600	Fructose intolerance
229700	Fructose-bisphosphatase deficiency
229800	[Fructosuria]
230000	Fucosidosis
230200	Galactokinase deficiency with cataracts
230350	Galactose epimerase deficiency
230400	Galactosemia
230450	Hemolytic anemia due to gamma-glutamylcysteine synthetase deficiency
230800	Gaucher disease
230800	Gaucher disease with cardiovascular calcification
231550	Achalasia-addisonianism-alacrimia syndrome
231670	Glutaricaciduria, type I
231680	Glutaricaciduria, type IIA
231950	Glutathioninuria
232000	Propionicacidemia, type I or pccA type
232050	Propionicacidemia, type II or pccB type
232200	Glycogen storage disease I
232400	Glycogen storage disease IIIa
232400	Glycogen storage disease IIIb
232500	Glycogen storage disease IV
232600	McArdle disease
232700	Glycogen storage disease VI
233100	[Renal glucosuria]
233700	Chronic granulomatous disease due to deficiency of NCF-1
233710	Chronic granulomatous disease due to deficiency of NCF-2
234000	Factor XII deficiency
234200	Neurodegeneration with brain iron accumulation
235200	Hemochromatosis
235800	[Histidinemia]
236100	Holoprosencephaly-1
236200	Homocystinuria, B6-responsive and nonresponsive types
236730	Urofacial syndrome
237300	Carbamoylphosphate synthetase I deficiency
238310	Hyperglycinemia, nonketotic, type II
238600	Chylomicronemia syndrome, familial
238600	Combined hyperlipemia, familial
238600	Hyperlipoproteinemia I
238600	Lipoprotein lipase deficiency
238970	HHH syndrome
239100	Van Buchem disease
239500	Hyperprolinemia, type I
240300	Autoimmune polyglandular disease, type I
243500	Isovalericacidemia

245050	Ketoacidosis due to SCOT deficiency
245200	Krabbe disease
245349	Lacticacidemia due to PDX1 deficiency
245900	Norum disease
245900	Fish-eye disease
246450	HMG-CoA lyase deficiency
246530	Leukotriene C4 synthase deficiency
246600	Pancreatic lipase deficiency
246900	Lipoamide dehydrogenase deficiency
247200	Miller-Dieker lissencephaly syndrome
247640	Leukemia, acute lymphoblastic
248510	Mannosidosis, beta-
248600	Maple syrup urine disease, type Ia
248610	Maple syrup urine disease, type II
248611	Maple syrup urine disease, type Ib
249000	Meckel syndrome
250100	Metachromatic leukodystrophy
250250	Cartilage-hair hypoplasia
250790	Methemoglobinemia due to cytochrome b5 deficiency
250800	Methemoglobinemia, type I
250800	Methemoglobinemia, type II
251000	Methylmalonicaciduria, mutase deficiency type
251600	Microphthalmia, autosomal recessive
252500	Mucopolipidosis II
252500	Mucopolipidosis III
252800	Mucopolysaccharidosis Ih
252800	Mucopolysaccharidosis Ih/s
252800	Mucopolysaccharidosis Is
252920	Sanfilippo syndrome, type B
252940	Sanfilippo syndrome, type D
253200	Maroteaux-Lamy syndrome, several forms
253250	Mulibrey nanism
253270	Multiple carboxylase deficiency, biotin-responsive
253601	Miyoshi myopathy, 254130
253601	Muscular dystrophy, limb-girdle, type 2B
254210	Myasthenia gravis, familial infantile
254770	Epilepsy, juvenile myoclonic
254780	Myoclonus epilepsy, Lafora type
255800	Schwartz-Jampel syndrome
256030	Nemaline myopathy-2
256100	Nephronophthisis, juvenile
256540	Galactosialidosis
256550	Sialidosis, type I
256550	Sialidosis, type II
256700	Neuroblastoma
256731	Ceroid-lipofuscinosis, neuronal-5, variant late infantile

257220	Niemann-Pick disease, type C
257220	Niemann-Pick disease, type D, 257250
258501	3-methylglutaconicaciduria, type III
259700	Osteopetrosis, recessive
259730	Renal tubular acidosis-osteopetrosis syndrome
259770	Osteoporosis-pseudoglioma syndrome
259900	Hyperoxaluria, primary, type 1
261510	Pseudo-Zellweger syndrome
261515	Peroxisomal bifunctional enzyme deficiency
261600	Phenylketonuria
261600	[Hyperphenylalaninemia, mild]
261640	Phenylketonuria due to PTS deficiency
261670	Myopathy due to phosphoglycerate mutase deficiency
261680	Hypoglycemia due to PCK1 deficiency
262000	Bjornstad syndrome
263200	Polycystic kidney disease, autosomal recessive
263700	Porphyria, congenital erythropoietic
264300	Pseudohermaphroditism, male, with gynecomastia
264470	Adrenoleukodystrophy, pseudoneonatal
264700	Pseudo-vitamin D dependency rickets 1
266100	Pyridoxine dependency with seizures
266150	Pyruvate carboxylase deficiency
266200	Anemia, hemolytic, due to PK deficiency
266600	Inflammatory bowel disease-1
267750	Knobloch syndrome
268900	[Sarcosinemia]
270100	Situs inversus viscerum
270800	Spastic paraplegia-5A
271245	Spinocerebellar ataxia-8, infantile, with sensory neuropathy
271900	Canavan disease
272750	GM2-gangliosidosis, AB variant
272800	Tay-Sachs disease
272800	[Hex A pseudodeficiency]
272800	GM2-gangliosidosis, juvenile, adult
273300	Male germ cell tumor
274180	Thromboxane synthase deficiency
274270	Thymine-uraciluria
274270	Fluorouracil toxicity, sensitivity to
274600	Pendred syndrome
274600	Deafness, autosomal recessive 4
275350	Transcobalamin II deficiency
276000	Pancreatitis, hereditary, 167800
276000	Trypsinogen deficiency
276600	Tyrosinemia, type II
276700	Tyrosinemia, type I
276900	Usher syndrome, type 1A

276901	Usher syndrome, type 2
276902	Usher syndrome, type 3
276903	Usher syndrome, type 1B
276903	Deafness, autosomal dominant 11, neurosensory, 601317
276903	Deafness, autosomal recessive 2, neurosensory, 600060
277700	Werner syndrome
277730	Wernicke-Korsakoff syndrome, susceptibility to
278000	Wolman disease
278000	Cholesteryl ester storage disease
278250	Wrinkly skin syndrome
278300	Xanthinuria, type I
300008	Nephrolithiasis, type I, 310468
300008	Proteinuria, low molecular weight, with hypercalciuric nephrocalcinosis
300008	Dent disease, 300009
300008	Hypophosphatemia, type III
300011	Menkes disease, 309400
300011	Occipital horn syndrome, 304150
300011	Cutis laxa, neonatal
300031	Mental retardation, X-linked, FRAXF type
300032	Alpha-thalassemia/mental retardation syndrome, type 2, 301040
300032	Juberg-Marsidi syndrome, 309590
300037	Simpson dysmorphia syndrome, 312870
300039	Deafness, X-linked 3, conductive, with stapes fixation, 304400
300044	Wernicke-Korsakoff syndrome, susceptibility to
300047	Mental retardation, X-linked 20
300048	Intestinal pseudoobstruction, neuronal, X-linked
300049	Nodular heterotopia, bilateral periventricular
300049	BPNH/MR syndrome
300055	Mental retardation with psychosis, pyramidal signs, and macroorchidism
300062	Mental retardation, X-linked 14
300071	Night blindness, congenital stationary, type 2
300075	Coffin-Lowry syndrome, 303600
300076	Wood neuroimmunologic syndrome
300077	Mental retardation, X-linked 29
300100	Adrenoleukodystrophy
300100	Adrenomyeloneuropathy
300104	Mental retardation, X-linked nonspecific, 309541
300110	Night blindness, congenital stationary, X-linked incomplete, 300071
300123	Mental retardation with isolated growth hormone deficiency
300126	Dyskeratosis congenita-1, 305000
300136	Diabetes mellitus, insulin-dependent, X-linked, susceptibility to
300300	XLA and isolated growth hormone deficiency, 307200
300300	Agammaglobulinemia, type 1, X-linked
300600	Ocular albinism, Forsius-Eriksson type
301000	Thrombocytopenia, X-linked, 313900

301000	Wiskott-Aldrich syndrome
301200	Amelogenesis imperfecta
301201	Amelogenesis imperfecta-3, hypoplastic type
301220	Partington syndrome II
301310	Anemia, sideroblastic, with spinocerebellar ataxia
301590	Anophthalmos-1
301830	Arthrogryposis, X-linked (spinal muscular atrophy, infantile, X-linked)
301835	Arts syndrome
301845	Bazex syndrome
301900	Borjeson-Forssman-Lehmann syndrome
302060	Noncompaction of left ventricular myocardium, isolated
302060	Barth syndrome
302060	Cardiomyopathy, X-linked dilated, 300069
302060	Endocardial fibroelastosis-2
302350	Nance-Horan syndrome
302801	Charcot-Marie-Tooth neuropathy, X-linked-2, recessive
302960	Chondrodysplasia punctata, X-linked dominant
303700	Colorblindness, blue monochromatic
303800	Colorblindness, deutan
303900	Colorblindness, protan
304020	Cone dystrophy, progressive X-linked, 1
304040	Charcot-Marie-Tooth neuropathy, X-linked-1, dominant, 302800
304340	Mental retardation, X-linked, syndromic-5, with Dandy-Walker malformation, basal ganglia disease, and seizures
304800	Diabetes insipidus, nephrogenic
305100	Anhidrotic ectodermal dysplasia
305435	Heterocellular hereditary persistence of fetal hemoglobin, Swiss type
305450	FG syndrome
305900	Favism
305900	G6PD deficiency
305900	Hemolytic anemia due to G6PD deficiency
306000	Glycogenosis, X-linked hepatic, type I
306000	Glycogenosis, X-linked hepatic, type II
306100	Gonadal dysgenesis, XY female type
306400	Chronic granulomatous disease, X-linked
306700	Hemophilia A
306995	[Homosexuality, male]
307150	Hypertrichosis, congenital generalized
307700	Hypoparathyroidism, X-linked
307800	Hypophosphatemia, hereditary
308000	HPRT-related gout
308000	Lesch-Nyhan syndrome
308230	Immunodeficiency, X-linked, with hyper-IgM
308240	Lymphoproliferative syndrome, X-linked
308310	Incontinentia pigmenti, familial
308380	Severe combined immunodeficiency, X-linked, 300400
308380	Combined immunodeficiency, X-linked, moderate, 312863

308800	Keratosis follicularis spinulosa decalvans
308840	Spastic paraplegia, 312900
308840	Hydrocephalus due to aqueductal stenosis, 307000
308840	MASA syndrome, 303350
309200	Manic-depressive illness, X-linked
309300	Megalocornea, X-linked
309470	Mental retardation, X-linked, syndromic-3, with spastic diplegia
309500	Renpenning syndrome-1
309510	Mental retardation, X-linked, syndromic-1, with dystonic movements, ataxia, and seizures
309545	Mental retardation, X-linked nonspecific, with aphasia
309548	Mental retardation, X-linked, FRAXE type
309555	Gustavson syndrome
309585	Mental retardation, X-linked, syndromic-6, with gynecomastia and obesity
309600	Allan-Herndon syndrome
309605	Mental retardation, X-linked, syndromic-4, with congenital contractures and low fingertip arches
309610	Mental retardation, X-linked, syndromic-2, with dysmorphism and cerebral atrophy
309620	Mental retardation-skeletal dysplasia
309850	Brunner syndrome
309900	Mucopolysaccharidosis II
310300	Emery-Dreifuss muscular dystrophy
310400	Myotubular myopathy, X-linked
310460	Myopia-1
310460	Bornholm eye disease
310490	Cowchock syndrome
310500	Night blindness, congenital stationary, type 1
311050	Optic atrophy, X-linked
311200	Oral-facial-digital syndrome 1
311250	Ornithine transcarbamylase deficiency
311300	Otopalatodigital syndrome, type I
311360	Ovarian failure, premature
311510	Waisman parkinsonism-mental retardation syndrome
311770	Paroxysmal nocturnal hemoglobinuria
311800	Myoglobinuria/hemolysis due to PGK deficiency
311800	Hemolytic anemia due to PGK deficiency
311850	Phosphoribosyl pyrophosphate synthetase-related gout
311870	Muscle glycogenosis
312000	Panhypopituitarism, X-linked
312040	N syndrome, 310465
312060	Properdin deficiency, X-linked
312170	Pyruvate dehydrogenase deficiency
312600	Retinitis pigmentosa-2
312610	Retinitis pigmentosa-3
312700	Retinoschisis

312760	Turner syndrome
313350	Split hand/foot malformation, type 2
313400	Spondyloepiphyseal dysplasia tarda
313850	Thoracoabdominal syndrome
314200	[Euthyroidal hyper- and hypothyroxinemia]
314250	Dystonia-3, torsion, with parkinsonism, Filipino type
314300	Goeminne TKCR syndrome
314400	Cardiac valvular dysplasia-1
314580	Wieacker-Wolff syndrome
314850	McLeod phenotype
600020	Prostate cancer, 176807
600035	Schizencephaly
600040	Colorectal cancer
600044	Thrombocythemia, essential, 187950
600045	Xeroderma pigmentosum, group E, subtype 2
600048	Breast cancer-3
600059	Retinitis pigmentosa-13
600065	Leukocyte adhesion deficiency, 116920
600079	Colon cancer
600095	Split hand/foot malformation, type 3
600101	Deafness, autosomal dominant 2
600105	Retinitis pigmentosa-12, autosomal recessive
600119	Muscular dystrophy, Duchenne-like, type 2
600119	Adhalinopathy, primary
600138	Retinitis pigmentosa-11
600140	Rubenstein-Taybi syndrome, 180849
600143	Epilepsy, progressive, with mental retardation
600151	Bardet-Biedl syndrome 3
600160	Melanoma, 155601
600163	Long QT syndrome-3
600173	SCID, autosomal recessive, T-negative/B-positive type
600175	Spinal muscular atrophy, congenital nonprogressive, of lower limbs
600179	Leber congenital amaurosis, type I, 204000
600184	Carnitine acetyltransferase deficiency
600194	Ichthyosis bullosa of Siemens, 146800
600202	Dyslexia, specific, 2
600211	Cleidocranial dysplasia, 119600
600221	Venous malformations, multiple cutaneous and mucosal, 600195
600223	Spinocerebellar ataxia-4
600225	Phenylketonuria, atypical, due to GCH1 deficiency, 233910
600225	Dystonia, DOPA-responsive, 128230
600228	Pseudohypoaldosteronism, type I, 264350
600231	Palmoplantar keratoderma, Bothnia type
600234	HMG-CoA synthase-2 deficiency
600243	Temperature-sensitive apoptosis
600258	Colorectal cancer, hereditary nonpolyposis, type 3

600261	Ehlers-Danlos-like syndrome
600266	Resistance/susceptibility to TB, etc.
600273	Polycystic kidney disease, infantile severe, with tuberous sclerosis
600276	Cerebral arteriopathy with subcortical infarcts and leukoencephalopathy, 125310
600281	Non-insulin-dependent diabetes mellitus, 125853
600281	MODY, type 1, 125850
600309	Atrioventricular canal defect-1
600310	Pseudoachondroplasia, 177170
600310	Epiphyseal dysplasia, multiple 1, 132400
600319	Diabetes mellitus, insulin-dependent, 4
600320	Insulin-dependent diabetes mellitus-5
600321	Diabetes mellitus, insulin-dependent, 7
600332	Rippling muscle disease-1
600354	Spinal muscular atrophy-1, 253300
600354	Spinal muscular atrophy-2, 253550
600354	Spinal muscular atrophy-3, 253400
600374	Bardet-Biedl syndrome 4
600414	Adrenoleukodystrophy, neonatal, 202370
600429	[Ii blood group, 110800]
600430	Brachydactyly-mental retardation syndrome
600510	Pigment dispersion syndrome
600511	Schizophrenia-3
600512	Epilepsy, partial
600525	Trichodontoosseous syndrome, 190320
600528	CPT deficiency, hepatic, type I, 255120
600536	Myopathy, congenital
600542	Chondrosarcoma, extraskeletal myxoid
600584	Atrial septal defect with atrioventricular conduction defects, 108900
600593	Craniosynostosis, Adelaide type
600617	Lipoid adrenal hyperplasia, 201710
600618	Leukemia, acute lymphoblastic
600623	Prostate cancer, 176807
600624	Cone-rod retinal dystrophy-1
600631	Enuresis, nocturnal, 1
600650	Myopathy due to CPT II deficiency, 255110
600650	CPT deficiency, hepatic, type II, 600649
600669	Epilepsy, generalized, idiopathic
600698	Salivary adenoma
600698	Uterine leiomyoma
600698	Lipoma
600698	Lipomatosis, multiple, 151900
600700	Lipoma
600701	Lipoma
600722	Ceroid lipofuscinosis, neuronal, variant juvenile type, with granular osmiophilic deposits
600722	Ceroid lipofuscinosis, neuronal-1, infantile, 256730

600725	Holoprosencephaly-3, 142945
600759	Alzheimer disease-4
600760	Pseudohypoaldosteronism, type I, 264350
600760	Liddle syndrome, 177200
600761	Pseudohypoaldosteronism, type I, 264350
600761	Liddle syndrome, 177200
600795	Dementia, familial, nonspecific
600807	Bronchial asthma
600808	Enuresis, nocturnal, 2
600811	Xeroderma pigmentosum, group E, DDB-negative subtype, 278740
600837	Hirschsprung disease, 142623
600839	Bartter syndrome, 241200
600850	Schizophrenia disorder-4
600856	Beckwith-Wiedemann syndrome, 130650
600857	Leigh syndrome
600881	Cataract, congenital, zonular, with sutural opacities
600882	Charcot-Marie-Tooth neuropathy-2B
600883	Diabetes mellitus, insulin-dependent, 8
600887	Endometrial carcinoma
600897	Cataract, zonular pulverulent-1, 116200
600900	Muscular dystrophy, limb-girdle, type 2E
600918	Cystinuria, type III
600919	Long QT syndrome-4 with sinus bradycardia
600923	Porphyria variegata, 176200
600946	Short stature, autosomal dominant, with normal serum growth hormone binding protein
600946	Short stature, idiopathic
600946	Laron dwarfism, 262500
600956	Persistent Mullerian duct syndrome, type II, 261550
600957	Persistent Mullerian duct syndrome, type I, 261550
600958	Cardiomyopathy, familial hypertrophic, 4, 115197
600965	Deafness, autosomal dominant 6
600968	Gitelman syndrome, 263800
600971	Deafness, autosomal recessive 6
600974	Deafness, autosomal recessive 7
600975	Glaucoma 3, primary infantile, B
600977	Cone dystrophy, progressive
600993	Pancreatic cancer
600995	Nephrotic syndrome, idiopathic, steroid-resistant
600996	Arrhythmogenic right ventricular dysplasia-2
600998	Bleeding diathesis due to GNAQ deficiency
601002	5-oxoprolinuria, 266130
601002	Hemolytic anemia due to glutathione synthetase deficiency, 231900
601011	Spinocerebellar ataxia-6, 183086
601011	Cerebellar ataxia, pure
601011	Episodic ataxia, type 2, 108500
601011	Hemiplegic migraine, familial, 141500

601071	Deafness, autosomal recessive 9
601072	Deafness, autosomal recessive 8
601090	Iridogoniodysgenesis, 601631
601105	Pycnodysostosis, 265800
601107	Dubin-Johnson syndrome, 237500
601130	Tolbutamide poor metabolizer
601145	Epilepsy, progressive myoclonic 1, 254800
601146	Brachydactyly, type C, 113100
601146	Acromesomelic dysplasia, Hunter-Thompson type, 201250
601146	Chondrodysplasia, Grebe type, 200700
601154	Cardiomyopathy, dilated, 1E
601199	Neonatal hyperparathyroidism, 239200
601199	Hypocalcemia, autosomal dominant, 601198
601199	Hypocalciuric hypercalcemia, type I, 145980
601202	Cataract, anterior polar-2
601208	Insulin-dependent diabetes mellitus-11
601226	Progressive external ophthalmoplegia, type 2
601238	Cerebellar ataxia, Cayman type
601267	HIV infection, susceptibility/resistance to
601277	Ichthyosis, lamellar, type 2
601282	Muscular dystrophy with epidermolysis bullosa simplex, 226670
601284	Hereditary hemorrhagic telangiectasia-2, 600376
601313	Polycystic kidney disease, adult type I, 173900
601316	Deafness, autosomal dominant 10
601318	Diabetes mellitus, insulin-dependent, 13
601363	Wilms tumor, type 4
601373	HIV infection, susceptibility/resistance to
601382	Charcot-Marie-Tooth neuropathy-4B
601385	Prostate cancer
601399	Platelet disorder, familial, with associated myeloid malignancy
601402	Leukemia, myeloid, acute
601406	B-cell non-Hodgkin lymphoma, high-grade
601410	Diabetes mellitus, transient neonatal
601411	Muscular dystrophy, limb-girdle, type 2F, 601287
601412	Deafness, autosomal dominant 7
601414	Retinitis pigmentosa-18
601455	Hereditary motor and sensory neuropathy, Lom type
601458	Inflammatory bowel disease-2
601471	Moebius syndrome-2
601493	Cardiomyopathy, dilated 1C
601494	Cardiomyopathy, familial, dilated-2
601518	Prostate cancer, hereditary, 1, 176807
601542	Rieger syndrome, type 1, 180500
601545	Lissencephaly-1
601556	Spinocerebellar ataxia-1, 164400
601567	Combined factor V and VIII deficiency, 227300

601596	Charcot-Marie-Tooth neuropathy, demyelinating
601604	Mycobacterial and salmonella infections, susceptibility to
601606	Trichoepithelioma, multiple familial
601607	Rhabdoid tumors
601620	Holt-Oram syndrome, 142900
601621	Ulnar-mammary syndrome, 181450
601622	Saethre-Chotzen syndrome, 101400
601623	Angelman syndrome
601649	Blepharophimosis, epicanthus inversus, and ptosis, type 2
601650	Paraganglioma, familial nonchromaffin, 2
601652	Glaucoma 1A, primary open angle, juvenile-onset, 137750
601666	Insulin-dependent diabetes mellitus-15
601669	Hirschsprung disease, one form
601676	Acute insulin response
601680	Distal arthrogryposis, type 2B
601682	Glaucoma 1C, primary open angle
601687	Meesmann corneal dystrophy, 122100
601690	Platelet-activating factor acetylhydrolase deficiency
601691	Retinitis pigmentosa-19, 601718
601691	Stargardt disease-1, 248200
601691	Cone-rod dystrophy 3
601691	Fundus flavimaculatus with macular dystrophy, 248200
601692	Reis-Bucklers corneal dystrophy
601692	Corneal dystrophy, Avellino type
601692	Corneal dystrophy, Groenouw type I, 121900
601692	Corneal dystrophy, lattice type I, 122200
601718	Retinitis pigmentosa-19
601744	Systemic lupus erythematosus, susceptibility to, 1
601757	Rhizomelic chondrodysplasia punctata, type 1, 215100
601768	Leukemia, acute myeloid
601769	Osteoporosis, involutional
601769	Rickets, vitamin D-resistant, 277440
601771	Glaucoma 3A, primary infantile, 231300
601777	Cone dystrophy, progressive
601780	Ceroid-lipofuscinosis, neuronal-6, variant late infantile
601785	Carbohydrate-deficient glycoprotein syndrome, type I, 212065
601800	[Hair color, brown]
601843	Hypothyroidism, congenital, 274400
601844	Pseudohypoaldosteronism type II
601846	Muscular dystrophy with rimmed vacuoles
601850	Retinitis pigmentosa-deafness syndrome
601863	Bare lymphocyte syndrome, complementation group C
601868	Deafness, autosomal dominant 13
601884	[High bone mass]
601889	Lymphoma, diffuse large cell
601916	Pancreatic cancer

601928	Monilethrix, 158000
601941	Insulin-dependent diabetes mellitus-6
601954	Muscular dystrophy, limb-girdle, type 2G
601969	Medulloblastoma, 155255
601969	Glioblastoma multiforme, 137800
601975	Ectodermal dysplasia/skin fragility syndrome
602014	Hypomagnesemia with secondary hypocalcemia
602025	Obesity/hyperinsulinism, susceptibility to
602028	Multiple myeloma
602066	Convulsions, infantile and paroxysmal choreoathetosis
602067	Cardiomyopathy, dilated, 1F
602078	Fibrosis of extraocular muscles, congenital, 2
602080	Paget disease of bone-2
602081	Speech-language disorder-1
602082	Corneal dystrophy, Thiel-Behnke type
602085	Postaxial polydactyly, type A2
602086	Arrhythmogenic right ventricular dysplasia-3
602087	Arrhythmogenic right ventricular dysplasia-4
602088	Nephronophthisis, infantile
602089	Hemangioma, capillary, hereditary
602091	Marfan syndrome, atypical
602094	Lipodystrophy, familial partial
602096	Alzheimer disease-5
602099	Amyotrophic lateral sclerosis-5
602116	Glioma
602117	Prader-Willi syndrome
602121	Deafness, autosomal dominant nonsyndromic sensorineural, 1, 124900
602134	Tremor, familial essential, 2
602136	Refsum disease, infantile, 266510
602136	Zellweger syndrome-1, 214100
602136	Adrenoleukodystrophy, neonatal, 202370
602153	Monilethrix, 158000
602216	Peutz-Jeghers syndrome, 175200
602225	Cone-rod retinal dystrophy-2, 120970
602225	Leber congenital amaurosis, type III
602229	Waardenburg-Shah syndrome, 277580
602232	Epilepsy, benign neonatal, type 2, 121201
602235	Epilepsy, benign, neonatal, type 1, 121200
602279	Oculopharyngeal muscular dystrophy, 164300
602279	Oculopharyngeal muscular dystrophy, autosomal recessive, 257950
602280	Retinitis pigmentosa-14, 600132
602363	Ellis-van Creveld-like syndrome
602397	Cholestasis, benign recurrent intrahepatic, 243300
602397	Cholestasis, progressive familial intrahepatic-1, 211600
602403	Alzheimer disease, susceptibility to
602404	Parkinson disease, type 3

602447	Coronary artery disease, susceptibility to
602460	Deafness, autosomal dominant 15, 602459
602475	Ossification of posterior longitudinal ligament of spine
602477	Febrile convulsions, familial, 2
602491	Hyperlipidemia, familial combined, 1
602522	Bartter syndrome, infantile, with sensorineural deafness
602544	Parkinson disease, juvenile, type 2, 600116
602574	Deafness, autosomal dominant 12, 601842
602574	Deafness, autosomal dominant 8, 601543
602575	Nail-patella syndrome with open-angle glaucoma, 137750
602575	Nail-patella syndrome, 161200
602594	Retinitis pigmentosa-22
602616	Carbohydrate-deficient glycoprotein syndrome, type II, 212066
602629	Dystonia-6, torsion
602631	Rhabdomyosarcoma, 268210
602631	Breast Cancer
602669	Anterior segment mesenchymal dysgenesis and cataract, 107250
602669	Cataract, congenital
602685	Mental retardation, severe, with spasticity and tapetoretinal degeneration
602716	Nephrosis-1, congenital, Finnish type, 256300
602759	Prostate cancer, hereditary, 2, 176807
602771	Muscular dystrophy, congenital, with early spine rigidity
602772	Retinitis pigmentosa-24
602782	Faisalabad histiocytosis

Mature Polypeptides

[155] The present invention also encompasses mature forms of a polypeptide having the amino acid sequence of SEQ ID NO:Y and/or the amino acid sequence encoded by the cDNA in a deposited clone. Polynucleotides encoding the mature forms (such as, for example, the polynucleotide sequence in SEQ ID NO:X and/or the polynucleotide sequence contained in the cDNA of a deposited clone) are also encompassed by the invention. Moreover, fragments or variants of these polypeptides (such as, fragments as described herein, polypeptides at least 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, or 100% identical to these polypeptides, or polypeptides encoded by a polynucleotide that hybridizes under stringent conditions to the complementary strand of the polynucleotide encoding these polypeptides) are also encompassed by the invention. In preferred embodiments, these fragments or variants retain one or more functional activities of the full-length or mature form of the polypeptide (e.g., biological activity, antigenicity [ability to bind (or compete

with a polypeptide of the invention for binding) to an anti-polypeptide of the invention antibody], immunogenicity (ability to generate antibody which binds to a specific polypeptide of the invention), ability to form multimers with polypeptides of the invention, and ability to bind to a receptor or ligand for a polypeptide of the invention). Antibodies that bind the polypeptides of the invention, and polynucleotides encoding these polypeptides are also encompassed by the invention.

[156] According to the signal hypothesis, proteins secreted by mammalian cells have a signal or secretary leader sequence which is cleaved from the mature protein once export of the growing protein chain across the rough endoplasmic reticulum has been initiated. Most mammalian cells and even insect cells cleave secreted proteins with the same specificity. However, in some cases, cleavage of a secreted protein is not entirely uniform, which results in two or more mature species of the protein. Further, it has long been known that cleavage specificity of a secreted protein is ultimately determined by the primary structure of the complete protein, that is, it is inherent in the amino acid sequence of the polypeptide.

[157] Methods for predicting whether a protein has a signal sequence, as well as the cleavage point for that sequence, are available. For instance, the method of McGeoch, *Virus Res.* 3:271-286 (1985), uses the information from a short N-terminal charged region and a subsequent uncharged region of the complete (uncleaved) protein. The method of von Heinje, *Nucleic Acids Res.* 14:4683-4690 (1986) uses the information from the residues surrounding the cleavage site, typically residues -13 to +2, where +1 indicates the amino terminus of the secreted protein. The accuracy of predicting the cleavage points of known mammalian secretory proteins for each of these methods is in the range of 75-80%. (von Heinje, *supra.*) However, the two methods do not always produce the same predicted cleavage point(s) for a given protein.

[158] In the present case, the deduced amino acid sequence of the secreted polypeptide was analyzed by a computer program called SignalP (Henrik Nielsen et al., *Protein Engineering* 10:1-6 (1997)), which predicts the cellular location of a protein based on the amino acid sequence. As part of this computational prediction of localization, the methods of McGeoch and von Heinje are incorporated. The analysis of the amino acid sequences of the secreted proteins described herein by this program provided the results shown in Table 1A.

[159] In specific embodiments, polypeptides of the invention comprise, or alternatively consist of, the predicted mature form of the polypeptide as delineated in columns 14 and 15 of Table 1A. Moreover, fragments or variants of these polypeptides (such as, fragments as

described herein, polypeptides at least 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, or 100% identical to these polypeptides, or polypeptides encoded by a polynucleotide that hybridizes under stringent conditions to the complementary strand of the polynucleotide encoding these polypeptides) are also encompassed by the invention. In preferred embodiments, these fragments or variants retain one or more functional activities of the full-length or mature form of the polypeptide (e.g., biological activity, antigenicity [ability to bind (or compete with a polypeptide of the invention for binding) to an anti-polypeptide of the invention antibody], immunogenicity (ability to generate antibody which binds to a specific polypeptide of the invention), ability to form multimers with polypeptides of the invention, and ability to bind to a receptor or ligand for a polypeptide of the invention). Antibodies that bind the polypeptides of the invention, and polynucleotides encoding these polypeptides are also encompassed by the invention.

[160] Polynucleotides encoding proteins comprising, or consisting of, the predicted mature form of polypeptides of the invention (e.g., polynucleotides having the sequence of SEQ ID NO: X (Table 1A, column 4), the sequence delineated in columns 7 and 8 of Table 1A, and a sequence encoding the mature polypeptide delineated in columns 14 and 15 of Table 1A (e.g., the sequence of SEQ ID NO:X encoding the mature polypeptide delineated in columns 14 and 15 of Table 1)) are also encompassed by the invention, as are fragments or variants of these polynucleotides (such as, fragments as described herein, polynucleotides at least 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, or 100% identical to these polynucleotides, and nucleic acids which hybridizes under stringent conditions to the complementary strand of the polynucleotide).

[161] As one of ordinary skill would appreciate, however, cleavage sites sometimes vary from organism to organism and cannot be predicted with absolute certainty. Accordingly, the present invention provides secreted polypeptides having a sequence shown in SEQ ID NO:Y which have an N-terminus beginning within 15 residues of the predicted cleavage point (i.e., having 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 more or less contiguous residues of SEQ ID NO:Y at the N-terminus when compared to the predicted mature form of the polypeptide (e.g., the mature polypeptide delineated in columns 14 and 15 of Table 1). Similarly, it is also recognized that in some cases, cleavage of the signal sequence from a secreted protein is not entirely uniform, resulting in more than one secreted species. These polypeptides, and the polynucleotides encoding such polypeptides, are contemplated by the present invention.

[162] Moreover, the signal sequence identified by the above analysis may not necessarily predict the naturally occurring signal sequence. For example, the naturally occurring signal sequence may be further upstream from the predicted signal sequence. However, it is likely that the predicted signal sequence will be capable of directing the secreted protein to the ER. Nonetheless, the present invention provides the mature protein produced by expression of the polynucleotide sequence of SEQ ID NO:X and/or the polynucleotide sequence contained in the cDNA of a deposited clone, in a mammalian cell (e.g., COS cells, as described below). These polypeptides, and the polynucleotides encoding such polypeptides, are contemplated by the present invention.

Polynucleotide and Polypeptide Variants

[163] The present invention is also directed to variants of the polynucleotide sequence disclosed in SEQ ID NO:X or the complementary strand thereto, nucleotide sequences encoding the polypeptide of SEQ ID NO:Y, the nucleotide sequence of SEQ ID NO:X that encodes the polypeptide sequence as defined in columns 13 and 14 of Table 1A, nucleotide sequences encoding the polypeptide sequence as defined in columns 13 and 14 of Table 1A, the nucleotide sequence of SEQ ID NO:X encoding the polypeptide sequence as defined in column 7 of Table 1B, nucleotide sequences encoding the polypeptide as defined in column 7 of Table 1B, the nucleotide sequence as defined in columns 8 and 9 of Table 2, nucleotide sequences encoding the polypeptide encoded by the nucleotide sequence as defined in columns 8 and 9 of Table 2, the nucleotide sequence as defined in column 6 of Table 1C, nucleotide sequences encoding the polypeptide encoded by the nucleotide sequence as defined in column 6 of Table 1C, the cDNA sequence contained in Clone ID NO:Z, nucleotide sequences encoding the polypeptide encoded by the cDNA sequence contained in Clone ID NO:Z, and/or nucleotide sequences encoding a mature (secreted) polypeptide encoded by the cDNA sequence contained in Clone ID NO:Z.

[164] The present invention also encompasses variants of the polypeptide sequence disclosed in SEQ ID NO:Y, the polypeptide as defined in columns 13 and 14 of Table 1A, the polypeptide sequence as defined in column 7 of Table 1B, a polypeptide sequence encoded by the polynucleotide sequence in SEQ ID NO:X, a polypeptide sequence encoded by the nucleotide sequence as defined in columns 8 and 9 of Table 2, a polypeptide sequence encoded by the nucleotide sequence as defined in column 6 of Table 1C, a polypeptide sequence encoded by the complement of the polynucleotide sequence in SEQ

ID NO:X, the polypeptide sequence encoded by the cDNA sequence contained in Clone ID NO:Z and/or a mature (secreted) polypeptide encoded by the cDNA sequence contained in Clone ID NO:Z.

[165] "Variant" refers to a polynucleotide or polypeptide differing from the polynucleotide or polypeptide of the present invention, but retaining essential properties thereof. Generally, variants are overall closely similar, and, in many regions, identical to the polynucleotide or polypeptide of the present invention.

[166] Thus, one aspect of the invention provides an isolated nucleic acid molecule comprising, or alternatively consisting of, a polynucleotide having a nucleotide sequence selected from the group consisting of: (a) a nucleotide sequence described in SEQ ID NO:X or contained in the cDNA sequence of Clone ID NO:Z; (b) a nucleotide sequence in SEQ ID NO:X or the cDNA in Clone ID NO:Z which encodes the complete amino acid sequence of SEQ ID NO:Y or the complete amino acid sequence encoded by the cDNA in Clone ID NO:Z; (c) a nucleotide sequence in SEQ ID NO:X or the cDNA in Clone ID NO:Z which encodes a mature polypeptide (i.e., a secreted polypeptide (e.g., as delineated in columns 14 and 15 of Table 1A)); (d) a nucleotide sequence in SEQ ID NO:X or the cDNA sequence of Clone ID NO:Z, which encodes a biologically active fragment of a polypeptide; (e) a nucleotide sequence in SEQ ID NO:X or the cDNA sequence of Clone ID NO:Z, which encodes an antigenic fragment of a polypeptide; (f) a nucleotide sequence encoding a polypeptide comprising the complete amino acid sequence of SEQ ID NO:Y or the complete amino acid sequence encoded by the cDNA in Clone ID NO:Z; (g) a nucleotide sequence encoding a mature polypeptide of the amino acid sequence of SEQ ID NO:Y (i.e., a secreted polypeptide (e.g., as delineated in columns 14 and 15 of Table 1A)) or a mature polypeptide of the amino acid sequence encoded by the cDNA in Clone ID NO:Z ; (h) a nucleotide sequence encoding a biologically active fragment of a polypeptide having the complete amino acid sequence of SEQ ID NO:Y or the complete amino acid sequence encoded by the cDNA in Clone ID NO:Z; (i) a nucleotide sequence encoding an antigenic fragment of a polypeptide having the complete amino acid sequence of SEQ ID NO:Y or the complete amino acid sequence encoded by the cDNA in Clone ID NO:Z; and (j) a nucleotide sequence complementary to any of the nucleotide sequences in (a), (b), (c), (d), (e), (f), (g), (h), or (i) above.

[167] The present invention is also directed to nucleic acid molecules which comprise, or alternatively consist of, a nucleotide sequence which is at least 80%, 85%, 90%, 95%,

96%, 97%, 98%, 99% or 100%, identical to, for example, any of the nucleotide sequences in (a), (b), (c), (d), (e), (f), (g), (h), (i), or (j) above, the nucleotide coding sequence in SEQ ID NO:X or the complementary strand thereto, the nucleotide coding sequence of the cDNA contained in Clone ID NO:Z or the complementary strand thereto, a nucleotide sequence encoding the polypeptide of SEQ ID NO:Y, a nucleotide sequence encoding a polypeptide sequence encoded by the nucleotide sequence in SEQ ID NO:X, a polypeptide sequence encoded by the complement of the polynucleotide sequence in SEQ ID NO:X, a nucleotide sequence encoding the polypeptide encoded by the cDNA contained in Clone ID NO:Z, the nucleotide coding sequence in SEQ ID NO:X as defined in columns 8 and 9 of Table 2 or the complementary strand thereto, a nucleotide sequence encoding the polypeptide encoded by the nucleotide sequence in SEQ ID NO:X as defined in columns 8 and 9 of Table 2 or the complementary strand thereto, the nucleotide coding sequence in SEQ ID NO:B as defined in column 6 of Table 1C or the complementary strand thereto, a nucleotide sequence encoding the polypeptide encoded by the nucleotide sequence in SEQ ID NO:B as defined in column 6 of Table 1C or the complementary strand thereto, the nucleotide sequence in SEQ ID NO:X encoding the polypeptide sequence as defined in column 7 of Table 1B or the complementary strand thereto, nucleotide sequences encoding the polypeptide as defined in column 7 of Table 1B or the complementary strand thereto, and/or polynucleotide fragments of any of these nucleic acid molecules (e.g., those fragments described herein). Polynucleotides which hybridize to the complement of these nucleic acid molecules under stringent hybridization conditions or alternatively, under lower stringency conditions, are also encompassed by the invention, as are polypeptides encoded by these polynucleotides and nucleic acids.

[168] In a preferred embodiment, the invention encompasses nucleic acid molecules which comprise, or alternatively, consist of a polynucleotide which hybridizes under stringent hybridization conditions, or alternatively, under lower stringency conditions, to a polynucleotide in (a), (b), (c), (d), (e), (f), (g), (h), or (i), above, as are polypeptides encoded by these polynucleotides. In another preferred embodiment, polynucleotides which hybridize to the complement of these nucleic acid molecules under stringent hybridization conditions, or alternatively, under lower stringency conditions, are also encompassed by the invention, as are polypeptides encoded by these polynucleotides.

[169] In another embodiment, the invention provides a purified protein comprising, or alternatively consisting of, a polypeptide having an amino acid sequence selected from the

group consisting of: (a) the complete amino acid sequence of SEQ ID NO:Y or the complete amino acid sequence encoded by the cDNA in Clone ID NO:Z; (b) the amino acid sequence of a mature (secreted) form of a polypeptide having the amino acid sequence of SEQ ID NO:Y (e.g., as delineated in columns 14 and 15 of Table 1A) or a mature form of the amino acid sequence encoded by the cDNA in Clone ID NO:Z mature; (c) the amino acid sequence of a biologically active fragment of a polypeptide having the complete amino acid sequence of SEQ ID NO:Y or the complete amino acid sequence encoded by the cDNA in Clone ID NO:Z; and (d) the amino acid sequence of an antigenic fragment of a polypeptide having the complete amino acid sequence of SEQ ID NO:Y or the complete amino acid sequence encoded by the cDNA in Clone ID NO:Z.

[170] The present invention is also directed to proteins which comprise, or alternatively consist of, an amino acid sequence which is at least 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100%, identical to, for example, any of the amino acid sequences in (a), (b), (c), or (d), above, the amino acid sequence shown in SEQ ID NO:Y, the amino acid sequence encoded by the cDNA contained in Clone ID NO:Z, the amino acid sequence of the polypeptide encoded by the nucleotide sequence in SEQ ID NO:X as defined in columns 8 and 9 of Table 2, the amino acid sequence of the polypeptide encoded by the nucleotide sequence in SEQ ID NO:B as defined in column 6 of Table 1C, the amino acid sequence as defined in column 7 of Table 1B, an amino acid sequence encoded by the nucleotide sequence in SEQ ID NO:X, and an amino acid sequence encoded by the complement of the polynucleotide sequence in SEQ ID NO:X. Fragments of these polypeptides are also provided (e.g., those fragments described herein). Further proteins encoded by polynucleotides which hybridize to the complement of the nucleic acid molecules encoding these amino acid sequences under stringent hybridization conditions or alternatively, under lower stringency conditions, are also encompassed by the invention, as are the polynucleotides encoding these proteins.

[171] By a nucleic acid having a nucleotide sequence at least, for example, 95% "identical" to a reference nucleotide sequence of the present invention, it is intended that the nucleotide sequence of the nucleic acid is identical to the reference sequence except that the nucleotide sequence may include up to five point mutations per each 100 nucleotides of the reference nucleotide sequence encoding the polypeptide. In other words, to obtain a nucleic acid having a nucleotide sequence at least 95% identical to a reference nucleotide sequence, up to 5% of the nucleotides in the reference sequence may be deleted or substituted with

another nucleotide, or a number of nucleotides up to 5% of the total nucleotides in the reference sequence may be inserted into the reference sequence. The query sequence may be an entire sequence referred to in Table 1B or 2 as the ORF (open reading frame), or any fragment specified as described herein.

[172] As a practical matter, whether any particular nucleic acid molecule or polypeptide is at least 80%, 85%, 90%, 95%, 96%, 97%, 98% or 99% identical to a nucleotide sequence of the present invention can be determined conventionally using known computer programs. A preferred method for determining the best overall match between a query sequence (a sequence of the present invention) and a subject sequence, also referred to as a global sequence alignment, can be determined using the FASTDB computer program based on the algorithm of Brutlag et al. (Comp. App. Biosci. 6:237-245 (1990)). In a sequence alignment the query and subject sequences are both DNA sequences. An RNA sequence can be compared by converting U's to T's. The result of said global sequence alignment is expressed as percent identity. Preferred parameters used in a FASTDB alignment of DNA sequences to calculate percent identity are: Matrix=Unitary, k-tuple=4, Mismatch Penalty=1, Joining Penalty=30, Randomization Group Length=0, Cutoff Score=1, Gap Penalty=5, Gap Size Penalty 0.05, Window Size=500 or the length of the subject nucleotide sequence, whichever is shorter.

[173] If the subject sequence is shorter than the query sequence because of 5' or 3' deletions, not because of internal deletions, a manual correction must be made to the results. This is because the FASTDB program does not account for 5' and 3' truncations of the subject sequence when calculating percent identity. For subject sequences truncated at the 5' or 3' ends, relative to the query sequence, the percent identity is corrected by calculating the number of bases of the query sequence that are 5' and 3' of the subject sequence, which are not matched/aligned, as a percent of the total bases of the query sequence. Whether a nucleotide is matched/aligned is determined by results of the FASTDB sequence alignment. This percentage is then subtracted from the percent identity, calculated by the above FASTDB program using the specified parameters, to arrive at a final percent identity score. This corrected score is what is used for the purposes of the present invention. Only bases outside the 5' and 3' bases of the subject sequence, as displayed by the FASTDB alignment, which are not matched/aligned with the query sequence, are calculated for the purposes of manually adjusting the percent identity score.

[174] For example, a 90 base subject sequence is aligned to a 100 base query sequence

to determine percent identity. The deletions occur at the 5' end of the subject sequence and therefore, the FASTDB alignment does not show a matched/alignment of the first 10 bases at 5' end. The 10 unpaired bases represent 10% of the sequence (number of bases at the 5' and 3' ends not matched/total number of bases in the query sequence) so 10% is subtracted from the percent identity score calculated by the FASTDB program. If the remaining 90 bases were perfectly matched the final percent identity would be 90%. In another example, a 90 base subject sequence is compared with a 100 base query sequence. This time the deletions are internal deletions so that there are no bases on the 5' or 3' of the subject sequence which are not matched/aligned with the query. In this case the percent identity calculated by FASTDB is not manually corrected. Once again, only bases 5' and 3' of the subject sequence which are not matched/aligned with the query sequence are manually corrected for. No other manual corrections are to be made for the purposes of the present invention.

[175] By a polypeptide having an amino acid sequence at least, for example, 95% "identical" to a query amino acid sequence of the present invention, it is intended that the amino acid sequence of the subject polypeptide is identical to the query sequence except that the subject polypeptide sequence may include up to five amino acid alterations per each 100 amino acids of the query amino acid sequence. In other words, to obtain a polypeptide having an amino acid sequence at least 95% identical to a query amino acid sequence, up to 5% of the amino acid residues in the subject sequence may be inserted, deleted, (indels) or substituted with another amino acid. These alterations of the reference sequence may occur at the amino or carboxy terminal positions of the reference amino acid sequence or anywhere between those terminal positions, interspersed either individually among residues in the reference sequence or in one or more contiguous groups within the reference sequence.

[176] As a practical matter, whether any particular polypeptide is at least 80%, 85%, 90%, 95%, 96%, 97%, 98% or 99% identical to, for instance, the amino acid sequence of a polypeptide referred to in Table 1A (e.g., the amino acid sequence delineated in columns 14 and 15) or a fragment thereof, Table 1B (e.g., the amino acid sequence identified in column 6) or a fragment thereof, Table 2 (e.g., the amino acid sequence of the polypeptide encoded by the polynucleotide sequence defined in columns 8 and 9 of Table 2) or a fragment thereof, the amino acid sequence of the polypeptide encoded by the polynucleotide sequence in SEQ ID NO:B as defined in column 6 of Table 1C or a fragment thereof, the

amino acid sequence of the polypeptide encoded by the nucleotide sequence in SEQ ID NO:X or a fragment thereof, or the amino acid sequence of the polypeptide encoded by cDNA contained in Clone ID NO:Z, or a fragment thereof, the amino acid sequence of a mature (secreted) polypeptide encoded by cDNA contained in Clone ID NO:Z, or a fragment thereof, can be determined conventionally using known computer programs. A preferred method for determining the best overall match between a query sequence (a sequence of the present invention) and a subject sequence, also referred to as a global sequence alignment, can be determined using the FASTDB computer program based on the algorithm of Brutlag et al. (Comp. App. Biosci.6:237-245 (1990)). In a sequence alignment the query and subject sequences are either both nucleotide sequences or both amino acid sequences. The result of said global sequence alignment is expressed as percent identity. Preferred parameters used in a FASTDB amino acid alignment are: Matrix=PAM 0, k-tuple=2, Mismatch Penalty=1, Joining Penalty=20, Randomization Group Length=0, Cutoff Score=1, Window Size=sequence length, Gap Penalty=5, Gap Size Penalty=0.05, Window Size=500 or the length of the subject amino acid sequence, whichever is shorter.

[177] If the subject sequence is shorter than the query sequence due to N- or C-terminal deletions, not because of internal deletions, a manual correction must be made to the results. This is because the FASTDB program does not account for N- and C-terminal truncations of the subject sequence when calculating global percent identity. For subject sequences truncated at the N- and C-termini, relative to the query sequence, the percent identity is corrected by calculating the number of residues of the query sequence that are N- and C-terminal of the subject sequence, which are not matched/aligned with a corresponding subject residue, as a percent of the total bases of the query sequence. Whether a residue is matched/aligned is determined by results of the FASTDB sequence alignment. This percentage is then subtracted from the percent identity, calculated by the above FASTDB program using the specified parameters, to arrive at a final percent identity score. This final percent identity score is what is used for the purposes of the present invention. Only residues to the N- and C-termini of the subject sequence, which are not matched/aligned with the query sequence, are considered for the purposes of manually adjusting the percent identity score. That is, only query residue positions outside the farthest N- and C- terminal residues of the subject sequence.

[178] For example, a 90 amino acid residue subject sequence is aligned with a 100 residue query sequence to determine percent identity. The deletion occurs at the N-

of biological function. As an example, Ron et al. (J. Biol. Chem. 268: 2984-2988 (1993)) reported variant KGF proteins having heparin binding activity even after deleting 3, 8, or 27 amino-terminal amino acid residues. Similarly, Interferon gamma exhibited up to ten times higher activity after deleting 8-10 amino acid residues from the carboxy terminus of this protein. (Dobeli et al., J. Biotechnology 7:199-216 (1988).)

[182] Moreover, ample evidence demonstrates that variants often retain a biological activity similar to that of the naturally occurring protein. For example, Gayle and coworkers (J. Biol. Chem. 268:22105-22111 (1993)) conducted extensive mutational analysis of human cytokine IL-1a. They used random mutagenesis to generate over 3,500 individual IL-1a mutants that averaged 2.5 amino acid changes per variant over the entire length of the molecule. Multiple mutations were examined at every possible amino acid position. The investigators found that "[m]ost of the molecule could be altered with little effect on either [binding or biological activity]." In fact, only 23 unique amino acid sequences, out of more than 3,500 nucleotide sequences examined, produced a protein that significantly differed in activity from wild-type.

[183] Furthermore, even if deleting one or more amino acids from the N-terminus or C-terminus of a polypeptide results in modification or loss of one or more biological functions, other biological activities may still be retained. For example, the ability of a deletion variant to induce and/or to bind antibodies which recognize the secreted form will likely be retained when less than the majority of the residues of the secreted form are removed from the N-terminus or C-terminus. Whether a particular polypeptide lacking N- or C-terminal residues of a protein retains such immunogenic activities can readily be determined by routine methods described herein and otherwise known in the art.

[184] Thus, the invention further includes polypeptide variants which show a functional activity (e.g., biological activity) of the polypeptides of the invention. Such variants include deletions, insertions, inversions, repeats, and substitutions selected according to general rules known in the art so as have little effect on activity.

[185] The present application is directed to nucleic acid molecules at least 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% identical to the nucleic acid sequences disclosed herein, (e.g., encoding a polypeptide having the amino acid sequence of an N and/or C terminal deletion), irrespective of whether they encode a polypeptide having functional activity. This is because even where a particular nucleic acid molecule does not encode a polypeptide having functional activity, one of skill in the art would still know how to use

the nucleic acid molecule, for instance, as a hybridization probe or a polymerase chain reaction (PCR) primer. Uses of the nucleic acid molecules of the present invention that do not encode a polypeptide having functional activity include, inter alia, (1) isolating a gene or allelic or splice variants thereof in a cDNA library; (2) *in situ* hybridization (e.g., "FISH") to metaphase chromosomal spreads to provide precise chromosomal location of the gene, as described in Verma et al., Human Chromosomes: A Manual of Basic Techniques, Pergamon Press, New York (1988); (3) Northern Blot analysis for detecting mRNA expression in specific tissues (e.g., normal or diseased tissues); and (4) *in situ* hybridization (e.g., histochemistry) for detecting mRNA expression in specific tissues (e.g., normal or diseased tissues).

[186] Preferred, however, are nucleic acid molecules having sequences at least 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% identical to the nucleic acid sequences disclosed herein, which do, in fact, encode a polypeptide having functional activity. By a polypeptide having "functional activity" is meant, a polypeptide capable of displaying one or more known functional activities associated with a full-length (complete) protein and/or a mature (secreted) protein of the invention. Such functional activities include, but are not limited to, biological activity, antigenicity [ability to bind (or compete with a polypeptide of the invention for binding) to an anti-polypeptide of the invention antibody], immunogenicity (ability to generate antibody which binds to a specific polypeptide of the invention), ability to form multimers with polypeptides of the invention, and ability to bind to a receptor or ligand for a polypeptide of the invention.

[187] The functional activity of the polypeptides, and fragments, variants and derivatives of the invention, can be assayed by various methods.

[188] For example, in one embodiment where one is assaying for the ability to bind or compete with a full-length polypeptide of the present invention for binding to an anti-polypeptide antibody, various immunoassays known in the art can be used, including but not limited to, competitive and non-competitive assay systems using techniques such as radioimmunoassays, ELISA (enzyme linked immunosorbent assay), "sandwich" immunoassays, immunoradiometric assays, gel diffusion precipitation reactions, immunodiffusion assays, *in situ* immunoassays (using colloidal gold, enzyme or radioisotope labels, for example), western blots, precipitation reactions, agglutination assays (e.g., gel agglutination assays, hemagglutination assays), complement fixation assays, immunofluorescence assays, protein A assays, and immunoelectrophoresis assays,

etc. In one embodiment, antibody binding is detected by detecting a label on the primary antibody. In another embodiment, the primary antibody is detected by detecting binding of a secondary antibody or reagent to the primary antibody. In a further embodiment, the secondary antibody is labeled. Many means are known in the art for detecting binding in an immunoassay and are within the scope of the present invention.

[189] In another embodiment, where a ligand is identified, or the ability of a polypeptide fragment, variant or derivative of the invention to multimerize is being evaluated, binding can be assayed, e.g., by means well-known in the art, such as, for example, reducing and non-reducing gel chromatography, protein affinity chromatography, and affinity blotting. See generally, Phizicky et al., Microbiol. Rev. 59:94-123 (1995). In another embodiment, the ability of physiological correlates of a polypeptide of the present invention to bind to a substrate(s) of the polypeptide of the invention can be routinely assayed using techniques known in the art.

[190] In addition, assays described herein (see Examples) and otherwise known in the art may routinely be applied to measure the ability of polypeptides of the present invention and fragments, variants and derivatives thereof to elicit polypeptide related biological activity (either *in vitro* or *in vivo*). Other methods will be known to the skilled artisan and are within the scope of the invention.

[191] Of course, due to the degeneracy of the genetic code, one of ordinary skill in the art will immediately recognize that a large number of the nucleic acid molecules having a sequence at least 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, or 100% identical to, for example, the nucleic acid sequence of the cDNA contained in Clone ID NO:Z, the nucleic acid sequence referred to in Table 1B (SEQ ID NO:X),), the nucleic acid sequence disclosed in Table 1A (e.g., the nucleic acid sequence delineated in columns 7 and 8), the nucleic acid sequence disclosed in Table 2 (e.g., the nucleic acid sequence delineated in columns 8 and 9) or fragments thereof, will encode polypeptides "having functional activity." In fact, since degenerate variants of any of these nucleotide sequences all encode the same polypeptide, in many instances, this will be clear to the skilled artisan even without performing the above described comparison assay. It will be further recognized in the art that, for such nucleic acid molecules that are not degenerate variants, a reasonable number will also encode a polypeptide having functional activity. This is because the skilled artisan is fully aware of amino acid substitutions that are either less likely or not likely to significantly effect protein function (e.g., replacing one aliphatic amino acid with a

second aliphatic amino acid), as further described below.

[192] For example, guidance concerning how to make phenotypically silent amino acid substitutions is provided in Bowie et al., "Deciphering the Message in Protein Sequences: Tolerance to Amino Acid Substitutions," *Science* 247:1306-1310 (1990), wherein the authors indicate that there are two main strategies for studying the tolerance of an amino acid sequence to change.

[193] The first strategy exploits the tolerance of amino acid substitutions by natural selection during the process of evolution. By comparing amino acid sequences in different species, conserved amino acids can be identified. These conserved amino acids are likely important for protein function. In contrast, the amino acid positions where substitutions have been tolerated by natural selection indicates that these positions are not critical for protein function. Thus, positions tolerating amino acid substitution could be modified while still maintaining biological activity of the protein.

[194] The second strategy uses genetic engineering to introduce amino acid changes at specific positions of a cloned gene to identify regions critical for protein function. For example, site directed mutagenesis or alanine-scanning mutagenesis (introduction of single alanine mutations at every residue in the molecule) can be used. See Cunningham and Wells, *Science* 244:1081-1085 (1989). The resulting mutant molecules can then be tested for biological activity.

[195] As the authors state, these two strategies have revealed that proteins are surprisingly tolerant of amino acid substitutions. The authors further indicate which amino acid changes are likely to be permissive at certain amino acid positions in the protein. For example, most buried (within the tertiary structure of the protein) amino acid residues require nonpolar side chains, whereas few features of surface side chains are generally conserved. Moreover, tolerated conservative amino acid substitutions involve replacement of the aliphatic or hydrophobic amino acids Ala, Val, Leu and Ile; replacement of the hydroxyl residues Ser and Thr; replacement of the acidic residues Asp and Glu; replacement of the amide residues Asn and Gln, replacement of the basic residues Lys, Arg, and His; replacement of the aromatic residues Phe, Tyr, and Trp, and replacement of the small-sized amino acids Ala, Ser, Thr, Met, and Gly.

[196] Besides conservative amino acid substitution, variants of the present invention include (i) substitutions with one or more of the non-conserved amino acid residues, where the substituted amino acid residues may or may not be one encoded by the genetic code, or

(ii) substitutions with one or more of the amino acid residues having a substituent group, or (iii) fusion of the mature polypeptide with another compound, such as a compound to increase the stability and/or solubility of the polypeptide (for example, polyethylene glycol), (iv) fusion of the polypeptide with additional amino acids, such as, for example, an IgG Fc fusion region peptide, serum albumin (preferably human serum albumin) or a fragment thereof, or leader or secretory sequence, or a sequence facilitating purification, or (v) fusion of the polypeptide with another compound, such as albumin (including but not limited to recombinant albumin (see, e.g., U.S. Patent No. 5,876,969, issued March 2, 1999, EP Patent 0 413 622, and U.S. Patent No. 5,766,883, issued June 16, 1998, herein incorporated by reference in their entirety)). Such variant polypeptides are deemed to be within the scope of those skilled in the art from the teachings herein.

[197] For example, polypeptide variants containing amino acid substitutions of charged amino acids with other charged or neutral amino acids may produce proteins with improved characteristics, such as less aggregation. Aggregation of pharmaceutical formulations both reduces activity and increases clearance due to the aggregate's immunogenic activity. See Pinckard et al., Clin. Exp. Immunol. 2:331-340 (1967); Robbins et al., Diabetes 36: 838-845 (1987); Cleland et al., Crit. Rev. Therapeutic Drug Carrier Systems 10:307-377 (1993).

[198] A further embodiment of the invention relates to polypeptides which comprise the amino acid sequence of a polypeptide having an amino acid sequence which contains at least one amino acid substitution, but not more than 50 amino acid substitutions, even more preferably, not more than 40 amino acid substitutions, still more preferably, not more than 30 amino acid substitutions, and still even more preferably, not more than 20 amino acid substitutions from a polypeptide sequence disclosed herein. Of course it is highly preferable for a polypeptide to have an amino acid sequence which, for example, comprises the amino acid sequence of a polypeptide of SEQ ID NO:Y, the amino acid sequence of the mature (e.g., secreted) polypeptide of SEQ ID NO:Y, an amino acid sequence encoded by SEQ ID NO:X, an amino acid sequence encoded by the portion of SEQ ID NO:X as defined in columns 8 and 9 of Table 2, an amino acid sequence encoded by the complement of SEQ ID NO:X, an amino acid sequence encoded by cDNA contained in Clone ID NO:Z, and/or the amino acid sequence of a mature (secreted) polypeptide encoded by cDNA contained in Clone ID NO:Z, or a fragment thereof, which contains, in order of ever-increasing preference, at least one, but not more than 10, 9, 8, 7, 6, 5, 4, 3, 2 or 1 amino acid substitutions.

[199] In specific embodiments, the polypeptides of the invention comprise, or alternatively, consist of, fragments or variants of a reference amino acid sequence selected from: (a) the amino acid sequence of SEQ ID NO:Y or fragments thereof (e.g., the mature form and/or other fragments described herein); (b) the amino acid sequence encoded by SEQ ID NO:X or fragments thereof; (c) the amino acid sequence encoded by the complement of SEQ ID NO:X or fragments thereof; (d) the amino acid sequence encoded by the portion of SEQ ID NO:X as defined in columns 8 and 9 of Table 2 or fragments thereof; and (e) the amino acid sequence encoded by cDNA contained in Clone ID NO:Z or fragments thereof; wherein the fragments or variants have 1-5, 5-10, 5-25, 5-50, 10-50 or 50-150, amino acid residue additions, substitutions, and/or deletions when compared to the reference amino acid sequence. In preferred embodiments, the amino acid substitutions are conservative. Polynucleotides encoding these polypeptides are also encompassed by the invention.

Polynucleotide and Polypeptide Fragments

[200] The present invention is also directed to polynucleotide fragments of the polynucleotides (nucleic acids) of the invention. In the present invention, a "polynucleotide fragment" refers to a polynucleotide having a nucleic acid sequence which, for example: is a portion of the cDNA contained in Clone ID NO:Z or the complementary strand thereto; is a portion of the polynucleotide sequence encoding the polypeptide encoded by the cDNA contained in Clone ID NO:Z or the complementary strand thereto; is a portion of the polynucleotide sequence encoding the mature (secreted) polypeptide encoded by the cDNA contained in Clone ID NO:Z or the complementary strand thereto; is a portion of a polynucleotide sequence encoding the mature amino acid sequence as defined in columns 14 and 15 of Table 1A or the complementary strand thereto; is a portion of a polynucleotide sequence encoding the amino acid sequence encoded by the region of SEQ ID NO:X as defined in columns 8 and 9 of Table 2 or the complementary strand thereto; is a portion of the polynucleotide sequence of SEQ ID NO:X as defined in columns 8 and 9 of Table 2 or the complementary strand thereto; is a portion of the polynucleotide sequence in SEQ ID NO:X or the complementary strand thereto; is a polynucleotide sequence encoding a portion of the polypeptide of SEQ ID NO:Y; is a polynucleotide sequence encoding a portion of a polypeptide encoded by SEQ ID NO:X; is a polynucleotide sequence encoding a portion of a polypeptide encoded by the complement of the polynucleotide sequence in SEQ ID NO:X; is a portion of a polynucleotide sequence encoding the amino acid sequence

0950083-091201

encoded by the region of SEQ ID NO:B as defined in column 6 of Table 1C or the complementary strand thereto; or is a portion of the polynucleotide sequence of SEQ ID NO:B as defined in column 6 of Table 1C or the complementary strand thereto.

[201] The polynucleotide fragments of the invention are preferably at least about 15 nt, and more preferably at least about 20 nt, still more preferably at least about 30 nt, and even more preferably, at least about 40 nt, at least about 50 nt, at least about 75 nt, or at least about 150 nt in length. A fragment "at least 20 nt in length," for example, is intended to include 20 or more contiguous bases from the cDNA sequence contained in Clone ID NO:Z, or the nucleotide sequence shown in SEQ ID NO:X or the complementary stand thereto. In this context "about" includes the particularly recited value or a value larger or smaller by several (5, 4, 3, 2, or 1) nucleotides, at either terminus or at both termini. These nucleotide fragments have uses that include, but are not limited to, as diagnostic probes and primers as discussed herein. Of course, larger fragments (e.g., at least 160, 170, 180, 190, 200, 250, 500, 600, 1000, or 2000 nucleotides in length) are also encompassed by the invention.

[202] Moreover, representative examples of polynucleotide fragments of the invention comprise, or alternatively consist of, a sequence from about nucleotide number 1-50, 51-100, 101-150, 151-200, 201-250, 251-300, 301-350, 351-400, 401-450, 451-500, 501-550, 551-600, 601-650, 651-700, 701-750, 751-800, 801-850, 851-900, 901-950, 951-1000, 1001-1050, 1051-1100, 1101-1150, 1151-1200, 1201-1250, 1251-1300, 1301-1350, 1351-1400, 1401-1450, 1451-1500, 1501-1550, 1551-1600, 1601-1650, 1651-1700, 1701-1750, 1751-1800, 1801-1850, 1851-1900, 1901-1950, 1951-2000, 2001-2050, 2051-2100, 2101-2150, 2151-2200, 2201-2250, 2251-2300, 2301-2350, 2351-2400, 2401-2450, 2451-2500, 2501-2550, 2551-2600, 2601-2650, 2651-2700, 2701-2750, 2751-2800, 2801-2850, 2851-2900, 2901-2950, 2951-3000, 3001-3050, 3051-3100, 3101-3150, 3151-3200, 3201-3250, 3251-3300, 3301-3350, 3351-3400, 3401-3450, 3451-3500, 3501-3550, 3551-3600, 3601-3650, 3651-3700, 3701-3750, 3751-3800, 3801-3850, 3851-3900, 3901-3950, 3951-4000, 4001-4050, 4051-4100, 4101-4150, 4151-4200, 4201-4250, 4251-4300, 4301-4350, 4351-4400, 4401-4450, 4451-4500, 4501-4550, 4551-4600, 4601-4650, 4651-4700, 4701-4750, 4751-4800, 4801-4850, 4851-4900, 4901-4950, 4951-5000, 5001-5050, 5051-5100, 5101-5150, 5151-5200, 5201-5250, 5251-5300, 5301-5350, 5351-5400, 5401-5450, 5451-5500, 5501-5550, 5551-5600, 5601-5650, 5651-5700, 5701-5750, 5751-5800, 5801-5850, 5851-5900, 5901-5950, 5951-6000, 6001-6050, 6051-6100, 6101-6150, 6151-6200, 6201-6250,

6251-6300, 6301-6350, 6351-6400, 6401-6450, 6451-6500, 6501-6550, 6551-6600, 6601-6650, 6651-6700, 6701-6750, 6751-6800, 6801-6850, 6851-6900, 6901-6950, 6951-7000, 7001-7050, 7051-7100, 7101-7150, 7151-7200, 7201-7250, 7251-7300 or 7301 to the end of SEQ ID NO:X, or the complementary strand thereto. In this context "about" includes the particularly recited range or a range larger or smaller by several (5, 4, 3, 2, or 1) nucleotides, at either terminus or at both termini. Preferably, these fragments encode a polypeptide which has a functional activity (e.g., biological activity). More preferably, these polynucleotides can be used as probes or primers as discussed herein. Polynucleotides which hybridize to one or more of these polynucleotides under stringent hybridization conditions or alternatively, under lower stringency conditions are also encompassed by the invention, as are polypeptides encoded by these polynucleotides.

[203] Further representative examples of polynucleotide fragments of the invention comprise, or alternatively consist of, a sequence from about nucleotide number 1-50, 51-100, 101-150, 151-200, 201-250, 251-300, 301-350, 351-400, 401-450, 451-500, 501-550, 551-600, 601-650, 651-700, 701-750, 751-800, 801-850, 851-900, 901-950, 951-1000, 1001-1050, 1051-1100, 1101-1150, 1151-1200, 1201-1250, 1251-1300, 1301-1350, 1351-1400, 1401-1450, 1451-1500, 1501-1550, 1551-1600, 1601-1650, 1651-1700, 1701-1750, 1751-1800, 1801-1850, 1851-1900, 1901-1950, 1951-2000, 2001-2050, 2051-2100, 2101-2150, 2151-2200, 2201-2250, 2251-2300, 2301-2350, 2351-2400, 2401-2450, 2451-2500, 2501-2550, 2551-2600, 2601-2650, 2651-2700, 2701-2750, 2751-2800, 2801-2850, 2851-2900, 2901-2950, 2951-3000, 3001-3050, 3051-3100, 3101-3150, 3151-3200, 3201-3250, 3251-3300, 3301-3350, 3351-3400, 3401-3450, 3451-3500, 3501-3550, 3551-3600, 3601-3650, 3651-3700, 3701-3750, 3751-3800, 3801-3850, 3851-3900, 3901-3950, 3951-4000, 4001-4050, 4051-4100, 4101-4150, 4151-4200, 4201-4250, 4251-4300, 4301-4350, 4351-4400, 4401-4450, 4451-4500, 4501-4550, 4551-4600, 4601-4650, 4651-4700, 4701-4750, 4751-4800, 4801-4850, 4851-4900, 4901-4950, 4951-5000, 5001-5050, 5051-5100, 5101-5150, 5151-5200, 5201-5250, 5251-5300, 5301-5350, 5351-5400, 5401-5450, 5451-5500, 5501-5550, 5551-5600, 5601-5650, 5651-5700, 5701-5750, 5751-5800, 5801-5850, 5851-5900, 5901-5950, 5951-6000, 6001-6050, 6051-6100, 6101-6150, 6151-6200, 6201-6250, 6251-6300, 6301-6350, 6351-6400, 6401-6450, 6451-6500, 6501-6550, 6551-6600, 6601-6650, 6651-6700, 6701-6750, 6751-6800, 6801-6850, 6851-6900, 6901-6950, 6951-7000, 7001-7050, 7051-7100, 7101-7150, 7151-7200, 7201-7250, 7251-7300 or 7301 to the end of the cDNA sequence contained in Clone ID NO:Z, or the complementary strand thereto.

In this context "about" includes the particularly recited range or a range larger or smaller by several (5, 4, 3, 2, or 1) nucleotides, at either terminus or at both termini. Preferably, these fragments encode a polypeptide which has a functional activity (e.g., biological activity). More preferably, these polynucleotides can be used as probes or primers as discussed herein. Polynucleotides which hybridize to one or more of these polynucleotides under stringent hybridization conditions or alternatively, under lower stringency conditions are also encompassed by the invention, as are polypeptides encoded by these polynucleotides.

[204] Moreover, representative examples of polynucleotide fragments of the invention comprise, or alternatively consist of, a nucleic acid sequence comprising one, two, three, four, five, six, seven, eight, nine, ten, or more of the above described polynucleotide fragments of the invention in combination with a polynucleotide sequence delineated in Table 1C column 6. Additional, representative examples of polynucleotide fragments of the invention comprise, or alternatively consist of, a nucleic acid sequence comprising one, two, three, four, five, six, seven, eight, nine, ten, or more of the above described polynucleotide fragments of the invention in combination with a polynucleotide sequence that is the complementary strand of a sequence delineated in column 6 of Table 1C. In further embodiments, the above-described polynucleotide fragments of the invention comprise, or alternatively consist of, sequences delineated in Table 1C, column 6, and have a nucleic acid sequence which is different from that of the BAC fragment having the sequence disclosed in SEQ ID NO:B (see Table 1C, column 5). In additional embodiments, the above-described polynucleotide fragments of the invention comprise, or alternatively consist of, sequences delineated in Table 1C, column 6, and have a nucleic acid sequence which is different from that published for the BAC clone identified as BAC ID NO:A (see Table 1C, column 4). In additional embodiments, the above-described polynucleotides of the invention comprise, or alternatively consist of, sequences delineated Table 1C, column 6, and have a nucleic acid sequence which is different from that contained in the BAC clone identified as BAC ID NO:A (see Table 1C, column 4). Polypeptides encoded by these polynucleotides, other polynucleotides that encode these polypeptides, and antibodies that bind these polypeptides are also encompassed by the invention. Additionally, fragments and variants of the above-described polynucleotides and polypeptides are also encompassed by the invention.

[205] In additional specific embodiments, polynucleotides of the invention comprise, or alternatively consist of, one, two, three, four, five, six, seven, eight, nine, ten, or more

fragments of the sequences delineated in column 6 of Table 1C, and the polynucleotide sequence of SEQ ID NO:X (e.g., as defined in Table 1C, column 2) or fragments or variants thereof. Polypeptides encoded by these polynucleotides, other polynucleotides that encode these polypeptides, and antibodies that bind these polypeptides are also encompassed by the invention.

[206] In additional specific embodiments, polynucleotides of the invention comprise, or alternatively consist of, one, two, three, four, five, six, seven, eight, nine, ten, or more fragments of the sequences delineated in column 6 of Table 1C which correspond to the same Clone ID NO:Z (see Table 1C, column 1), and the polynucleotide sequence of SEQ ID NO:X (e.g., as defined in Table 1A, 1B, or 1C) or fragments or variants thereof. Polypeptides encoded by these polynucleotides, other polynucleotides that encode these polypeptides, and antibodies that bind these polypeptides are also encompassed by the invention.

[207] In further specific embodiments, polynucleotides of the invention comprise, or alternatively consist of, one, two, three, four, five, six, seven, eight, nine, ten, or more fragments of the sequences delineated in the same row of column 6 of Table 1C, and the polynucleotide sequence of SEQ ID NO:X (e.g., as defined in Table 1A, 1B, or 1C) or fragments or variants thereof. Polypeptides encoded by these polynucleotides, other polynucleotides that encode these polypeptides, and antibodies that bind these polypeptides are also encompassed by the invention.

[208] In additional specific embodiments, polynucleotides of the invention comprise, or alternatively consist of a polynucleotide sequence in which the 3' 10 polynucleotides of one of the sequences delineated in column 6 of Table 1C and the 5' 10 polynucleotides of the sequence of SEQ ID NO:X are directly contiguous. Nucleic acids which hybridize to the complement of these 20 contiguous polynucleotides under stringent hybridization conditions or alternatively, under lower stringency conditions, are also encompassed by the invention. Polypeptides encoded by these polynucleotides and/or nucleic acids, other polynucleotides and/or nucleic acids that encode these polypeptides, and antibodies that bind these polypeptides are also encompassed by the invention. Additionally, fragments and variants of the above-described polynucleotides, nucleic acids, and polypeptides are also encompassed by the invention.

[209] In additional specific embodiments, polynucleotides of the invention comprise, or alternatively consist of a polynucleotide sequence in which the 3' 10 polynucleotides of one

of the sequences delineated in column 6 of Table 1C and the 5' 10 polynucleotides of a fragment or variant of the sequence of SEQ ID NO:X (e.g., as described herein) are directly contiguous Nucleic acids which hybridize to the complement of these 20 contiguous polynucleotides under stringent hybridization conditions or alternatively, under lower stringency conditions, are also encompassed by the invention. Polypeptides encoded by these polynucleotides and/or nucleic acids, other polynucleotides and/or nucleic acids encoding these polypeptides, and antibodies that bind these polypeptides are also encompassed by the invention. Additionally, fragments and variants of the above-described polynucleotides, nucleic acids, and polypeptides are also encompassed by the invention.

[210] In further specific embodiments, polynucleotides of the invention comprise, or alternatively consist of a polynucleotide sequence in which the 3' 10 polynucleotides of a fragment or variant of the sequence of SEQ ID NO:X and the 5' 10 polynucleotides of the sequence of one of the sequences delineated in column 6 of Table 1C are directly contiguous. Nucleic acids which hybridize to the complement of these 20 contiguous polynucleotides under stringent hybridization conditions or alternatively, under lower stringency conditions, are also encompassed by the invention. Polypeptides encoded by these polynucleotides and/or nucleic acids, other polynucleotides and/or nucleic acids encoding these polypeptides, and antibodies that bind these polypeptides are also encompassed by the invention. Additionally, fragments and variants of the above-described polynucleotides, nucleic acids, and polypeptides are also encompassed by the invention.

[211] In specific embodiments, polynucleotides of the invention comprise, or alternatively consist of a polynucleotide sequence in which the 3' 10 polynucleotides of one of the sequences delineated in column 6 of Table 1C and the 5' 10 polynucleotides of another sequence in column 6 are directly contiguous. In preferred embodiments, the 3' 10 polynucleotides of one of the sequences delineated in column 6 of Table 1C is directly contiguous with the 5' 10 polynucleotides of the next sequential exon delineated in Table 1C, column 6. Nucleic acids which hybridize to the complement of these 20 contiguous polynucleotides under stringent hybridization conditions or alternatively, under lower stringency conditions, are also encompassed by the invention. Polypeptides encoded by these polynucleotides and/or nucleic acids, other polynucleotides and/or nucleic acids encoding these polypeptides, and antibodies that bind these polypeptides are also encompassed by the invention. Additionally, fragments and variants of the above-described polynucleotides, nucleic acids, and polypeptides are also encompassed by the invention.

[212] In the present invention, a "polypeptide fragment" refers to an amino acid sequence which is a portion of the amino acid sequence contained in SEQ ID NO:Y, is a portion of the mature form of SEQ ID NO:Y as defined in columns 14 and 15 of Table 1A, a portion of an amino acid sequence encoded by the portion of SEQ ID NO:X as defined in columns 8 and 9 of Table 2, is a portion of an amino acid sequence encoded by the polynucleotide sequence of SEQ ID NO:X, is a portion of an amino acid sequence encoded by the complement of the polynucleotide sequence in SEQ ID NO:X, is a portion of the amino acid sequence of a mature (secreted) polypeptide encoded by the cDNA contained in Clone ID NO:Z, and/or is a portion of an amino acid sequence encoded by the cDNA contained in Clone ID NO:Z. Protein (polypeptide) fragments may be "free-standing," or comprised within a larger polypeptide of which the fragment forms a part or region, most preferably as a single continuous region. Representative examples of polypeptide fragments of the invention, include, for example, fragments comprising, or alternatively consisting of, from about amino acid number 1-20, 21-40, 41-60, 61-80, 81-100, 101-120, 121-140, 141-160, 161-180, 181-200, 201-220, 221-240, 241-260, 261-280, 281-300, 301-320, 321-340, 341-360, 361-380, 381-400, 401-420, 421-440, 441-460, 461-480, 481-500, 501-520, 521-540, 541-560, 561-580, 581-600, 601-620, 621-640, 641-660, 661-680, 681-700, 701-720, 721-740, 741-760, 761-780, 781-800, 801-820, 821-840, 841-860, 861-880, 881-900, 901-920, 921-940, 941-960, 961-980, 981-1000, 1001-1020, 1021-1040, 1041-1060, 1061-1080, 1081-1100, 1101-1120, 1121-1140, 1141-1160, 1161-1180, 1181-1200, 1201-1220, 1221-1240, 1241-1260, 1261-1280, 1281-1300, 1301-1320, 1321-1340, 1341-1360, 1361-1380, 1381-1400, 1401-1420, 1421-1440, or 1441 to the end of the coding region of cDNA and SEQ ID NO: Y. In a preferred embodiment, polypeptide fragments of the invention include, for example, fragments comprising, or alternatively consisting of, from about amino acid number 1-20, 21-40, 41-60, 61-80, 81-100, 101-120, 121-140, 141-160, 161-180, 181-200, 201-220, 221-240, 241-260, 261-280, 281-300, 301-320, 321-340, 341-360, 361-380, 381-400, 401-420, 421-440, 441-460, 461-480, 481-500, 501-520, 521-540, 541-560, 561-580, 581-600, 601-620, 621-640, 641-660, 661-680, 681-700, 701-720, 721-740, 741-760, 761-780, 781-800, 801-820, 821-840, 841-860, 861-880, 881-900, 901-920, 921-940, 941-960, 961-980, 981-1000, 1001-1020, 1021-1040, 1041-1060, 1061-1080, 1081-1100, 1101-1120, 1121-1140, 1141-1160, 1161-1180, 1181-1200, 1201-1220, 1221-1240, 1241-1260, 1261-1280, 1281-1300, 1301-1320, 1321-1340, 1341-1360, 1361-1380, 1381-1400, 1401-1420, 1421-1440, or 1441 to the end of the coding region of SEQ ID

NO:Y. Moreover, polypeptide fragments of the invention may be at least about 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 100, 110, 120, 130, 140, or 150 amino acids in length. In this context "about" includes the particularly recited ranges or values, or ranges or values larger or smaller by several (5, 4, 3, 2, or 1) amino acids, at either extreme or at both extremes. Polynucleotides encoding these polypeptide fragments are also encompassed by the invention.

[213] Even if deletion of one or more amino acids from the N-terminus of a protein results in modification or loss of one or more biological functions of the protein, other functional activities (e.g., biological activities, ability to multimerize, ability to bind a ligand) may still be retained. For example, the ability of shortened muteins to induce and/or bind to antibodies which recognize the complete or mature forms of the polypeptides generally will be retained when less than the majority of the residues of the complete or mature polypeptide are removed from the N-terminus. Whether a particular polypeptide lacking N-terminal residues of a complete polypeptide retains such immunologic activities can readily be determined by routine methods described herein and otherwise known in the art. It is not unlikely that a mutein with a large number of deleted N-terminal amino acid residues may retain some biological or immunogenic activities. In fact, peptides composed of as few as six amino acid residues may often evoke an immune response.

[214] Accordingly, polypeptide fragments include the secreted protein as well as the mature form. Further preferred polypeptide fragments include the secreted protein or the mature form having a continuous series of deleted residues from the amino or the carboxy terminus, or both. For example, any number of amino acids, ranging from 1-60, can be deleted from the amino terminus of either the secreted polypeptide or the mature form. Similarly, any number of amino acids, ranging from 1-30, can be deleted from the carboxy terminus of the secreted protein or mature form. Furthermore, any combination of the above amino and carboxy terminus deletions are preferred. Similarly, polynucleotides encoding these polypeptide fragments are also preferred.

[215] The present invention further provides polypeptides having one or more residues deleted from the amino terminus of the amino acid sequence of a polypeptide disclosed herein (e.g., a polypeptide of SEQ ID NO:Y, a polypeptide as defined in columns 14 and 15 of Table 1A, a polypeptide encoded by the polynucleotide sequence contained in SEQ ID NO:X or the complement thereof, a polypeptide encoded by the portion of SEQ ID NO:X as defined in columns 8 and 9 of Table 2, a polypeptide encoded by the portion of SEQ ID

NO:B as defined in column 6 of Table 1C, a polypeptide encoded by the cDNA contained in Clone ID NO:Z, and/or a mature polypeptide encoded by the cDNA contained in Clone ID NO:Z). In particular, N-terminal deletions may be described by the general formula m-q, where q is a whole integer representing the total number of amino acid residues in a polypeptide of the invention (e.g., the polypeptide disclosed in SEQ ID NO:Y, the mature (secreted) portion of SEQ ID NO:Y as defined in columns 14 and 15 of Table 1A, or the polypeptide encoded by the portion of SEQ ID NO:X as defined in columns 8 and 9 of Table 2), and m is defined as any integer ranging from 2 to q-6. Polynucleotides encoding these polypeptides are also encompassed by the invention.

[216] The present invention further provides polypeptides having one or more residues from the carboxy terminus of the amino acid sequence of a polypeptide disclosed herein (e.g., a polypeptide of SEQ ID NO:Y, the mature (secreted) portion of SEQ ID NO:Y as defined in columns 14 and 15 of Table 1A, a polypeptide encoded by the polynucleotide sequence contained in SEQ ID NO:X, a polypeptide encoded by the portion of SEQ ID NO:X as defined in columns 8 and 9 of Table 2, a polypeptide encoded by the portion of SEQ ID NO:B as defined in column 6 of Table 1C, a polypeptide encoded by the cDNA contained in Clone ID NO:Z, and/or a mature polypeptide encoded by the cDNA contained in Clone ID NO:Z). In particular, C-terminal deletions may be described by the general formula 1-n, where n is any whole integer ranging from 6 to q-1, and where n corresponds to the position of amino acid residue in a polypeptide of the invention. Polynucleotides encoding these polypeptides are also encompassed by the invention.

[217] In addition, any of the above described N- or C-terminal deletions can be combined to produce a N- and C-terminal deleted polypeptide. The invention also provides polypeptides having one or more amino acids deleted from both the amino and the carboxyl termini, which may be described generally as having residues m-n of a polypeptide encoded by SEQ ID NO:X (e.g., including, but not limited to, the preferred polypeptide disclosed as SEQ ID NO:Y, the mature (secreted) portion of SEQ ID NO:Y as defined in columns 14 and 15 of Table 1A, and the polypeptide encoded by the portion of SEQ ID NO:X as defined in columns 8 and 9 of Table 2), the cDNA contained in Clone ID NO:Z, and/or the complement thereof, where n and m are integers as described above. Polynucleotides encoding these polypeptides are also encompassed by the invention.

[218] Also as mentioned above, even if deletion of one or more amino acids from the C-terminus of a protein results in modification of loss of one or more biological functions

of the protein, other functional activities (e.g., biological activities, ability to multimerize, ability to bind a ligand) may still be retained. For example the ability of the shortened mutein to induce and/or bind to antibodies which recognize the complete or mature forms of the polypeptide generally will be retained when less than the majority of the residues of the complete or mature polypeptide are removed from the C-terminus. Whether a particular polypeptide lacking C-terminal residues of a complete polypeptide retains such immunologic activities can readily be determined by routine methods described herein and otherwise known in the art. It is not unlikely that a mutein with a large number of deleted C-terminal amino acid residues may retain some biological or immunogenic activities. In fact, peptides composed of as few as six amino acid residues may often evoke an immune response.

[219] The present application is also directed to proteins containing polypeptides at least 80%, 85%, 90%, 95%, 96%, 97%, 98% or 99% identical to a polypeptide sequence set forth herein. In preferred embodiments, the application is directed to proteins containing polypeptides at least 80%, 85%, 90%, 95%, 96%, 97%, 98% or 99% identical to polypeptides having the amino acid sequence of the specific N- and C-terminal deletions. Polynucleotides encoding these polypeptides are also encompassed by the invention.

[220] Any polypeptide sequence encoded by, for example, the polynucleotide sequences set forth as SEQ ID NO:X or the complement thereof, (presented, for example, in Tables 1A and 2), the cDNA contained in Clone ID NO:Z, or the polynucleotide sequence as defined in column 6 of Table 1C, may be analyzed to determine certain preferred regions of the polypeptide. For example, the amino acid sequence of a polypeptide encoded by a polynucleotide sequence of SEQ ID NO:X (e.g., the polypeptide of SEQ ID NO:Y and the polypeptide encoded by the portion of SEQ ID NO:X as defined in columns 8 and 9 of Table 2) or the cDNA contained in Clone ID NO:Z may be analyzed using the default parameters of the DNASTAR computer algorithm (DNASTAR, Inc., 1228 S. Park St., Madison, WI 53715 USA; <http://www.dnastar.com/>).

[221] Polypeptide regions that may be routinely obtained using the DNASTAR computer algorithm include, but are not limited to, Garnier-Robson alpha-regions, beta-regions, turn-regions, and coil-regions; Chou-Fasman alpha-regions, beta-regions, and turn-regions; Kyte-Doolittle hydrophilic regions and hydrophobic regions; Eisenberg alpha- and beta-amphipathic regions; Karplus-Schulz flexible regions; Emini surface-forming regions; and Jameson-Wolf regions of high antigenic index. Among highly preferred

polynucleotides of the invention in this regard are those that encode polypeptides comprising regions that combine several structural features, such as several (e.g., 1, 2, 3 or 4) of the features set out above.

[222] Additionally, Kyte-Doolittle hydrophilic regions and hydrophobic regions, Emimi surface-forming regions, and Jameson-Wolf regions of high antigenic index (i.e., containing four or more contiguous amino acids having an antigenic index of greater than or equal to 1.5, as identified using the default parameters of the Jameson-Wolf program) can routinely be used to determine polypeptide regions that exhibit a high degree of potential for antigenicity. Regions of high antigenicity are determined from data by DNASTAR analysis by choosing values which represent regions of the polypeptide which are likely to be exposed on the surface of the polypeptide in an environment in which antigen recognition may occur in the process of initiation of an immune response.

[223] Preferred polypeptide fragments of the invention are fragments comprising, or alternatively, consisting of, an amino acid sequence that displays a functional activity (e.g. biological activity) of the polypeptide sequence of which the amino acid sequence is a fragment. By a polypeptide displaying a "functional activity" is meant a polypeptide capable of one or more known functional activities associated with a full-length protein, such as, for example, biological activity, antigenicity, immunogenicity, and/or multimerization, as described herein.

[224] Other preferred polypeptide fragments are biologically active fragments. Biologically active fragments are those exhibiting activity similar, but not necessarily identical, to an activity of the polypeptide of the present invention. The biological activity of the fragments may include an improved desired activity, or a decreased undesirable activity.

[225] In preferred embodiments, polypeptides of the invention comprise, or alternatively consist of, one, two, three, four, five or more of the antigenic fragments of the polypeptide of SEQ ID NO:Y, or portions thereof. Polynucleotides encoding these polypeptides are also encompassed by the invention.

Epitopes and Antibodies

[226] The present invention encompasses polypeptides comprising, or alternatively consisting of, an epitope of: the polypeptide sequence shown in SEQ ID NO:Y; a polypeptide sequence encoded by SEQ ID NO:X or the complementary strand thereto; the

polypeptide sequence encoded by the portion of SEQ ID NO:X as defined in columns 8 and 9 of Table 2; the polypeptide sequence encoded by the portion of SEQ ID NO:B as defined in column 6 of Table 1C or the complement thereto; the polypeptide sequence encoded by the cDNA contained in Clone ID NO:Z; or the polypeptide sequence encoded by a polynucleotide that hybridizes to the sequence of SEQ ID NO:X, the complement of the sequence of SEQ ID NO:X, the complement of a portion of SEQ ID NO:X as defined in columns 8 and 9 of Table 2, or the cDNA sequence contained in Clone ID NO:Z under stringent hybridization conditions or alternatively, under lower stringency hybridization as defined *supra*. The present invention further encompasses polynucleotide sequences encoding an epitope of a polypeptide sequence of the invention (such as, for example, the sequence disclosed in SEQ ID NO:X, or a fragment thereof), polynucleotide sequences of the complementary strand of a polynucleotide sequence encoding an epitope of the invention, and polynucleotide sequences which hybridize to the complementary strand under stringent hybridization conditions or alternatively, under lower stringency hybridization conditions defined *supra*.

[227] The term "epitopes," as used herein, refers to portions of a polypeptide having antigenic or immunogenic activity in an animal, preferably a mammal, and most preferably in a human. In a preferred embodiment, the present invention encompasses a polypeptide comprising an epitope, as well as the polynucleotide encoding this polypeptide. An "immunogenic epitope," as used herein, is defined as a portion of a protein that elicits an antibody response in an animal, as determined by any method known in the art, for example, by the methods for generating antibodies described *infra*. (See, for example, Geysen et al., Proc. Natl. Acad. Sci. USA 81:3998- 4002 (1983)). The term "antigenic epitope," as used herein, is defined as a portion of a protein to which an antibody can immunospecifically bind its antigen as determined by any method well known in the art, for example, by the immunoassays described herein. Immunospecific binding excludes non-specific binding but does not necessarily exclude cross- reactivity with other antigens. Antigenic epitopes need not necessarily be immunogenic.

[228] Fragments which function as epitopes may be produced by any conventional means. (See, e.g., Houghten, R. A., Proc. Natl. Acad. Sci. USA 82:5131-5135 (1985) further described in U.S. Patent No. 4,631,211.)

[229] In the present invention, antigenic epitopes preferably contain a sequence of at least 4, at least 5, at least 6, at least 7, more preferably at least 8, at least 9, at least 10, at

least 11, at least 12, at least 13, at least 14, at least 15, at least 20, at least 25, at least 30, at least 40, at least 50, and, most preferably, between about 15 to about 30 amino acids. Preferred polypeptides comprising immunogenic or antigenic epitopes are at least 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, or 100 amino acid residues in length. Additional non-exclusive preferred antigenic epitopes include the antigenic epitopes disclosed herein, as well as portions thereof. Antigenic epitopes are useful, for example, to raise antibodies, including monoclonal antibodies, that specifically bind the epitope. Preferred antigenic epitopes include the antigenic epitopes disclosed herein, as well as any combination of two, three, four, five or more of these antigenic epitopes. Antigenic epitopes can be used as the target molecules in immunoassays. (See, for instance, Wilson et al., Cell 37:767-778 (1984); Sutcliffe et al., Science 219:660-666 (1983)).

[230] Non-limiting examples of epitopes of polypeptides that can be used to generate antibodies of the invention include a polypeptide comprising, or alternatively consisting of, at least one, two, three, four, five, six or more of the portion(s) of SEQ ID NO:Y specified in column 7 of Table 1B. These polypeptide fragments have been determined to bear antigenic epitopes of the proteins of the invention by the analysis of the Jameson-Wolf antigenic index which is included in the DNASTar suite of computer programs. By "comprise" it is intended that a polypeptide contains at least one, two, three, four, five, six or more of the portion(s) of SEQ ID NO:Y shown in column 7 of Table 1B, but it may contain additional flanking residues on either the amino or carboxyl termini of the recited portion. Such additional flanking sequences are preferably sequences naturally found adjacent to the portion; i.e., contiguous sequence shown in SEQ ID NO:Y. The flanking sequence may, however, be sequences from a heterologous polypeptide, such as from another protein described herein or from a heterologous polypeptide not described herein. In particular embodiments, epitope portions of a polypeptide of the invention comprise one, two, three, or more of the portions of SEQ ID NO:Y shown in column 7 of Table 1B.

[231] Similarly, immunogenic epitopes can be used, for example, to induce antibodies according to methods well known in the art. See, for instance, Sutcliffe et al., *supra*; Wilson et al., *supra*; Chow et al., Proc. Natl. Acad. Sci. USA 82:910-914; and Bittle et al., J. Gen. Virol. 66:2347-2354 (1985). Preferred immunogenic epitopes include the immunogenic epitopes disclosed herein, as well as any combination of two, three, four, five or more of these immunogenic epitopes. The polypeptides comprising one or more immunogenic epitopes may be presented for eliciting an antibody response together with a

carrier protein, such as an albumin, to an animal system (such as rabbit or mouse), or, if the polypeptide is of sufficient length (at least about 25 amino acids), the polypeptide may be presented without a carrier. However, immunogenic epitopes comprising as few as 8 to 10 amino acids have been shown to be sufficient to raise antibodies capable of binding to, at the very least, linear epitopes in a denatured polypeptide (e.g., in Western blotting).

[232] Epitope-bearing polypeptides of the present invention may be used to induce antibodies according to methods well known in the art including, but not limited to, *in vivo* immunization, *in vitro* immunization, and phage display methods. See, e.g., Sutcliffe et al., *supra*; Wilson et al., *supra*, and Bittle et al., J. Gen. Virol., 66:2347-2354 (1985). If *in vivo* immunization is used, animals may be immunized with free peptide; however, anti-peptide antibody titer may be boosted by coupling the peptide to a macromolecular carrier, such as keyhole limpet hemacyanin (KLH) or tetanus toxoid. For instance, peptides containing cysteine residues may be coupled to a carrier using a linker such as maleimidobenzoyl- N-hydroxysuccinimide ester (MBS), while other peptides may be coupled to carriers using a more general linking agent such as glutaraldehyde. Animals such as rabbits, rats and mice are immunized with either free or carrier- coupled peptides, for instance, by intraperitoneal and/or intradermal injection of emulsions containing about 100 μ g of peptide or carrier protein and Freund's adjuvant or any other adjuvant known for stimulating an immune response. Several booster injections may be needed, for instance, at intervals of about two weeks, to provide a useful titer of anti-peptide antibody which can be detected, for example, by ELISA assay using free peptide adsorbed to a solid surface. The titer of anti-peptide antibodies in serum from an immunized animal may be increased by selection of anti-peptide antibodies, for instance, by adsorption to the peptide on a solid support and elution of the selected antibodies according to methods well known in the art.

[233] As one of skill in the art will appreciate, and as discussed above, the polypeptides of the present invention (e.g., those comprising an immunogenic or antigenic epitope) can be fused to heterologous polypeptide sequences. For example, polypeptides of the present invention (including fragments or variants thereof), may be fused with the constant domain of immunoglobulins (IgA, IgE, IgG, IgM), or portions thereof (CH1, CH2, CH3, or any combination thereof and portions thereof, resulting in chimeric polypeptides. By way of another non-limiting example, polypeptides and/or antibodies of the present invention (including fragments or variants thereof) may be fused with albumin (including but not limited to recombinant human serum albumin or fragments or variants thereof (see, e.g.,

U.S. Patent No. 5,876,969, issued March 2, 1999, EP Patent 0 413 622, and U.S. Patent No. 5,766,883, issued June 16, 1998, herein incorporated by reference in their entirety)). In a preferred embodiment, polypeptides and/or antibodies of the present invention (including fragments or variants thereof) are fused with the mature form of human serum albumin (i.e., amino acids 1 – 585 of human serum albumin as shown in Figures 1 and 2 of EP Patent 0 322 094) which is herein incorporated by reference in its entirety. In another preferred embodiment, polypeptides and/or antibodies of the present invention (including fragments or variants thereof) are fused with polypeptide fragments comprising, or alternatively consisting of, amino acid residues 1-z of human serum albumin, where z is an integer from 369 to 419, as described in U.S. Patent 5,766,883 herein incorporated by reference in its entirety. Polypeptides and/or antibodies of the present invention (including fragments or variants thereof) may be fused to either the N- or C-terminal end of the heterologous protein (e.g., immunoglobulin Fc polypeptide or human serum albumin polypeptide). Polynucleotides encoding fusion proteins of the invention are also encompassed by the invention.

[234] Such fusion proteins as those described above may facilitate purification and may increase half-life *in vivo*. This has been shown for chimeric proteins consisting of the first two domains of the human CD4-polypeptide and various domains of the constant regions of the heavy or light chains of mammalian immunoglobulins. See, e.g., EP 394,827; Traunecker et al., *Nature*, 331:84-86 (1988). Enhanced delivery of an antigen across the epithelial barrier to the immune system has been demonstrated for antigens (e.g., insulin) conjugated to an FcRn binding partner such as IgG or Fc fragments (see, e.g., PCT Publications WO 96/22024 and WO 99/04813). IgG fusion proteins that have a disulfide-linked dimeric structure due to the IgG portion disulfide bonds have also been found to be more efficient in binding and neutralizing other molecules than monomeric polypeptides or fragments thereof alone. See, e.g., Fountoulakis et al., *J. Biochem.*, 270:3958-3964 (1995). Nucleic acids encoding the above epitopes can also be recombined with a gene of interest as an epitope tag (e.g., the hemagglutinin (HA) tag or flag tag) to aid in detection and purification of the expressed polypeptide. For example, a system described by Janknecht et al. allows for the ready purification of non-denatured fusion proteins expressed in human cell lines (Janknecht et al., 1991, *Proc. Natl. Acad. Sci. USA* 88:8972-897). In this system, the gene of interest is subcloned into a vaccinia recombination plasmid such that the open reading frame of the gene is translationally fused to an amino-

terminal tag consisting of six histidine residues. The tag serves as a matrix binding domain for the fusion protein. Extracts from cells infected with the recombinant vaccinia virus are loaded onto Ni²⁺ nitriloacetic acid-agarose column and histidine-tagged proteins can be selectively eluted with imidazole-containing buffers.

Fusion Proteins

[235] Any polypeptide of the present invention can be used to generate fusion proteins. For example, the polypeptide of the present invention, when fused to a second protein, can be used as an antigenic tag. Antibodies raised against the polypeptide of the present invention can be used to indirectly detect the second protein by binding to the polypeptide. Moreover, because secreted proteins target cellular locations based on trafficking signals, polypeptides of the present invention which are shown to be secreted can be used as targeting molecules once fused to other proteins.

[236] Examples of domains that can be fused to polypeptides of the present invention include not only heterologous signal sequences, but also other heterologous functional regions. The fusion does not necessarily need to be direct, but may occur through linker sequences.

[237] In certain preferred embodiments, proteins of the invention are fusion proteins comprising an amino acid sequence that is an N and/or C- terminal deletion of a polypeptide of the invention. In preferred embodiments, the invention is directed to a fusion protein comprising an amino acid sequence that is at least 90%, 95%, 96%, 97%, 98% or 99% identical to a polypeptide sequence of the invention. Polynucleotides encoding these proteins are also encompassed by the invention.

[238] Moreover, fusion proteins may also be engineered to improve characteristics of the polypeptide of the present invention. For instance, a region of additional amino acids, particularly charged amino acids, may be added to the N-terminus of the polypeptide to improve stability and persistence during purification from the host cell or subsequent handling and storage. Also, peptide moieties may be added to the polypeptide to facilitate purification. Such regions may be removed prior to final preparation of the polypeptide. The addition of peptide moieties to facilitate handling of polypeptides are familiar and routine techniques in the art.

[239] As one of skill in the art will appreciate that, as discussed above, polypeptides of the present invention, and epitope-bearing fragments thereof, can be combined with

heterologous polypeptide sequences. For example, the polypeptides of the present invention may be fused with heterologous polypeptide sequences, for example, the polypeptides of the present invention may be fused with the constant domain of immunoglobulins (IgA, IgE, IgG, IgM) or portions thereof (CH1, CH2, CH3, and any combination thereof, including both entire domains and portions thereof), or albumin (including, but not limited to, native or recombinant human albumin or fragments or variants thereof (see, e.g., U.S. Patent No. 5,876,969, issued March 2, 1999, EP Patent 0 413 622, and U.S. Patent No. 5,766,883, issued June 16, 1998, herein incorporated by reference in their entirety)), resulting in chimeric polypeptides. For example, EP-A-O 464 533 (Canadian counterpart 2045869) discloses fusion proteins comprising various portions of constant region of immunoglobulin molecules together with another human protein or part thereof. In many cases, the Fc part in a fusion protein is beneficial in therapy and diagnosis, and thus can result in, for example, improved pharmacokinetic properties (EP-A 0232 262). Alternatively, deleting the Fc part after the fusion protein has been expressed, detected, and purified, would be desired. For example, the Fc portion may hinder therapy and diagnosis if the fusion protein is used as an antigen for immunizations. In drug discovery, for example, human proteins, such as hIL-5, have been fused with Fc portions for the purpose of high-throughput screening assays to identify antagonists of hIL-5. See, D. Bennett et al., *J. Molecular Recognition* 8:52-58 (1995); K. Johanson et al., *J. Biol. Chem.* 270:9459-9471 (1995).

[240] Moreover, the polypeptides of the present invention can be fused to marker sequences, such as a polypeptide which facilitates purification of the fused polypeptide. In preferred embodiments, the marker amino acid sequence is a hexa-histidine peptide, such as the tag provided in a pQE vector (QIAGEN, Inc., 9259 Eton Avenue, Chatsworth, CA, 91311), among others, many of which are commercially available. As described in Gentz et al., *Proc. Natl. Acad. Sci. USA* 86:821-824 (1989), for instance, hexa-histidine provides for convenient purification of the fusion protein. Another peptide tag useful for purification, the "HA" tag, corresponds to an epitope derived from the influenza hemagglutinin protein (Wilson et al., *Cell* 37:767 (1984)).

[241] Additional fusion proteins of the invention may be generated through the techniques of gene-shuffling, motif-shuffling, exon-shuffling, and/or codon-shuffling (collectively referred to as "DNA shuffling"). DNA shuffling may be employed to modulate the activities of polypeptides of the invention, such methods can be used to

generate polypeptides with altered activity, as well as agonists and antagonists of the polypeptides. See, generally, U.S. Patent Nos. 5,605,793; 5,811,238; 5,830,721; 5,834,252; and 5,837,458, and Patten et al., Curr. Opinion Biotechnol. 8:724-33 (1997); Harayama, Trends Biotechnol. 16(2):76-82 (1998); Hansson, et al., J. Mol. Biol. 287:265-76 (1999); and Lorenzo and Blasco, Biotechniques 24(2):308- 13 (1998) (each of these patents and publications are hereby incorporated by reference in its entirety). In one embodiment, alteration of polynucleotides corresponding to SEQ ID NO:X and the polypeptides encoded by these polynucleotides may be achieved by DNA shuffling. DNA shuffling involves the assembly of two or more DNA segments by homologous or site-specific recombination to generate variation in the polynucleotide sequence. In another embodiment, polynucleotides of the invention, or the encoded polypeptides, may be altered by being subjected to random mutagenesis by error-prone PCR, random nucleotide insertion or other methods prior to recombination. In another embodiment, one or more components, motifs, sections, parts, domains, fragments, etc., of a polynucleotide encoding a polypeptide of the invention may be recombined with one or more components, motifs, sections, parts, domains, fragments, etc. of one or more heterologous molecules.

[242] Thus, any of these above fusions can be engineered using the polynucleotides or the polypeptides of the present invention.

Recombinant and Synthetic Production of Polypeptides of the Invention

[243] The present invention also relates to vectors containing the polynucleotide of the present invention, host cells, and the production of polypeptides by synthetic and recombinant techniques. The vector may be, for example, a phage, plasmid, viral, or retroviral vector. Retroviral vectors may be replication competent or replication defective. In the latter case, viral propagation generally will occur only in complementing host cells.

[244] The polynucleotides of the invention may be joined to a vector containing a selectable marker for propagation in a host. Generally, a plasmid vector is introduced in a precipitate, such as a calcium phosphate precipitate, or in a complex with a charged lipid. If the vector is a virus, it may be packaged in vitro using an appropriate packaging cell line and then transduced into host cells.

[245] The polynucleotide insert should be operatively linked to an appropriate promoter, such as the phage lambda PL promoter, the E. coli lac, trp, phoA and tac promoters, the SV40 early and late promoters and promoters of retroviral LTRs, to name a

few. Other suitable promoters will be known to the skilled artisan. The expression constructs will further contain sites for transcription initiation, termination, and, in the transcribed region, a ribosome binding site for translation. The coding portion of the transcripts expressed by the constructs will preferably include a translation initiating codon at the beginning and a termination codon (UAA, UGA or UAG) appropriately positioned at the end of the polypeptide to be translated.

[246] As indicated, the expression vectors will preferably include at least one selectable marker. Such markers include dihydrofolate reductase, G418, glutamine synthase, or neomycin resistance for eukaryotic cell culture, and tetracycline, kanamycin or ampicillin resistance genes for culturing in *E. coli* and other bacteria. Representative examples of appropriate hosts include, but are not limited to, bacterial cells, such as *E. coli*, *Streptomyces* and *Salmonella typhimurium* cells; fungal cells, such as yeast cells (e.g., *Saccharomyces cerevisiae* or *Pichia pastoris* (ATCC Accession No. 201178)); insect cells such as *Drosophila* S2 and *Spodoptera Sf9* cells; animal cells such as CHO, COS, 293, and Bowes melanoma cells; and plant cells. Appropriate culture mediums and conditions for the above-described host cells are known in the art.

[247] Among vectors preferred for use in bacteria include pQE70, pQE60 and pQE-9, available from QIAGEN, Inc.; pBluescript vectors, Phagescript vectors, pNH8A, pNH16a, pNH18A, pNH46A, available from Stratagene Cloning Systems, Inc.; and ptrc99a, pKK223-3, pKK233-3, pDR540, pRIT5 available from Pharmacia Biotech, Inc. Among preferred eukaryotic vectors are pWLNEO, pSV2CAT, pOG44, pXT1 and pSG available from Stratagene; and pSVK3, pBPV, pMSG and pSVL available from Pharmacia. Preferred expression vectors for use in yeast systems include, but are not limited to pYES2, pYD1, pTEF1/Zeo, pYES2/GS, pPICZ, pGAPZ, pGAPZalph, pPIC9, pPIC3.5, pHIL-D2, pHIL-S1, pPIC3.5K, pPIC9K, and PAO815 (all available from Invitrogen, Carlsbad, CA). Other suitable vectors will be readily apparent to the skilled artisan.

[248] Vectors which use glutamine synthase (GS) or DHFR as the selectable markers can be amplified in the presence of the drugs methionine sulfoximine or methotrexate, respectively. An advantage of glutamine synthase based vectors are the availability of cell lines (e.g., the murine myeloma cell line, NS0) which are glutamine synthase negative. Glutamine synthase expression systems can also function in glutamine synthase expressing cells (e.g., Chinese Hamster Ovary (CHO) cells) by providing additional inhibitor to prevent the functioning of the endogenous gene. A glutamine synthase expression system

and components thereof are detailed in PCT publications: WO87/04462; WO86/05807; WO89/01036; WO89/10404; and WO91/06657, which are hereby incorporated in their entireties by reference herein. Additionally, glutamine synthase expression vectors can be obtained from Lonza Biologics, Inc. (Portsmouth, NH). Expression and production of monoclonal antibodies using a GS expression system in murine myeloma cells is described in Bebbington *et al.*, *Bio/technology* 10:169(1992) and in Biblia and Robinson *Biotechnol. Prog.* 11:1 (1995) which are herein incorporated by reference.

[249] The present invention also relates to host cells containing the above-described vector constructs described herein, and additionally encompasses host cells containing nucleotide sequences of the invention that are operably associated with one or more heterologous control regions (e.g., promoter and/or enhancer) using techniques known of in the art. The host cell can be a higher eukaryotic cell, such as a mammalian cell (e.g., a human derived cell), or a lower eukaryotic cell, such as a yeast cell, or the host cell can be a prokaryotic cell, such as a bacterial cell. A host strain may be chosen which modulates the expression of the inserted gene sequences, or modifies and processes the gene product in the specific fashion desired. Expression from certain promoters can be elevated in the presence of certain inducers; thus expression of the genetically engineered polypeptide may be controlled. Furthermore, different host cells have characteristics and specific mechanisms for the translational and post-translational processing and modification (e.g., phosphorylation, cleavage) of proteins. Appropriate cell lines can be chosen to ensure the desired modifications and processing of the foreign protein expressed.

[250] Introduction of the nucleic acids and nucleic acid constructs of the invention into the host cell can be effected by calcium phosphate transfection, DEAE-dextran mediated transfection, cationic lipid-mediated transfection, electroporation, transduction, infection, or other methods. Such methods are described in many standard laboratory manuals, such as Davis *et al.*, *Basic Methods In Molecular Biology* (1986). It is specifically contemplated that the polypeptides of the present invention may in fact be expressed by a host cell lacking a recombinant vector.

[251] In addition to encompassing host cells containing the vector constructs discussed herein, the invention also encompasses primary, secondary, and immortalized host cells of vertebrate origin, particularly mammalian origin, that have been engineered to delete or replace endogenous genetic material (e.g., the coding sequence), and/or to include genetic material (e.g., heterologous polynucleotide sequences) that is operably associated with

polynucleotides of the invention, and which activates, alters, and/or amplifies endogenous polynucleotides. For example, techniques known in the art may be used to operably associate heterologous control regions (e.g., promoter and/or enhancer) and endogenous polynucleotide sequences via homologous recombination (see, e.g., US Patent Number 5,641,670, issued June 24, 1997; International Publication Number WO 96/29411; International Publication Number WO 94/12650; Koller *et al.*, *Proc. Natl. Acad. Sci. USA* 86:8932-8935 (1989); and Zijlstra *et al.*, *Nature* 342:435-438 (1989), the disclosures of each of which are incorporated by reference in their entireties).

[252] Polypeptides of the invention can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Most preferably, high performance liquid chromatography ("HPLC") is employed for purification.

[253] Polypeptides of the present invention can also be recovered from: products purified from natural sources, including bodily fluids, tissues and cells, whether directly isolated or cultured; products of chemical synthetic procedures; and products produced by recombinant techniques from a prokaryotic or eukaryotic host, including, for example, bacterial, yeast, higher plant, insect, and mammalian cells. Depending upon the host employed in a recombinant production procedure, the polypeptides of the present invention may be glycosylated or may be non-glycosylated. In addition, polypeptides of the invention may also include an initial modified methionine residue, in some cases as a result of host-mediated processes. Thus, it is well known in the art that the N-terminal methionine encoded by the translation initiation codon generally is removed with high efficiency from any protein after translation in all eukaryotic cells. While the N-terminal methionine on most proteins also is efficiently removed in most prokaryotes, for some proteins, this prokaryotic removal process is inefficient, depending on the nature of the amino acid to which the N-terminal methionine is covalently linked.

[254] In one embodiment, the yeast *Pichia pastoris* is used to express polypeptides of the invention in a eukaryotic system. *Pichia pastoris* is a methylotrophic yeast which can metabolize methanol as its sole carbon source. A main step in the methanol metabolism pathway is the oxidation of methanol to formaldehyde using O₂. This reaction is catalyzed by the enzyme alcohol oxidase. In order to metabolize methanol as its sole carbon source,

Pichia pastoris must generate high levels of alcohol oxidase due, in part, to the relatively low affinity of alcohol oxidase for O₂. Consequently, in a growth medium depending on methanol as a main carbon source, the promoter region of one of the two alcohol oxidase genes (*AOX1*) is highly active. In the presence of methanol, alcohol oxidase produced from the *AOX1* gene comprises up to approximately 30% of the total soluble protein in *Pichia pastoris*. See Ellis, S.B., *et al.*, *Mol. Cell. Biol.* 5:1111-21 (1985); Koutz, P.J., *et al.*, *Yeast* 5:167-77 (1989); Tschopp, J.F., *et al.*, *Nucl. Acids Res.* 15:3859-76 (1987). Thus, a heterologous coding sequence, such as, for example, a polynucleotide of the present invention, under the transcriptional regulation of all or part of the *AOX1* regulatory sequence is expressed at exceptionally high levels in *Pichia* yeast grown in the presence of methanol.

[255] In one example, the plasmid vector pPIC9K is used to express DNA encoding a polypeptide of the invention, as set forth herein, in a *Pichea* yeast system essentially as described in "*Pichia* Protocols: Methods in Molecular Biology," D.R. Higgins and J. Cregg, eds. The Humana Press, Totowa, NJ, 1998. This expression vector allows expression and secretion of a polypeptide of the invention by virtue of the strong *AOX1* promoter linked to the *Pichia pastoris* alkaline phosphatase (PHO) secretory signal peptide (i.e., leader) located upstream of a multiple cloning site.

[256] Many other yeast vectors could be used in place of pPIC9K, such as, pYES2, pYD1, pTEF1/Zeo, pYES2/GS, pPICZ, pGAPZ, pGAPZalpha, pPIC9, pPIC3.5, pHIL-D2, pHIL-S1, pPIC3.5K, and PAO815, as one skilled in the art would readily appreciate, as long as the proposed expression construct provides appropriately located signals for transcription, translation, secretion (if desired), and the like, including an in-frame AUG as required.

[257] In another embodiment, high-level expression of a heterologous coding sequence, such as, for example, a polynucleotide of the present invention, may be achieved by cloning the heterologous polynucleotide of the invention into an expression vector such as, for example, pGAPZ or pGAPZalpha, and growing the yeast culture in the absence of methanol.

[258] In addition to encompassing host cells containing the vector constructs discussed herein, the invention also encompasses primary, secondary, and immortalized host cells of

vertebrate origin, particularly mammalian origin, that have been engineered to delete or replace endogenous genetic material (e.g., coding sequence), and/or to include genetic material (e.g., heterologous polynucleotide sequences) that is operably associated with polynucleotides of the invention, and which activates, alters, and/or amplifies endogenous polynucleotides. For example, techniques known in the art may be used to operably associate heterologous control regions (e.g., promoter and/or enhancer) and endogenous polynucleotide sequences via homologous recombination (see, e.g., U.S. Patent No. 5,641,670, issued June 24, 1997; International Publication No. WO 96/29411, published September 26, 1996; International Publication No. WO 94/12650, published August 4, 1994; Koller et al., *Proc. Natl. Acad. Sci. USA* 86:8932-8935 (1989); and Zijlstra et al., *Nature* 342:435-438 (1989), the disclosures of each of which are incorporated by reference in their entireties).

[259] In addition, polypeptides of the invention can be chemically synthesized using techniques known in the art (e.g., see Creighton, 1983, *Proteins: Structures and Molecular Principles*, W.H. Freeman & Co., N.Y., and Hunkapiller et al., *Nature*, 310:105-111 (1984)). For example, a polypeptide corresponding to a fragment of a polypeptide can be synthesized by use of a peptide synthesizer. Furthermore, if desired, nonclassical amino acids or chemical amino acid analogs can be introduced as a substitution or addition into the polypeptide sequence. Non-classical amino acids include, but are not limited to, to the D-isomers of the common amino acids, 2,4-diaminobutyric acid, α -amino isobutyric acid, 4-aminobutyric acid, Abu, 2-amino butyric acid, g-Abu, e-Ahx, 6-amino hexanoic acid, Aib, 2-amino isobutyric acid, 3-amino propionic acid, ornithine, norleucine, norvaline, hydroxyproline, sarcosine, citrulline, homocitrulline, cysteic acid, t-butylglycine, t-butylalanine, phenylglycine, cyclohexylalanine, b-alanine, fluoro-amino acids, designer amino acids such as b-methyl amino acids, Ca-methyl amino acids, Na-methyl amino acids, and amino acid analogs in general. Furthermore, the amino acid can be D (dextrorotary) or L (levorotary).

[260] The invention encompasses polypeptides of the present invention which are differentially modified during or after translation, e.g., by glycosylation, acetylation, phosphorylation, amidation, derivatization by known protecting/blocking groups, proteolytic cleavage, linkage to an antibody molecule or other cellular ligand, etc. Any of numerous chemical modifications may be carried out by known techniques, including but not limited, to specific chemical cleavage by cyanogen bromide, trypsin, chymotrypsin,

papain, V8 protease, NaBH₄; acetylation, formylation, oxidation, reduction; metabolic synthesis in the presence of tunicamycin; etc.

[261] Additional post-translational modifications encompassed by the invention include, for example, e.g., N-linked or O-linked carbohydrate chains, processing of N-terminal or C-terminal ends), attachment of chemical moieties to the amino acid backbone, chemical modifications of N-linked or O-linked carbohydrate chains, and addition or deletion of an N-terminal methionine residue as a result of procaryotic host cell expression. The polypeptides may also be modified with a detectable label, such as an enzymatic, fluorescent, isotopic or affinity label to allow for detection and isolation of the protein.

[262] Examples of suitable enzymes include horseradish peroxidase, alkaline phosphatase, beta-galactosidase, or acetylcholinesterase; examples of suitable prosthetic group complexes include streptavidin/biotin and avidin/biotin; examples of suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; an example of a luminescent material includes luminol; examples of bioluminescent materials include luciferase, luciferin, and aequorin; and examples of suitable radioactive material include iodine (¹²¹I, ¹²³I, ¹²⁵I, ¹³¹I), carbon (¹⁴C), sulfur (³⁵S), tritium (³H), indium (¹¹¹In, ¹¹²In, ^{113m}In, ^{115m}In), technetium (⁹⁹Tc, ^{99m}Tc), thallium (²⁰¹Tl), gallium (⁶⁸Ga, ⁶⁷Ga), palladium (¹⁰³Pd), molybdenum (⁹⁹Mo), xenon (¹³³Xe), fluorine (¹⁸F), ¹⁵³Sm, ¹⁷⁷Lu, ¹⁵⁹Gd, ¹⁴⁹Pm, ¹⁴⁰La, ¹⁷⁵Yb, ¹⁶⁶Ho, ⁹⁰Y, ⁴⁷Sc, ¹⁸⁶Re, ¹⁸⁸Re, ¹⁴²Pr, ¹⁰⁵Rh, and ⁹⁷Ru.

[263] In specific embodiments, a polypeptide of the present invention or fragment or variant thereof is attached to macrocyclic chelators that associate with radiometal ions, including but not limited to, ¹⁷⁷Lu, ⁹⁰Y, ¹⁶⁶Ho, and ¹⁵³Sm, to polypeptides. In a preferred embodiment, the radiometal ion associated with the macrocyclic chelators is ¹¹¹In. In another preferred embodiment, the radiometal ion associated with the macrocyclic chelator is ⁹⁰Y. In specific embodiments, the macrocyclic chelator is 1,4,7,10-tetraazacyclododecane-N,N',N'',N'''-tetraacetic acid (DOTA). In other specific embodiments, DOTA is attached to an antibody of the invention or fragment thereof via a linker molecule. Examples of linker molecules useful for conjugating DOTA to a polypeptide are commonly known in the art - see, for example, DeNardo et al., Clin Cancer Res. 4(10):2483-90 (1998); Peterson et al., Bioconjug. Chem. 10(4):553-7 (1999); and

Zimmerman et al, Nucl. Med. Biol. 26(8):943-50 (1999); which are hereby incorporated by reference in their entirety.

[264] As mentioned, the proteins of the invention may be modified by either natural processes, such as posttranslational processing, or by chemical modification techniques which are well known in the art. It will be appreciated that the same type of modification may be present in the same or varying degrees at several sites in a given polypeptide. Polypeptides of the invention may be branched, for example, as a result of ubiquitination, and they may be cyclic, with or without branching. Cyclic, branched, and branched cyclic polypeptides may result from posttranslation natural processes or may be made by synthetic methods. Modifications include acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent cross-links, formation of cysteine, formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, pegylation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins such as arginylation, and ubiquitination. (See, for instance, PROTEINS - STRUCTURE AND MOLECULAR PROPERTIES, 2nd Ed., T. E. Creighton, W. H. Freeman and Company, New York (1993); POSTTRANSLATIONAL COVALENT MODIFICATION OF PROTEINS, B. C. Johnson, Ed., Academic Press, New York, pgs. 1-12 (1983); Seifter et al., Meth. Enzymol. 182:626-646 (1990); Rattan et al., Ann. N.Y. Acad. Sci. 663:48-62 (1992)).

[265] Also provided by the invention are chemically modified derivatives of the polypeptides of the invention which may provide additional advantages such as increased solubility, stability and circulating time of the polypeptide, or decreased immunogenicity (see U.S. Patent No. 4,179,337). The chemical moieties for derivitization may be selected from water soluble polymers such as polyethylene glycol, ethylene glycol/propylene glycol copolymers, carboxymethylcellulose, dextran, polyvinyl alcohol and the like. The polypeptides may be modified at random positions within the molecule, or at predetermined positions within the molecule and may include one, two, three or more attached chemical moieties.

[269] As suggested above, polyethylene glycol may be attached to proteins via linkage to any of a number of amino acid residues. For example, polyethylene glycol can be linked to proteins via covalent bonds to lysine, histidine, aspartic acid, glutamic acid, or cysteine residues. One or more reaction chemistries may be employed to attach polyethylene glycol to specific amino acid residues (e.g., lysine, histidine, aspartic acid, glutamic acid, or cysteine) of the protein or to more than one type of amino acid residue (e.g., lysine, histidine, aspartic acid, glutamic acid, cysteine and combinations thereof) of the protein.

[270] One may specifically desire proteins chemically modified at the N-terminus. Using polyethylene glycol as an illustration of the present composition, one may select from a variety of polyethylene glycol molecules (by molecular weight, branching, etc.), the proportion of polyethylene glycol molecules to protein (polypeptide) molecules in the reaction mix, the type of pegylation reaction to be performed, and the method of obtaining the selected N-terminally pegylated protein. The method of obtaining the N-terminally pegylated preparation (i.e., separating this moiety from other monopegylated moieties if necessary) may be by purification of the N-terminally pegylated material from a population of pegylated protein molecules. Selective proteins chemically modified at the N-terminus modification may be accomplished by reductive alkylation which exploits differential reactivity of different types of primary amino groups (lysine versus the N-terminal) available for derivatization in a particular protein. Under the appropriate reaction conditions, substantially selective derivatization of the protein at the N-terminus with a carbonyl group containing polymer is achieved.

[271] As indicated above, pegylation of the proteins of the invention may be accomplished by any number of means. For example, polyethylene glycol may be attached to the protein either directly or by an intervening linker. Linkerless systems for attaching polyethylene glycol to proteins are described in Delgado et al., Crit. Rev. Thera. Drug Carrier Sys. 9:249-304 (1992); Francis et al., Intern. J. of Hematol. 68:1-18 (1998); U.S. Patent No. 4,002,531; U.S. Patent No. 5,349,052; WO 95/06058; and WO 98/32466, the disclosures of each of which are incorporated herein by reference.

[272] One system for attaching polyethylene glycol directly to amino acid residues of proteins without an intervening linker employs tresylated MPEG, which is produced by the modification of monmethoxy polyethylene glycol (MPEG) using tresylchloride ($\text{ClSO}_2\text{CH}_2\text{CF}_3$). Upon reaction of protein with tresylated MPEG, polyethylene glycol is directly attached to amine groups of the protein. Thus, the invention includes protein-

polyethylene glycol conjugates produced by reacting proteins of the invention with a polyethylene glycol molecule having a 2,2,2-trifluoroethane sulphonyl group.

[273] Polyethylene glycol can also be attached to proteins using a number of different intervening linkers. For example, U.S. Patent No. 5,612,460, the entire disclosure of which is incorporated herein by reference, discloses urethane linkers for connecting polyethylene glycol to proteins. Protein-polyethylene glycol conjugates wherein the polyethylene glycol is attached to the protein by a linker can also be produced by reaction of proteins with compounds such as MPEG-succinimidylsuccinate, MPEG activated with 1,1'-carbonyldiimidazole, MPEG-2,4,5-trichloropenylcarbonate, MPEG-p-nitrophenolcarbonate, and various MPEG-succinate derivatives. A number of additional polyethylene glycol derivatives and reaction chemistries for attaching polyethylene glycol to proteins are described in International Publication No. WO 98/32466, the entire disclosure of which is incorporated herein by reference. Pegylated protein products produced using the reaction chemistries set out herein are included within the scope of the invention.

[274] The number of polyethylene glycol moieties attached to each protein of the invention (i.e., the degree of substitution) may also vary. For example, the pegylated proteins of the invention may be linked, on average, to 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 12, 15, 17, 20, or more polyethylene glycol molecules. Similarly, the average degree of substitution within ranges such as 1-3, 2-4, 3-5, 4-6, 5-7, 6-8, 7-9, 8-10, 9-11, 10-12, 11-13, 12-14, 13-15, 14-16, 15-17, 16-18, 17-19, or 18-20 polyethylene glycol moieties per protein molecule. Methods for determining the degree of substitution are discussed, for example, in Delgado et al., Crit. Rev. Thera. Drug Carrier Sys. 9:249-304 (1992).

[275] The polypeptides of the invention can be recovered and purified from chemical synthesis and recombinant cell cultures by standard methods which include, but are not limited to, ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Most preferably, high performance liquid chromatography ("HPLC") is employed for purification. Well known techniques for refolding protein may be employed to regenerate active conformation when the polypeptide is denatured during isolation and/or purification.

[276] The polypeptides of the invention may be in monomers or multimers (i.e., dimers, trimers, tetramers and higher multimers). Accordingly, the present invention relates to monomers and multimers of the polypeptides of the invention, their preparation, and compositions (preferably, Therapeutics) containing them. In specific embodiments, the polypeptides of the invention are monomers, dimers, trimers or tetramers. In additional embodiments, the multimers of the invention are at least dimers, at least trimers, or at least tetramers.

[277] Multimers encompassed by the invention may be homomers or heteromers. As used herein, the term homomer refers to a multimer containing only polypeptides corresponding to a protein of the invention (e.g., the amino acid sequence of SEQ ID NO:Y, an amino acid sequence encoded by SEQ ID NO:X or the complement of SEQ ID NO:X, the amino acid sequence encoded by the portion of SEQ ID NO:X as defined in columns 8 and 9 of Table 2, and/or an amino acid sequence encoded by cDNA contained in Clone ID NO:Z (including fragments, variants, splice variants, and fusion proteins, corresponding to these as described herein)). These homomers may contain polypeptides having identical or different amino acid sequences. In a specific embodiment, a homomer of the invention is a multimer containing only polypeptides having an identical amino acid sequence. In another specific embodiment, a homomer of the invention is a multimer containing polypeptides having different amino acid sequences. In specific embodiments, the multimer of the invention is a homodimer (e.g., containing two polypeptides having identical or different amino acid sequences) or a homotrimer (e.g., containing three polypeptides having identical and/or different amino acid sequences). In additional embodiments, the homomeric multimer of the invention is at least a homodimer, at least a homotrimer, or at least a homotetramer.

[278] As used herein, the term heteromer refers to a multimer containing one or more heterologous polypeptides (i.e., polypeptides of different proteins) in addition to the polypeptides of the invention. In a specific embodiment, the multimer of the invention is a heterodimer, a heterotrimer, or a heterotetramer. In additional embodiments, the heteromeric multimer of the invention is at least a heterodimer, at least a heterotrimer, or at least a heterotetramer.

[279] Multimers of the invention may be the result of hydrophobic, hydrophilic, ionic and/or covalent associations and/or may be indirectly linked by, for example, liposome formation. Thus, in one embodiment, multimers of the invention, such as, for example,

homodimers or homotrimers, are formed when polypeptides of the invention contact one another in solution. In another embodiment, heteromultimers of the invention, such as, for example, heterotrimers or heterotetramers, are formed when polypeptides of the invention contact antibodies to the polypeptides of the invention (including antibodies to the heterologous polypeptide sequence in a fusion protein of the invention) in solution. In other embodiments, multimers of the invention are formed by covalent associations with and/or between the polypeptides of the invention. Such covalent associations may involve one or more amino acid residues contained in the polypeptide sequence (e.g., that recited in SEQ ID NO:Y, encoded by the portion of SEQ ID NO:X as defined in columns 8 and 9 of Table 2, and/or encoded by the cDNA contained in Clone ID NO:Z). In one instance, the covalent associations are cross-linking between cysteine residues located within the polypeptide sequences which interact in the native (i.e., naturally occurring) polypeptide. In another instance, the covalent associations are the consequence of chemical or recombinant manipulation. Alternatively, such covalent associations may involve one or more amino acid residues contained in the heterologous polypeptide sequence in a fusion protein. In one example, covalent associations are between the heterologous sequence contained in a fusion protein of the invention (see, e.g., US Patent Number 5,478,925). In a specific example, the covalent associations are between the heterologous sequence contained in a Fc fusion protein of the invention (as described herein). In another specific example, covalent associations of fusion proteins of the invention are between heterologous polypeptide sequence from another protein that is capable of forming covalently associated multimers, such as for example, osteoprotegerin (see, e.g., International Publication NO: WO 98/49305, the contents of which are herein incorporated by reference in its entirety). In another embodiment, two or more polypeptides of the invention are joined through peptide linkers. Examples include those peptide linkers described in U.S. Pat. No. 5,073,627 (hereby incorporated by reference). Proteins comprising multiple polypeptides of the invention separated by peptide linkers may be produced using conventional recombinant DNA technology.

[280] Another method for preparing multimer polypeptides of the invention involves use of polypeptides of the invention fused to a leucine zipper or isoleucine zipper polypeptide sequence. Leucine zipper and isoleucine zipper domains are polypeptides that promote multimerization of the proteins in which they are found. Leucine zippers were originally identified in several DNA-binding proteins (Landschulz et al., Science 240:1759,

(1988)), and have since been found in a variety of different proteins. Among the known leucine zippers are naturally occurring peptides and derivatives thereof that dimerize or trimerize. Examples of leucine zipper domains suitable for producing soluble multimeric proteins of the invention are those described in PCT application WO 94/10308, hereby incorporated by reference. Recombinant fusion proteins comprising a polypeptide of the invention fused to a polypeptide sequence that dimerizes or trimerizes in solution are expressed in suitable host cells, and the resulting soluble multimeric fusion protein is recovered from the culture supernatant using techniques known in the art.

[281] Trimeric polypeptides of the invention may offer the advantage of enhanced biological activity. Preferred leucine zipper moieties and isoleucine moieties are those that preferentially form trimers. One example is a leucine zipper derived from lung surfactant protein D (SPD), as described in Hoppe et al. (FEBS Letters 344:191, (1994)) and in U.S. patent application Ser. No. 08/446,922, hereby incorporated by reference. Other peptides derived from naturally occurring trimeric proteins may be employed in preparing trimeric polypeptides of the invention.

[282] In another example, proteins of the invention are associated by interactions between Flag® polypeptide sequence contained in fusion proteins of the invention containing Flag® polypeptide sequence. In a further embodiment, proteins of the invention are associated by interactions between heterologous polypeptide sequence contained in Flag® fusion proteins of the invention and anti-Flag® antibody.

[283] The multimers of the invention may be generated using chemical techniques known in the art. For example, polypeptides desired to be contained in the multimers of the invention may be chemically cross-linked using linker molecules and linker molecule length optimization techniques known in the art (see, e.g., US Patent Number 5,478,925, which is herein incorporated by reference in its entirety). Additionally, multimers of the invention may be generated using techniques known in the art to form one or more inter-molecule cross-links between the cysteine residues located within the sequence of the polypeptides desired to be contained in the multimer (see, e.g., US Patent Number 5,478,925, which is herein incorporated by reference in its entirety). Further, polypeptides of the invention may be routinely modified by the addition of cysteine or biotin to the C-terminus or N-terminus of the polypeptide and techniques known in the art may be applied to generate multimers containing one or more of these modified polypeptides (see, e.g., US Patent Number 5,478,925, which is herein incorporated by reference in its entirety).

09950063-091201

Additionally, techniques known in the art may be applied to generate liposomes containing the polypeptide components desired to be contained in the multimer of the invention (see, e.g., US Patent Number 5,478,925, which is herein incorporated by reference in its entirety).

[284] Alternatively, multimers of the invention may be generated using genetic engineering techniques known in the art. In one embodiment, polypeptides contained in multimers of the invention are produced recombinantly using fusion protein technology described herein or otherwise known in the art (see, e.g., US Patent Number 5,478,925, which is herein incorporated by reference in its entirety). In a specific embodiment, polynucleotides coding for a homodimer of the invention are generated by ligating a polynucleotide sequence encoding a polypeptide of the invention to a sequence encoding a linker polypeptide and then further to a synthetic polynucleotide encoding the translated product of the polypeptide in the reverse orientation from the original C-terminus to the N-terminus (lacking the leader sequence) (see, e.g., US Patent Number 5,478,925, which is herein incorporated by reference in its entirety). In another embodiment, recombinant techniques described herein or otherwise known in the art are applied to generate recombinant polypeptides of the invention which contain a transmembrane domain (or hydrophobic or signal peptide) and which can be incorporated by membrane reconstitution techniques into liposomes (see, e.g., US Patent Number 5,478,925, which is herein incorporated by reference in its entirety).

Antibodies

[285] Further polypeptides of the invention relate to antibodies and T-cell antigen receptors (TCR) which immunospecifically bind a polypeptide, polypeptide fragment, or variant of the invention (e.g., a polypeptide or fragment or variant of the amino acid sequence of SEQ ID NO:Y or a polypeptide encoded by the cDNA contained in Clone ID No:Z, and/or an epitope, of the present invention) as determined by immunoassays well known in the art for assaying specific antibody-antigen binding. Antibodies of the invention include, but are not limited to, polyclonal, monoclonal, multispecific, human, humanized or chimeric antibodies, single chain antibodies, Fab fragments, F(ab') fragments, fragments produced by a Fab expression library, anti-idiotypic (anti-Id) antibodies (including, e.g., anti-Id antibodies to antibodies of the invention), intracellularly-made antibodies (i.e., intrabodies), and epitope-binding fragments of any of the above. The

term "antibody," as used herein, refers to immunoglobulin molecules and immunologically active portions of immunoglobulin molecules, i.e., molecules that contain an antigen binding site that immunospecifically binds an antigen. The immunoglobulin molecules of the invention can be of any type (e.g., IgG, IgE, IgM, IgD, IgA and IgY), class (e.g., IgG1, IgG2, IgG3, IgG4, IgA1 and IgA2) or subclass of immunoglobulin molecule. In preferred embodiments, the immunoglobulin molecules of the invention are IgG1. In other preferred embodiments, the immunoglobulin molecules of the invention are IgG4.

[286] Most preferably the antibodies are human antigen-binding antibody fragments of the present invention and include, but are not limited to, Fab, Fab' and F(ab')₂, Fd, single-chain Fvs (scFv), single-chain antibodies, disulfide-linked Fvs (sdFv) and fragments comprising either a VL or VH domain. Antigen-binding antibody fragments, including single-chain antibodies, may comprise the variable region(s) alone or in combination with the entirety or a portion of the following: hinge region, CH1, CH2, and CH3 domains. Also included in the invention are antigen-binding fragments also comprising any combination of variable region(s) with a hinge region, CH1, CH2, and CH3 domains. The antibodies of the invention may be from any animal origin including birds and mammals. Preferably, the antibodies are human, murine (e.g., mouse and rat), donkey, sheep rabbit, goat, guinea pig, camel, horse, or chicken. As used herein, "human" antibodies include antibodies having the amino acid sequence of a human immunoglobulin and include antibodies isolated from human immunoglobulin libraries or from animals transgenic for one or more human immunoglobulin and that do not express endogenous immunoglobulins, as described infra and, for example in, U.S. Patent No. 5,939,598 by Kucherlapati et al.

[287] The antibodies of the present invention may be monospecific, bispecific, trispecific or of greater multispecificity. Multispecific antibodies may be specific for different epitopes of a polypeptide of the present invention or may be specific for both a polypeptide of the present invention as well as for a heterologous epitope, such as a heterologous polypeptide or solid support material. See, e.g., PCT publications WO 93/17715; WO 92/08802; WO 91/00360; WO 92/05793; Tutt, et al., J. Immunol. 147:60-69 (1991); U.S. Patent Nos. 4,474,893; 4,714,681; 4,925,648; 5,573,920; 5,601,819; Kostelny et al., J. Immunol. 148:1547-1553 (1992).

[288] Antibodies of the present invention may be described or specified in terms of the epitope(s) or portion(s) of a polypeptide of the present invention which they recognize or specifically bind. The epitope(s) or polypeptide portion(s) may be specified as described

herein, e.g., by N-terminal and C-terminal positions, or by size in contiguous amino acid residues, or listed in the Tables and Figures. Preferred epitopes of the invention include the predicted epitopes shown in column 7 of Table 1B, as well as polynucleotides that encode these epitopes. Antibodies which specifically bind any epitope or polypeptide of the present invention may also be excluded. Therefore, the present invention includes antibodies that specifically bind polypeptides of the present invention, and allows for the exclusion of the same.

[289] Antibodies of the present invention may also be described or specified in terms of their cross-reactivity. Antibodies that do not bind any other analog, ortholog, or homolog of a polypeptide of the present invention are included. Antibodies that bind polypeptides with at least 95%, at least 90%, at least 85%, at least 80%, at least 75%, at least 70%, at least 65%, at least 60%, at least 55%, and at least 50% identity (as calculated using methods known in the art and described herein) to a polypeptide of the present invention are also included in the present invention. In specific embodiments, antibodies of the present invention cross-react with murine, rat and/or rabbit homologs of human proteins and the corresponding epitopes thereof. Antibodies that do not bind polypeptides with less than 95%, less than 90%, less than 85%, less than 80%, less than 75%, less than 70%, less than 65%, less than 60%, less than 55%, and less than 50% identity (as calculated using methods known in the art and described herein) to a polypeptide of the present invention are also included in the present invention. In a specific embodiment, the above-described cross-reactivity is with respect to any single specific antigenic or immunogenic polypeptide, or combination(s) of 2, 3, 4, 5, or more of the specific antigenic and/or immunogenic polypeptides disclosed herein. Further included in the present invention are antibodies which bind polypeptides encoded by polynucleotides which hybridize to a polynucleotide of the present invention under stringent hybridization conditions (as described herein). Antibodies of the present invention may also be described or specified in terms of their binding affinity to a polypeptide of the invention. Preferred binding affinities include those with a dissociation constant or K_d less than 5×10^{-2} M, 10^{-2} M, 5×10^{-3} M, 10^{-3} M, 5×10^{-4} M, 10^{-4} M, 5×10^{-5} M, 10^{-5} M, 5×10^{-6} M, 10^{-6} M, 5×10^{-7} M, 10^{-7} M, 5×10^{-8} M, 10^{-8} M, 5×10^{-9} M, 10^{-9} M, 5×10^{-10} M, 10^{-10} M, 5×10^{-11} M, 10^{-11} M, 5×10^{-12} M, 10^{-12} M, 5×10^{-13} M, 10^{-13} M, 5×10^{-14} M, 10^{-14} M, 5×10^{-15} M, or 10^{-15} M.

[290] The invention also provides antibodies that competitively inhibit binding of an antibody to an epitope of the invention as determined by any method known in the art for

determining competitive binding, for example, the immunoassays described herein. In preferred embodiments, the antibody competitively inhibits binding to the epitope by at least 95%, at least 90%, at least 85 %, at least 80%, at least 75%, at least 70%, at least 60%, or at least 50%.

[291] Antibodies of the present invention may act as agonists or antagonists of the polypeptides of the present invention. For example, the present invention includes antibodies which disrupt the receptor/ligand interactions with the polypeptides of the invention either partially or fully. Preferably, antibodies of the present invention bind an antigenic epitope disclosed herein, or a portion thereof. The invention features both receptor-specific antibodies and ligand-specific antibodies. The invention also features receptor-specific antibodies which do not prevent ligand binding but prevent receptor activation. Receptor activation (i.e., signaling) may be determined by techniques described herein or otherwise known in the art. For example, receptor activation can be determined by detecting the phosphorylation (e.g., tyrosine or serine/threonine) of the receptor or its substrate by immunoprecipitation followed by western blot analysis (for example, as described *supra*). In specific embodiments, antibodies are provided that inhibit ligand activity or receptor activity by at least 95%, at least 90%, at least 85%, at least 80%, at least 75%, at least 70%, at least 60%, or at least 50% of the activity in absence of the antibody.

[292] The invention also features receptor-specific antibodies which both prevent ligand binding and receptor activation as well as antibodies that recognize the receptor-ligand complex, and, preferably, do not specifically recognize the unbound receptor or the unbound ligand. Likewise, included in the invention are neutralizing antibodies which bind the ligand and prevent binding of the ligand to the receptor, as well as antibodies which bind the ligand, thereby preventing receptor activation, but do not prevent the ligand from binding the receptor. Further included in the invention are antibodies which activate the receptor. These antibodies may act as receptor agonists, i.e., potentiate or activate either all or a subset of the biological activities of the ligand-mediated receptor activation, for example, by inducing dimerization of the receptor. The antibodies may be specified as agonists, antagonists or inverse agonists for biological activities comprising the specific biological activities of the peptides of the invention disclosed herein. The above antibody agonists can be made using methods known in the art. See, e.g., PCT publication WO 96/40281; U.S. Patent No. 5,811,097; Deng et al., Blood 92(6):1981-1988 (1998); Chen et al., Cancer Res. 58(16):3668-3678 (1998); Harrop et al., J. Immunol. 161(4):1786-1794

(1998); Zhu et al., Cancer Res. 58(15):3209-3214 (1998); Yoon et al., J. Immunol. 160(7):3170-3179 (1998); Prat et al., J. Cell. Sci. 111(Pt2):237-247 (1998); Pitard et al., J. Immunol. Methods 205(2):177-190 (1997); Liautard et al., Cytokine 9(4):233-241 (1997); Carlson et al., J. Biol. Chem. 272(17):11295-11301 (1997); Taryman et al., Neuron 14(4):755-762 (1995); Muller et al., Structure 6(9):1153-1167 (1998); Bartunek et al., Cytokine 8(1):14-20 (1996) (which are all incorporated by reference herein in their entireties).

[293] Antibodies of the present invention may be used, for example, to purify, detect, and target the polypeptides of the present invention, including both *in vitro* and *in vivo* diagnostic and therapeutic methods. For example, the antibodies have utility in immunoassays for qualitatively and quantitatively measuring levels of the polypeptides of the present invention in biological samples. See, e.g., Harlow et al., Antibodies: A Laboratory Manual, (Cold Spring Harbor Laboratory Press, 2nd ed. 1988); incorporated by reference herein in its entirety.

[294] As discussed in more detail below, the antibodies of the present invention may be used either alone or in combination with other compositions. The antibodies may further be recombinantly fused to a heterologous polypeptide at the N- or C-terminus or chemically conjugated (including covalent and non-covalent conjugations) to polypeptides or other compositions. For example, antibodies of the present invention may be recombinantly fused or conjugated to molecules useful as labels in detection assays and effector molecules such as heterologous polypeptides, drugs, radionuclides, or toxins. See, e.g., PCT publications WO 92/08495; WO 91/14438; WO 89/12624; U.S. Patent No. 5,314,995; and EP 396,387; the disclosures of which are incorporated herein by reference in their entireties.

[295] The antibodies of the invention include derivatives that are modified, i.e., by the covalent attachment of any type of molecule to the antibody such that covalent attachment does not prevent the antibody from generating an anti-idiotypic response. For example, but not by way of limitation, the antibody derivatives include antibodies that have been modified, e.g., by glycosylation, acetylation, pegylation, phosphorylation, amidation, derivatization by known protecting/blocking groups, proteolytic cleavage, linkage to a cellular ligand or other protein, etc. Any of numerous chemical modifications may be carried out by known techniques, including, but not limited to specific chemical cleavage, acetylation, formylation, metabolic synthesis of tunicamycin, etc. Additionally, the derivative may contain one or more non-classical amino acids.

[296] The antibodies of the present invention may be generated by any suitable method known in the art. Polyclonal antibodies to an antigen-of- interest can be produced by various procedures well known in the art. For example, a polypeptide of the invention can be administered to various host animals including, but not limited to, rabbits, mice, rats, etc. to induce the production of sera containing polyclonal antibodies specific for the antigen. Various adjuvants may be used to increase the immunological response, depending on the host species, and include but are not limited to, Freund's (complete and incomplete), mineral gels such as aluminum hydroxide, surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanins, dinitrophenol, and potentially useful human adjuvants such as BCG (bacille Calmette-Guerin) and corynebacterium parvum. Such adjuvants are also well known in the art.

[297] Monoclonal antibodies can be prepared using a wide variety of techniques known in the art including the use of hybridoma, recombinant, and phage display technologies, or a combination thereof. For example, monoclonal antibodies can be produced using hybridoma techniques including those known in the art and taught, for example, in Harlow et al., *Antibodies: A Laboratory Manual*, (Cold Spring Harbor Laboratory Press, 2nd ed. 1988); Hammerling, et al., in: *Monoclonal Antibodies and T-Cell Hybridomas* 563-681 (Elsevier, N.Y., 1981) (said references incorporated by reference in their entireties). The term "monoclonal antibody" as used herein is not limited to antibodies produced through hybridoma technology. The term "monoclonal antibody" refers to an antibody that is derived from a single clone, including any eukaryotic, prokaryotic, or phage clone, and not the method by which it is produced.

[298] Methods for producing and screening for specific antibodies using hybridoma technology are routine and well known in the art and are discussed in detail in the Examples. In a non-limiting example, mice can be immunized with a polypeptide of the invention or a cell expressing such peptide. Once an immune response is detected, e.g., antibodies specific for the antigen are detected in the mouse serum, the mouse spleen is harvested and splenocytes isolated. The splenocytes are then fused by well known techniques to any suitable myeloma cells, for example cells from cell line SP20 available from the ATCC. Hybridomas are selected and cloned by limited dilution. The hybridoma clones are then assayed by methods known in the art for cells that secrete antibodies capable of binding a polypeptide of the invention. Ascites fluid, which generally contains

high levels of antibodies, can be generated by immunizing mice with positive hybridoma clones.

[299] Accordingly, the present invention provides methods of generating monoclonal antibodies as well as antibodies produced by the method comprising culturing a hybridoma cell secreting an antibody of the invention wherein, preferably, the hybridoma is generated by fusing splenocytes isolated from a mouse immunized with an antigen of the invention with myeloma cells and then screening the hybridomas resulting from the fusion for hybridoma clones that secrete an antibody able to bind a polypeptide of the invention.

[300] Another well known method for producing both polyclonal and monoclonal human B cell lines is transformation using Epstein Barr Virus (EBV). Protocols for generating EBV-transformed B cell lines are commonly known in the art, such as, for example, the protocol outlined in Chapter 7.22 of Current Protocols in Immunology, Coligan et al., Eds., 1994, John Wiley & Sons, NY, which is hereby incorporated in its entirety by reference. The source of B cells for transformation is commonly human peripheral blood, but B cells for transformation may also be derived from other sources including, but not limited to, lymph nodes, tonsil, spleen, tumor tissue, and infected tissues. Tissues are generally made into single cell suspensions prior to EBV transformation. Additionally, steps may be taken to either physically remove or inactivate T cells (e.g., by treatment with cyclosporin A) in B cell-containing samples, because T cells from individuals seropositive for anti-EBV antibodies can suppress B cell immortalization by EBV.

[301] In general, the sample containing human B cells is inoculated with EBV, and cultured for 3-4 weeks. A typical source of EBV is the culture supernatant of the B95-8 cell line (ATCC #VR-1492). Physical signs of EBV transformation can generally be seen towards the end of the 3-4 week culture period. By phase-contrast microscopy, transformed cells may appear large, clear, hairy and tend to aggregate in tight clusters of cells. Initially, EBV lines are generally polyclonal. However, over prolonged periods of cell cultures, EBV lines may become monoclonal or polyclonal as a result of the selective outgrowth of particular B cell clones. Alternatively, polyclonal EBV transformed lines may be subcloned (e.g., by limiting dilution culture) or fused with a suitable fusion partner and plated at limiting dilution to obtain monoclonal B cell lines. Suitable fusion partners for EBV transformed cell lines include mouse myeloma cell lines (e.g., SP2/0, X63-Ag8.653), heteromyeloma cell lines (human x mouse; e.g., SPAM-8, SBC-H20, and CB-F7), and

human cell lines (e.g., GM 1500, SKO-007, RPMI 8226, and KR-4). Thus, the present invention also provides a method of generating polyclonal or monoclonal human antibodies against polypeptides of the invention or fragments thereof, comprising EBV-transformation of human B cells.

[302] Antibody fragments which recognize specific epitopes may be generated by known techniques. For example, Fab and F(ab')₂ fragments of the invention may be produced by proteolytic cleavage of immunoglobulin molecules, using enzymes such as papain (to produce Fab fragments) or pepsin (to produce F(ab')₂ fragments). F(ab')₂ fragments contain the variable region, the light chain constant region and the CH1 domain of the heavy chain.

[303] For example, the antibodies of the present invention can also be generated using various phage display methods known in the art. In phage display methods, functional antibody domains are displayed on the surface of phage particles which carry the polynucleotide sequences encoding them. In a particular embodiment, such phage can be utilized to display antigen binding domains expressed from a repertoire or combinatorial antibody library (e.g., human or murine). Phage expressing an antigen binding domain that binds the antigen of interest can be selected or identified with antigen, e.g., using labeled antigen or antigen bound or captured to a solid surface or bead. Phage used in these methods are typically filamentous phage including fd and M13 binding domains expressed from phage with Fab, Fv or disulfide stabilized Fv antibody domains recombinantly fused to either the phage gene III or gene VIII protein. Examples of phage display methods that can be used to make the antibodies of the present invention include those disclosed in Brinkman et al., J. Immunol. Methods 182:41-50 (1995); Ames et al., J. Immunol. Methods 184:177-186 (1995); Kettleborough et al., Eur. J. Immunol. 24:952-958 (1994); Persic et al., Gene 187 9-18 (1997); Burton et al., Advances in Immunology 57:191-280 (1994); PCT application No. PCT/GB91/01134; PCT publications WO 90/02809; WO 91/10737; WO 92/01047; WO 92/18619; WO 93/11236; WO 95/15982; WO 95/20401; and U.S. Patent Nos. 5,698,426; 5,223,409; 5,403,484; 5,580,717; 5,427,908; 5,750,753; 5,821,047; 5,571,698; 5,427,908; 5,516,637; 5,780,225; 5,658,727; 5,733,743 and 5,969,108; each of which is incorporated herein by reference in its entirety.

[304] As described in the above references, after phage selection, the antibody coding regions from the phage can be isolated and used to generate whole antibodies, including human antibodies, or any other desired antigen binding fragment, and expressed in any

desired host, including mammalian cells, insect cells, plant cells, yeast, and bacteria, e.g., as described in detail below. For example, techniques to recombinantly produce Fab, Fab' and F(ab')₂ fragments can also be employed using methods known in the art such as those disclosed in PCT publication WO 92/22324; Mullinax et al., *BioTechniques* 12(6):864-869 (1992); and Sawai et al., *AJRI* 34:26-34 (1995); and Better et al., *Science* 240:1041-1043 (1988) (said references incorporated by reference in their entireties).

[305] Examples of techniques which can be used to produce single-chain Fvs and antibodies include those described in U.S. Patents 4,946,778 and 5,258,498; Huston et al., *Methods in Enzymology* 203:46-88 (1991); Shu et al., *PNAS* 90:7995-7999 (1993); and Skerra et al., *Science* 240:1038-1040 (1988). For some uses, including *in vivo* use of antibodies in humans and *in vitro* detection assays, it may be preferable to use chimeric, humanized, or human antibodies. A chimeric antibody is a molecule in which different portions of the antibody are derived from different animal species, such as antibodies having a variable region derived from a murine monoclonal antibody and a human immunoglobulin constant region. Methods for producing chimeric antibodies are known in the art. See e.g., Morrison, *Science* 229:1202 (1985); Oi et al., *BioTechniques* 4:214 (1986); Gillies et al., (1989) *J. Immunol. Methods* 125:191-202; U.S. Patent Nos. 5,807,715; 4,816,567; and 4,816,397, which are incorporated herein by reference in their entirety. Humanized antibodies are antibody molecules from non-human species antibody that binds the desired antigen having one or more complementarity determining regions (CDRs) from the non-human species and a framework regions from a human immunoglobulin molecule. Often, framework residues in the human framework regions will be substituted with the corresponding residue from the CDR donor antibody to alter, preferably improve, antigen binding. These framework substitutions are identified by methods well known in the art, e.g., by modeling of the interactions of the CDR and framework residues to identify framework residues important for antigen binding and sequence comparison to identify unusual framework residues at particular positions. (See, e.g., Queen et al., U.S. Patent No. 5,585,089; Riechmann et al., *Nature* 332:323 (1988), which are incorporated herein by reference in their entireties.) Antibodies can be humanized using a variety of techniques known in the art including, for example, CDR-grafting (EP 239,400; PCT publication WO 91/09967; U.S. Patent Nos. 5,225,539; 5,530,101; and 5,585,089), veneering or resurfacing (EP 592,106; EP 519,596; Padlan, *Molecular Immunology* 28(4/5):489-498 (1991); Studnicka et al., *Protein Engineering*

7(6):805-814 (1994); Roguska. et al., PNAS 91:969-973 (1994)), and chain shuffling (U.S. Patent No. 5,565,332).

[306] Completely human antibodies are particularly desirable for therapeutic treatment of human patients. Human antibodies can be made by a variety of methods known in the art including phage display methods described above using antibody libraries derived from human immunoglobulin sequences. See also, U.S. Patent Nos. 4,444,887 and 4,716,111; and PCT publications WO 98/46645, WO 98/50433, WO 98/24893, WO 98/16654, WO 96/34096, WO 96/33735, and WO 91/10741; each of which is incorporated herein by reference in its entirety.

[307] Human antibodies can also be produced using transgenic mice which are incapable of expressing functional endogenous immunoglobulins, but which can express human immunoglobulin genes. For example, the human heavy and light chain immunoglobulin gene complexes may be introduced randomly or by homologous recombination into mouse embryonic stem cells. Alternatively, the human variable region, constant region, and diversity region may be introduced into mouse embryonic stem cells in addition to the human heavy and light chain genes. The mouse heavy and light chain immunoglobulin genes may be rendered non-functional separately or simultaneously with the introduction of human immunoglobulin loci by homologous recombination. In particular, homozygous deletion of the JH region prevents endogenous antibody production. The modified embryonic stem cells are expanded and microinjected into blastocysts to produce chimeric mice. The chimeric mice are then bred to produce homozygous offspring which express human antibodies. The transgenic mice are immunized in the normal fashion with a selected antigen, e.g., all or a portion of a polypeptide of the invention. Monoclonal antibodies directed against the antigen can be obtained from the immunized, transgenic mice using conventional hybridoma technology. The human immunoglobulin transgenes harbored by the transgenic mice rearrange during B cell differentiation, and subsequently undergo class switching and somatic mutation. Thus, using such a technique, it is possible to produce therapeutically useful IgG, IgA, IgM and IgE antibodies. For an overview of this technology for producing human antibodies, see Lonberg and Huszar, Int. Rev. Immunol. 13:65-93 (1995). For a detailed discussion of this technology for producing human antibodies and human monoclonal antibodies and protocols for producing such antibodies, see, e.g., PCT publications WO 98/24893; WO 92/01047; WO 96/34096; WO 96/33735; European Patent No. 0 598 877; U.S. Patent Nos. 5,413,923; 5,625,126;

5,633,425; 5,569,825; 5,661,016; 5,545,806; 5,814,318; 5,885,793; 5,916,771; 5,939,598; 6,075,181; and 6,114,598, which are incorporated by reference herein in their entirety. In addition, companies such as Abgenix, Inc. (Freemont, CA) and Genpharm (San Jose, CA) can be engaged to provide human antibodies directed against a selected antigen using technology similar to that described above.

[308] Completely human antibodies which recognize a selected epitope can be generated using a technique referred to as "guided selection." In this approach a selected non-human monoclonal antibody, e.g., a mouse antibody, is used to guide the selection of a completely human antibody recognizing the same epitope. (Jespers et al., *Bio/technology* 12:899-903 (1988)).

[309] Further, antibodies to the polypeptides of the invention can, in turn, be utilized to generate anti-idiotypic antibodies that "mimic" polypeptides of the invention using techniques well known to those skilled in the art. (See, e.g., Greenspan & Bona, *FASEB J.* 7(5):437-444; (1989) and Nissinoff, J. *Immunol.* 147(8):2429-2438 (1991)). For example, antibodies which bind to and competitively inhibit polypeptide multimerization and/or binding of a polypeptide of the invention to a ligand can be used to generate anti-idiotypes that "mimic" the polypeptide multimerization and/or binding domain and, as a consequence, bind to and neutralize polypeptide and/or its ligand. Such neutralizing anti-idiotypes or Fab fragments of such anti-idiotypes can be used in therapeutic regimens to neutralize polypeptide ligand(s)/receptor(s). For example, such anti-idiotypic antibodies can be used to bind a polypeptide of the invention and/or to bind its ligand(s)/receptor(s), and thereby block its biological activity. Alternatively, antibodies which bind to and enhance polypeptide multimerization and/or binding, and/or receptor/ligand multimerization, binding and/or signaling can be used to generate anti-idiotypes that function as agonists of a polypeptide of the invention and/or its ligand/receptor. Such agonistic anti-idiotypes or Fab fragments of such anti-idiotypes can be used in therapeutic regimens as agonists of the polypeptides of the invention or its ligand(s)/receptor(s). For example, such anti-idiotypic antibodies can be used to bind a polypeptide of the invention and/or to bind its ligand(s)/receptor(s), and thereby promote or enhance its biological activity.

[310] Intrabodies of the invention can be produced using methods known in the art, such as those disclosed and reviewed in Chen et al., *Hum. Gene Ther.* 5:595-601 (1994); Marasco, W.A., *Gene Ther.* 4:11-15 (1997); Rondon and Marasco, *Annu. Rev. Microbiol.* 51:257-283 (1997); Proba et al., *J. Mol. Biol.* 275:245-253 (1998); Cohen et al., *Oncogene*

17:2445-2456 (1998); Ohage and Steipe, J. Mol. Biol. 291:1119-1128 (1999); Ohage et al., J. Mol. Biol. 291:1129-1134 (1999); Wirtz and Steipe, Protein Sci. 8:2245-2250 (1999); Zhu et al., J. Immunol. Methods 231:207-222 (1999); and references cited therein.

Polynucleotides Encoding Antibodies

[311] The invention further provides polynucleotides comprising a nucleotide sequence encoding an antibody of the invention and fragments thereof. The invention also encompasses polynucleotides that hybridize under stringent or alternatively, under lower stringency hybridization conditions, e.g., as defined *supra*, to polynucleotides that encode an antibody, preferably, that specifically binds to a polypeptide of the invention, preferably, an antibody that binds to a polypeptide having the amino acid sequence of SEQ ID NO:Y, to a polypeptide encoded by a portion of SEQ ID NO:X as defined in columns 8 and 9 of Table 2, and/or to a polypeptide encoded by the cDNA contained in Clone ID NO:Z.

[312] The polynucleotides may be obtained, and the nucleotide sequence of the polynucleotides determined, by any method known in the art. For example, if the nucleotide sequence of the antibody is known, a polynucleotide encoding the antibody may be assembled from chemically synthesized oligonucleotides (e.g., as described in Kutmeier et al., BioTechniques 17:242 (1994)), which, briefly, involves the synthesis of overlapping oligonucleotides containing portions of the sequence encoding the antibody, annealing and ligating of those oligonucleotides, and then amplification of the ligated oligonucleotides by PCR.

[313] Alternatively, a polynucleotide encoding an antibody may be generated from nucleic acid from a suitable source. If a clone containing a nucleic acid encoding a particular antibody is not available, but the sequence of the antibody molecule is known, a nucleic acid encoding the immunoglobulin may be chemically synthesized or obtained from a suitable source (e.g., an antibody cDNA library, or a cDNA library generated from, or nucleic acid, preferably poly A+ RNA, isolated from, any tissue or cells expressing the antibody, such as hybridoma cells selected to express an antibody of the invention) by PCR amplification using synthetic primers hybridizable to the 3' and 5' ends of the sequence or by cloning using an oligonucleotide probe specific for the particular gene sequence to identify, e.g., a cDNA clone from a cDNA library that encodes the antibody. Amplified nucleic acids generated by PCR may then be cloned into replicable cloning vectors using any method well known in the art.

[314] Once the nucleotide sequence and corresponding amino acid sequence of the antibody is determined, the nucleotide sequence of the antibody may be manipulated using methods well known in the art for the manipulation of nucleotide sequences, e.g., recombinant DNA techniques, site directed mutagenesis, PCR, etc. (see, for example, the techniques described in Sambrook et al., 1990, *Molecular Cloning, A Laboratory Manual*, 2d Ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, NY and Ausubel et al., eds., 1998, *Current Protocols in Molecular Biology*, John Wiley & Sons, NY, which are both incorporated by reference herein in their entireties), to generate antibodies having a different amino acid sequence, for example to create amino acid substitutions, deletions, and/or insertions.

[315] In a specific embodiment, the amino acid sequence of the heavy and/or light chain variable domains may be inspected to identify the sequences of the complementarity determining regions (CDRs) by methods that are well known in the art, e.g., by comparison to known amino acid sequences of other heavy and light chain variable regions to determine the regions of sequence hypervariability. Using routine recombinant DNA techniques, one or more of the CDRs may be inserted within framework regions, e.g., into human framework regions to humanize a non-human antibody, as described *supra*. The framework regions may be naturally occurring or consensus framework regions, and preferably human framework regions (see, e.g., Chothia et al., *J. Mol. Biol.* 278: 457-479 (1998) for a listing of human framework regions). Preferably, the polynucleotide generated by the combination of the framework regions and CDRs encodes an antibody that specifically binds a polypeptide of the invention. Preferably, as discussed *supra*, one or more amino acid substitutions may be made within the framework regions, and, preferably, the amino acid substitutions improve binding of the antibody to its antigen. Additionally, such methods may be used to make amino acid substitutions or deletions of one or more variable region cysteine residues participating in an intrachain disulfide bond to generate antibody molecules lacking one or more intrachain disulfide bonds. Other alterations to the polynucleotide are encompassed by the present invention and within the skill of the art.

[316] In addition, techniques developed for the production of "chimeric antibodies" (Morrison et al., *Proc. Natl. Acad. Sci.* 81:851-855 (1984); Neuberger et al., *Nature* 312:604-608 (1984); Takeda et al., *Nature* 314:452-454 (1985)) by splicing genes from a mouse antibody molecule of appropriate antigen specificity together with genes from a human antibody molecule of appropriate biological activity can be used. As described

supra, a chimeric antibody is a molecule in which different portions are derived from different animal species, such as those having a variable region derived from a murine mAb and a human immunoglobulin constant region, e.g., humanized antibodies.

[317] Alternatively, techniques described for the production of single chain antibodies (U.S. Patent No. 4,946,778; Bird, Science 242:423- 42 (1988); Huston et al., Proc. Natl. Acad. Sci. USA 85:5879-5883 (1988); and Ward et al., Nature 334:544-54 (1989)) can be adapted to produce single chain antibodies. Single chain antibodies are formed by linking the heavy and light chain fragments of the Fv region via an amino acid bridge, resulting in a single chain polypeptide. Techniques for the assembly of functional Fv fragments in *E. coli* may also be used (Skerra et al., Science 242:1038- 1041 (1988)).

Methods of Producing Antibodies

[318] The antibodies of the invention can be produced by any method known in the art for the synthesis of antibodies, in particular, by chemical synthesis or preferably, by recombinant expression techniques. Methods of producing antibodies include, but are not limited to, hybridoma technology, EBV transformation, and other methods discussed herein as well as through the use recombinant DNA technology, as discussed below.

[319] Recombinant expression of an antibody of the invention, or fragment, derivative or analog thereof, (e.g., a heavy or light chain of an antibody of the invention or a single chain antibody of the invention), requires construction of an expression vector containing a polynucleotide that encodes the antibody. Once a polynucleotide encoding an antibody molecule or a heavy or light chain of an antibody, or portion thereof (preferably containing the heavy or light chain variable domain), of the invention has been obtained, the vector for the production of the antibody molecule may be produced by recombinant DNA technology using techniques well known in the art. Thus, methods for preparing a protein by expressing a polynucleotide containing an antibody encoding nucleotide sequence are described herein. Methods which are well known to those skilled in the art can be used to construct expression vectors containing antibody coding sequences and appropriate transcriptional and translational control signals. These methods include, for example, *in vitro* recombinant DNA techniques, synthetic techniques, and *in vivo* genetic recombination. The invention, thus, provides replicable vectors comprising a nucleotide sequence encoding an antibody molecule of the invention, or a heavy or light chain thereof, or a heavy or light chain variable domain, operably linked to a promoter. Such vectors

may include the nucleotide sequence encoding the constant region of the antibody molecule (see, e.g., PCT Publication WO 86/05807; PCT Publication WO 89/01036; and U.S. Patent No. 5,122,464) and the variable domain of the antibody may be cloned into such a vector for expression of the entire heavy or light chain.

[320] The expression vector is transferred to a host cell by conventional techniques and the transfected cells are then cultured by conventional techniques to produce an antibody of the invention. Thus, the invention includes host cells containing a polynucleotide encoding an antibody of the invention, or a heavy or light chain thereof, or a single chain antibody of the invention, operably linked to a heterologous promoter. In preferred embodiments for the expression of double-chained antibodies, vectors encoding both the heavy and light chains may be co-expressed in the host cell for expression of the entire immunoglobulin molecule, as detailed below.

[321] A variety of host-expression vector systems may be utilized to express the antibody molecules of the invention. Such host-expression systems represent vehicles by which the coding sequences of interest may be produced and subsequently purified, but also represent cells which may, when transformed or transfected with the appropriate nucleotide coding sequences, express an antibody molecule of the invention in situ. These include but are not limited to microorganisms such as bacteria (e.g., *E. coli*, *B. subtilis*) transformed with recombinant bacteriophage DNA, plasmid DNA or cosmid DNA expression vectors containing antibody coding sequences; yeast (e.g., *Saccharomyces*, *Pichia*) transformed with recombinant yeast expression vectors containing antibody coding sequences; insect cell systems infected with recombinant virus expression vectors (e.g., baculovirus) containing antibody coding sequences; plant cell systems infected with recombinant virus expression vectors (e.g., cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or transformed with recombinant plasmid expression vectors (e.g., Ti plasmid) containing antibody coding sequences; or mammalian cell systems (e.g., COS, CHO, BHK, 293, 3T3 cells) harboring recombinant expression constructs containing promoters derived from the genome of mammalian cells (e.g., metallothionein promoter) or from mammalian viruses (e.g., the adenovirus late promoter; the vaccinia virus 7.5K promoter). Preferably, bacterial cells such as *Escherichia coli*, and more preferably, eukaryotic cells, especially for the expression of whole recombinant antibody molecule, are used for the expression of a recombinant antibody molecule. For example, mammalian cells such as Chinese hamster ovary cells (CHO), in conjunction with a vector such as the major intermediate early gene

promoter element from human cytomegalovirus is an effective expression system for antibodies (Foecking et al., Gene 45:101 (1986); Cockett et al., Bio/Technology 8:2 (1990)).

[322] In bacterial systems, a number of expression vectors may be advantageously selected depending upon the use intended for the antibody molecule being expressed. For example, when a large quantity of such a protein is to be produced, for the generation of pharmaceutical compositions of an antibody molecule, vectors which direct the expression of high levels of fusion protein products that are readily purified may be desirable. Such vectors include, but are not limited, to the E. coli expression vector pUR278 (Ruther et al., EMBO J. 2:1791 (1983)), in which the antibody coding sequence may be ligated individually into the vector in frame with the lac Z coding region so that a fusion protein is produced; pIN vectors (Inouye & Inouye, Nucleic Acids Res. 13:3101-3109 (1985); Van Heeke & Schuster, J. Biol. Chem. 24:5503-5509 (1989)); and the like. pGEX vectors may also be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption and binding to matrix glutathione-agarose beads followed by elution in the presence of free glutathione. The pGEX vectors are designed to include thrombin or factor Xa protease cleavage sites so that the cloned target gene product can be released from the GST moiety.

[323] In an insect system, Autographa californica nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes. The virus grows in *Spodoptera frugiperda* cells. The antibody coding sequence may be cloned individually into non-essential regions (for example the polyhedrin gene) of the virus and placed under control of an AcNPV promoter (for example the polyhedrin promoter).

[324] In mammalian host cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, the antibody coding sequence of interest may be ligated to an adenovirus transcription/translation control complex, e.g., the late promoter and tripartite leader sequence. This chimeric gene may then be inserted in the adenovirus genome by in vitro or *in vivo* recombination. Insertion in a non-essential region of the viral genome (e.g., region E1 or E3) will result in a recombinant virus that is viable and capable of expressing the antibody molecule in infected hosts. (e.g., see Logan & Shenk, Proc. Natl. Acad. Sci. USA 81:355-359 (1984)). Specific initiation signals may also be required for efficient translation of inserted antibody

coding sequences. These signals include the ATG initiation codon and adjacent sequences. Furthermore, the initiation codon must be in phase with the reading frame of the desired coding sequence to ensure translation of the entire insert. These exogenous translational control signals and initiation codons can be of a variety of origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of appropriate transcription enhancer elements, transcription terminators, etc. (see Bittner et al., *Methods, in Enzymol.* 153:51-544 (1987)).

[325] In addition, a host cell strain may be chosen which modulates the expression of the inserted sequences, or modifies and processes the gene product in the specific fashion desired. Such modifications (e.g., glycosylation) and processing (e.g., cleavage) of protein products may be important for the function of the protein. Different host cells have characteristic and specific mechanisms for the post-translational processing and modification of proteins and gene products. Appropriate cell lines or host systems can be chosen to ensure the correct modification and processing of the foreign protein expressed. To this end, eukaryotic host cells which possess the cellular machinery for proper processing of the primary transcript, glycosylation, and phosphorylation of the gene product may be used. Such mammalian host cells include but are not limited to CHO, VERY, BHK, Hela, COS, MDCK, 293, 3T3, WI38, and in particular, breast cancer cell lines such as, for example, BT483, Hs578T, HTB2, BT20 and T47D, and normal mammary gland cell line such as, for example, CRL7030 and Hs578Bst.

[326] For long-term, high-yield production of recombinant proteins, stable expression is preferred. For example, cell lines which stably express the antibody molecule may be engineered. Rather than using expression vectors which contain viral origins of replication, host cells can be transformed with DNA controlled by appropriate expression control elements (e.g., promoter, enhancer, sequences, transcription terminators, polyadenylation sites, etc.), and a selectable marker. Following the introduction of the foreign DNA, engineered cells may be allowed to grow for 1-2 days in an enriched media, and then are switched to a selective media. The selectable marker in the recombinant plasmid confers resistance to the selection and allows cells to stably integrate the plasmid into their chromosomes and grow to form foci which in turn can be cloned and expanded into cell lines. This method may advantageously be used to engineer cell lines which express the antibody molecule. Such engineered cell lines may be particularly useful in screening and evaluation of compounds that interact directly or indirectly with the antibody molecule.

[327] A number of selection systems may be used, including but not limited to the herpes simplex virus thymidine kinase (Wigler et al., *Cell* 11:223 (1977)), hypoxanthine-guanine phosphoribosyltransferase (Szybalska & Szybalski, *Proc. Natl. Acad. Sci. USA* 48:202 (1992)), and adenine phosphoribosyltransferase (Lowy et al., *Cell* 22:817 (1980)) genes can be employed in tk-, hgprt- or aprt- cells, respectively. Also, antimetabolite resistance can be used as the basis of selection for the following genes: dhfr, which confers resistance to methotrexate (Wigler et al., *Natl. Acad. Sci. USA* 77:357 (1980); O'Hare et al., *Proc. Natl. Acad. Sci. USA* 78:1527 (1981)); gpt, which confers resistance to mycophenolic acid (Mulligan & Berg, *Proc. Natl. Acad. Sci. USA* 78:2072 (1981)); neo, which confers resistance to the aminoglycoside G-418 *Clinical Pharmacy* 12:488-505; Wu and Wu, *Biotherapy* 3:87-95 (1991); Tolstoshev, *Ann. Rev. Pharmacol. Toxicol.* 32:573-596 (1993); Mulligan, *Science* 260:926-932 (1993); and Morgan and Anderson, *Ann. Rev. Biochem.* 62:191-217 (1993); May, 1993, *TIB TECH* 11(5):155-215 (1993)); and hygromycin (Santerre et al., *Gene* 30:147 (1984)). Methods commonly known in the art of recombinant DNA technology may be routinely applied to select the desired recombinant clone, and such methods are described, for example, in Ausubel et al. (eds.), *Current Protocols in Molecular Biology*, John Wiley & Sons, NY (1993); Kriegler, *Gene Transfer and Expression, A Laboratory Manual*, Stockton Press, NY (1990); and in Chapters 12 and 13, Dracopoli et al. (eds), *Current Protocols in Human Genetics*, John Wiley & Sons, NY (1994); Colberre-Garapin et al., *J. Mol. Biol.* 150:1 (1981), which are incorporated by reference herein in their entirety.

[328] The expression levels of an antibody molecule can be increased by vector amplification (for a review, see Bebbington and Hentschel, *The use of vectors based on gene amplification for the expression of cloned genes in mammalian cells in DNA cloning*, Vol.3. (Academic Press, New York, 1987)). When a marker in the vector system expressing antibody is amplifiable, increase in the level of inhibitor present in culture of host cell will increase the number of copies of the marker gene. Since the amplified region is associated with the antibody gene, production of the antibody will also increase (Crouse et al., *Mol. Cell. Biol.* 3:257 (1983)).

[329] Vectors which use glutamine synthase (GS) or DHFR as the selectable markers can be amplified in the presence of the drugs methionine sulphoximine or methotrexate, respectively. An advantage of glutamine synthase based vectors are the availability of cell lines (e.g., the murine myeloma cell line, NS0) which are glutamine synthase negative.

Glutamine synthase expression systems can also function in glutamine synthase expressing cells (e.g. Chinese Hamster Ovary (CHO) cells) by providing additional inhibitor to prevent the functioning of the endogenous gene. A glutamine synthase expression system and components thereof are detailed in PCT publications: WO87/04462; WO86/05807; WO89/01036; WO89/10404; and WO91/06657 which are incorporated in their entireties by reference herein. Additionally, glutamine synthase expression vectors that may be used according to the present invention are commercially available from suppliers, including, for example Lonza Biologics, Inc. (Portsmouth, NH). Expression and production of monoclonal antibodies using a GS expression system in murine myeloma cells is described in Bebbington *et al.*, *Bio/technology* 10:169(1992) and in Biblia and Robinson *Biotechnol. Prog.* 11:1 (1995) which are incorporated in their entireties by reference herein.

[330] The host cell may be co-transfected with two expression vectors of the invention, the first vector encoding a heavy chain derived polypeptide and the second vector encoding a light chain derived polypeptide. The two vectors may contain identical selectable markers which enable equal expression of heavy and light chain polypeptides. Alternatively, a single vector may be used which encodes, and is capable of expressing, both heavy and light chain polypeptides. In such situations, the light chain should be placed before the heavy chain to avoid an excess of toxic free heavy chain (Proudfoot, *Nature* 322:52 (1986); Kohler, *Proc. Natl. Acad. Sci. USA* 77:2197 (1980)). The coding sequences for the heavy and light chains may comprise cDNA or genomic DNA.

[331] Once an antibody molecule of the invention has been produced by an animal, chemically synthesized, or recombinantly expressed, it may be purified by any method known in the art for purification of an immunoglobulin molecule, for example, by chromatography (e.g., ion exchange, affinity, particularly by affinity for the specific antigen after Protein A, and sizing column chromatography), centrifugation, differential solubility, or by any other standard technique for the purification of proteins. In addition, the antibodies of the present invention or fragments thereof can be fused to heterologous polypeptide sequences described herein or otherwise known in the art, to facilitate purification.

[332] The present invention encompasses antibodies recombinantly fused or chemically conjugated (including both covalently and non-covalently conjugations) to a polypeptide (or portion thereof, preferably at least 10, 20, 30, 40, 50, 60, 70, 80, 90 or 100 amino acids of the polypeptide) of the present invention to generate fusion proteins. The fusion does

not necessarily need to be direct, but may occur through linker sequences. The antibodies may be specific for antigens other than polypeptides (or portion thereof, preferably at least 10, 20, 30, 40, 50, 60, 70, 80, 90 or 100 amino acids of the polypeptide) of the present invention. For example, antibodies may be used to target the polypeptides of the present invention to particular cell types, either *in vitro* or *in vivo*, by fusing or conjugating the polypeptides of the present invention to antibodies specific for particular cell surface receptors. Antibodies fused or conjugated to the polypeptides of the present invention may also be used in *in vitro* immunoassays and purification methods using methods known in the art. See e.g., Harbor et al., *supra*, and PCT publication WO 93/21232; EP 439,095; Naramura et al., Immunol. Lett. 39:91-99 (1994); U.S. Patent 5,474,981; Gillies et al., PNAS 89:1428-1432 (1992); Fell et al., J. Immunol. 146:2446-2452 (1991), which are incorporated by reference in their entireties.

[333] The present invention further includes compositions comprising the polypeptides of the present invention fused or conjugated to antibody domains other than the variable regions. For example, the polypeptides of the present invention may be fused or conjugated to an antibody Fc region, or portion thereof. The antibody portion fused to a polypeptide of the present invention may comprise the constant region, hinge region, CH1 domain, CH2 domain, and CH3 domain or any combination of whole domains or portions thereof. The polypeptides may also be fused or conjugated to the above antibody portions to form multimers. For example, Fc portions fused to the polypeptides of the present invention can form dimers through disulfide bonding between the Fc portions. Higher multimeric forms can be made by fusing the polypeptides to portions of IgA and IgM. Methods for fusing or conjugating the polypeptides of the present invention to antibody portions are known in the art. See, e.g., U.S. Patent Nos. 5,336,603; 5,622,929; 5,359,046; 5,349,053; 5,447,851; 5,112,946; EP 307,434; EP 367,166; PCT publications WO 96/04388; WO 91/06570; Ashkenazi et al., Proc. Natl. Acad. Sci. USA 88:10535-10539 (1991); Zheng et al., J. Immunol. 154:5590-5600 (1995); and Vil et al., Proc. Natl. Acad. Sci. USA 89:11337-11341 (1992) (said references incorporated by reference in their entireties).

[334] As discussed, *supra*, the polypeptides corresponding to a polypeptide, polypeptide fragment, or a variant of SEQ ID NO:Y may be fused or conjugated to the above antibody portions to increase the *in vivo* half life of the polypeptides or for use in immunoassays using methods known in the art. Further, the polypeptides corresponding to SEQ ID NO:Y may be fused or conjugated to the above antibody portions to facilitate

purification. One reported example describes chimeric proteins consisting of the first two domains of the human CD4-polypeptide and various domains of the constant regions of the heavy or light chains of mammalian immunoglobulins. See EP 394,827; and Traunecker et al., *Nature* 331:84-86 (1988). The polypeptides of the present invention fused or conjugated to an antibody having disulfide-linked dimeric structures (due to the IgG) may also be more efficient in binding and neutralizing other molecules, than the monomeric secreted protein or protein fragment alone. See, for example, Fountoulakis et al., *J. Biochem.* 270:3958-3964 (1995). In many cases, the Fc part in a fusion protein is beneficial in therapy and diagnosis, and thus can result in, for example, improved pharmacokinetic properties. See, for example, EP A 232,262. Alternatively, deleting the Fc part after the fusion protein has been expressed, detected, and purified, would be desired. For example, the Fc portion may hinder therapy and diagnosis if the fusion protein is used as an antigen for immunizations. In drug discovery, for example, human proteins, such as hIL-5, have been fused with Fc portions for the purpose of high-throughput screening assays to identify antagonists of hIL-5. (See, Bennett et al., *J. Molecular Recognition* 8:52-58 (1995); Johanson et al., *J. Biol. Chem.* 270:9459-9471 (1995)).

[335] Moreover, the antibodies or fragments thereof of the present invention can be fused to marker sequences, such as a peptide to facilitate purification. In preferred embodiments, the marker amino acid sequence is a hexa-histidine peptide, such as the tag provided in a pQE vector (QIAGEN, Inc., 9259 Eton Avenue, Chatsworth, CA, 91311), among others, many of which are commercially available. As described in Gentz et al., *Proc. Natl. Acad. Sci. USA* 86:821-824 (1989), for instance, hexa-histidine provides for convenient purification of the fusion protein. Other peptide tags useful for purification include, but are not limited to, the "HA" tag, which corresponds to an epitope derived from the influenza hemagglutinin protein (Wilson et al., *Cell* 37:767 (1984)) and the "flag" tag.

[336] The present invention further encompasses antibodies or fragments thereof conjugated to a diagnostic or therapeutic agent. The antibodies can be used diagnostically to, for example, monitor the development or progression of a tumor as part of a clinical testing procedure to, e.g., determine the efficacy of a given treatment regimen. Detection can be facilitated by coupling the antibody to a detectable substance. Examples of detectable substances include various enzymes, prosthetic groups, fluorescent materials, luminescent materials, bioluminescent materials, radioactive materials, positron emitting metals using various positron emission tomographies, and nonradioactive paramagnetic

metal ions. The detectable substance may be coupled or conjugated either directly to the antibody (or fragment thereof) or indirectly, through an intermediate (such as, for example, a linker known in the art) using techniques known in the art. See, for example, U.S. Patent No. 4,741,900 for metal ions which can be conjugated to antibodies for use as diagnostics according to the present invention. Examples of suitable enzymes include horseradish peroxidase, alkaline phosphatase, beta-galactosidase, or acetylcholinesterase; examples of suitable prosthetic group complexes include streptavidin/biotin and avidin/biotin; examples of suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; an example of a luminescent material includes luminol; examples of bioluminescent materials include luciferase, luciferin, and aequorin; and examples of suitable radioactive material include ^{125}I , ^{131}I , ^{111}In or ^{99}Tc .

[337] Further, an antibody or fragment thereof may be conjugated to a therapeutic moiety such as a cytotoxin, e.g., a cytostatic or cytotoxic agent, a therapeutic agent or a radioactive metal ion, e.g., alpha-emitters such as, for example, ^{213}Bi . A cytotoxin or cytotoxic agent includes any agent that is detrimental to cells. Examples include paclitaxol, cytochalasin B, gramicidin D, ethidium bromide, emetine, mitomycin, etoposide, teniposide, vincristine, vinblastine, colchicin, doxorubicin, daunorubicin, dihydroxy anthracin dione, mitoxantrone, mithramycin, actinomycin D, 1-dehydrotestosterone, glucocorticoids, procaine, tetracaine, lidocaine, propranolol, and puromycin and analogs or homologs thereof. Therapeutic agents include, but are not limited to, antimetabolites (e.g., methotrexate, 6-mercaptopurine, 6-thioguanine, cytarabine, 5-fluorouracil decarbazine), alkylating agents (e.g., mechlorethamine, thioepa chlorambucil, melphalan, carmustine (BSNU) and lomustine (CCNU), cyclophosphamide, busulfan, dibromomannitol, streptozotocin, mitomycin C, and cis- dichlorodiamine platinum (II) (DDP) cisplatin), anthracyclines (e.g., daunorubicin (formerly daunomycin) and doxorubicin), antibiotics (e.g., dactinomycin (formerly actinomycin), bleomycin, mithramycin, and anthramycin (AMC)), and anti-mitotic agents (e.g., vincristine and vinblastine).

[338] The conjugates of the invention can be used for modifying a given biological response, the therapeutic agent or drug moiety is not to be construed as limited to classical chemical therapeutic agents. For example, the drug moiety may be a protein or polypeptide possessing a desired biological activity. Such proteins may include, for example, a toxin such as abrin, ricin A, pseudomonas exotoxin, or diphtheria toxin; a protein such as tumor

necrosis factor, α -interferon, β -interferon, nerve growth factor, platelet derived growth factor, tissue plasminogen activator, an apoptotic agent, e.g., TNF- α , TNF- β , AIM I (See, International Publication No. WO 97/33899), AIM II (See, International Publication No. WO 97/34911), Fas Ligand (Takahashi *et al.*, *Int. Immunol.*, 6:1567-1574 (1994)), VEGI (See, International Publication No. WO 99/23105), a thrombotic agent or an anti-angiogenic agent, e.g., angiostatin or endostatin; or, biological response modifiers such as, for example, lymphokines, interleukin-1 ("IL-1"), interleukin-2 ("IL-2"), interleukin-6 ("IL-6"), granulocyte macrophage colony stimulating factor ("GM-CSF"), granulocyte colony stimulating factor ("G-CSF"), or other growth factors.

[339] Antibodies may also be attached to solid supports, which are particularly useful for immunoassays or purification of the target antigen. Such solid supports include, but are not limited to, glass, cellulose, polyacrylamide, nylon, polystyrene, polyvinyl chloride or polypropylene.

[340] Techniques for conjugating such therapeutic moiety to antibodies are well known. See, for example, Arnon *et al.*, "Monoclonal Antibodies For Immunotargeting Of Drugs In Cancer Therapy", in *Monoclonal Antibodies And Cancer Therapy*, Reisfeld *et al.* (eds.), pp. 243-56 (Alan R. Liss, Inc. 1985); Hellstrom *et al.*, "Antibodies For Drug Delivery", in *Controlled Drug Delivery* (2nd Ed.), Robinson *et al.* (eds.), pp. 623-53 (Marcel Dekker, Inc. 1987); Thorpe, "Antibody Carriers Of Cytotoxic Agents In Cancer Therapy: A Review", in *Monoclonal Antibodies '84: Biological And Clinical Applications*, Pinchera *et al.* (eds.), pp. 475-506 (1985); "Analysis, Results, And Future Prospective Of The Therapeutic Use Of Radiolabeled Antibody In Cancer Therapy", in *Monoclonal Antibodies For Cancer Detection And Therapy*, Baldwin *et al.* (eds.), pp. 303-16 (Academic Press 1985), and Thorpe *et al.*, "The Preparation And Cytotoxic Properties Of Antibody-Toxin Conjugates", *Immunol. Rev.* 62:119-58 (1982).

[341] Alternatively, an antibody can be conjugated to a second antibody to form an antibody heteroconjugate as described by Segal in U.S. Patent No. 4,676,980, which is incorporated herein by reference in its entirety.

[342] An antibody, with or without a therapeutic moiety conjugated to it, administered alone or in combination with cytotoxic factor(s) and/or cytokine(s) can be used as a therapeutic.

Immunophenotyping

[343] The antibodies of the invention may be utilized for immunophenotyping of cell lines and biological samples. Translation products of the gene of the present invention may be useful as cell-specific markers, or more specifically as cellular markers that are differentially expressed at various stages of differentiation and/or maturation of particular cell types. Monoclonal antibodies directed against a specific epitope, or combination of epitopes, will allow for the screening of cellular populations expressing the marker. Various techniques can be utilized using monoclonal antibodies to screen for cellular populations expressing the marker(s), and include magnetic separation using antibody-coated magnetic beads, "panning" with antibody attached to a solid matrix (i.e., plate), and flow cytometry (See, e.g., U.S. Patent 5,985,660; and Morrison *et al.*, *Cell*, 96:737-49 (1999)).

[344] These techniques allow for the screening of particular populations of cells, such as might be found with hematological malignancies (i.e. minimal residual disease (MRD) in acute leukemic patients) and "non-self" cells in transplantations to prevent Graft-versus-Host Disease (GVHD). Alternatively, these techniques allow for the screening of hematopoietic stem and progenitor cells capable of undergoing proliferation and/or differentiation, as might be found in human umbilical cord blood.

Assays For Antibody Binding

[345] The antibodies of the invention may be assayed for immunospecific binding by any method known in the art. The immunoassays which can be used include but are not limited to competitive and non-competitive assay systems using techniques such as western blots, radioimmunoassays, ELISA (enzyme linked immunosorbent assay), "sandwich" immunoassays, immunoprecipitation assays, precipitin reactions, gel diffusion precipitin reactions, immunodiffusion assays, agglutination assays, complement-fixation assays, immunoradiometric assays, fluorescent immunoassays, and protein A immunoassays, to name but a few. Such assays are routine and well known in the art (see, e.g., Ausubel *et al.*, eds, 1994, *Current Protocols in Molecular Biology*, Vol. 1, John Wiley & Sons, Inc., New York, which is incorporated by reference herein in its entirety). Exemplary immunoassays are described briefly below (but are not intended by way of limitation).

[346] Immunoprecipitation protocols generally comprise lysing a population of cells in a lysis buffer such as RIPA buffer (1% NP-40 or Triton X- 100, 1% sodium deoxycholate, 0.1% SDS, 0.15 M NaCl, 0.01 M sodium phosphate at pH 7.2, 1% Trasylol) supplemented

with protein phosphatase and/or protease inhibitors (e.g., EDTA, PMSF, aprotinin, sodium vanadate), adding the antibody of interest to the cell lysate, incubating for a period of time (e.g., 1-4 hours) at 4° C, adding protein A and/or protein G sepharose beads to the cell lysate, incubating for about an hour or more at 4° C, washing the beads in lysis buffer and resuspending the beads in SDS/sample buffer. The ability of the antibody of interest to immunoprecipitate a particular antigen can be assessed by, e.g., western blot analysis. One of skill in the art would be knowledgeable as to the parameters that can be modified to increase the binding of the antibody to an antigen and decrease the background (e.g., pre-clearing the cell lysate with sepharose beads). For further discussion regarding immunoprecipitation protocols see, e.g., Ausubel et al., eds., (1994), Current Protocols in Molecular Biology, Vol. 1, John Wiley & Sons, Inc., New York, section 10.16.1.

[347] Western blot analysis generally comprises preparing protein samples, electrophoresis of the protein samples in a polyacrylamide gel (e.g., 8%- 20% SDS-PAGE depending on the molecular weight of the antigen), transferring the protein sample from the polyacrylamide gel to a membrane such as nitrocellulose, PVDF or nylon, blocking the membrane in blocking solution (e.g., PBS with 3% BSA or non-fat milk), washing the membrane in washing buffer (e.g., PBS-Tween 20), blocking the membrane with primary antibody (the antibody of interest) diluted in blocking buffer, washing the membrane in washing buffer, blocking the membrane with a secondary antibody (which recognizes the primary antibody, e.g., an anti-human antibody) conjugated to an enzymatic substrate (e.g., horseradish peroxidase or alkaline phosphatase) or radioactive molecule (e.g., ³²P or ¹²⁵I) diluted in blocking buffer, washing the membrane in wash buffer, and detecting the presence of the antigen. One of skill in the art would be knowledgeable as to the parameters that can be modified to increase the signal detected and to reduce the background noise. For further discussion regarding western blot protocols see, e.g., Ausubel et al, eds, (1994), Current Protocols in Molecular Biology, Vol. 1, John Wiley & Sons, Inc., New York, section 10.8.1.

[348] ELISAs comprise preparing antigen, coating the well of a 96 well microtiter plate with the antigen, adding the antibody of interest conjugated to a detectable compound such as an enzymatic substrate (e.g., horseradish peroxidase or alkaline phosphatase) to the well and incubating for a period of time, and detecting the presence of the antigen. In ELISAs the antibody of interest does not have to be conjugated to a detectable compound; instead, a second antibody (which recognizes the antibody of interest) conjugated to a detectable

compound may be added to the well. Further, instead of coating the well with the antigen, the antibody may be coated to the well. In this case, a second antibody conjugated to a detectable compound may be added following the addition of the antigen of interest to the coated well. One of skill in the art would be knowledgeable as to the parameters that can be modified to increase the signal detected as well as other variations of ELISAs known in the art. For further discussion regarding ELISAs see, e.g., Ausubel et al, eds, (1994), Current Protocols in Molecular Biology, Vol. 1, John Wiley & Sons, Inc., New York, section 11.2.1.

[349] The binding affinity of an antibody to an antigen and the off-rate of an antibody-antigen interaction can be determined by competitive binding assays. One example of a competitive binding assay is a radioimmunoassay comprising the incubation of labeled antigen (e.g., ^3H or ^{125}I) with the antibody of interest in the presence of increasing amounts of unlabeled antigen, and the detection of the antibody bound to the labeled antigen. The affinity of the antibody of interest for a particular antigen and the binding off-rates can be determined from the data by scatchard plot analysis. Competition with a second antibody can also be determined using radioimmunoassays. In this case, the antigen is incubated with antibody of interest conjugated to a labeled compound (e.g., ^3H or ^{125}I) in the presence of increasing amounts of an unlabeled second antibody.

[350] Antibodies of the invention may be characterized using immunocytochemistry methods on cells (e.g., mammalian cells, such as CHO cells) transfected with a vector enabling the expression of an antigen or with vector alone using techniques commonly known in the art. Antibodies that bind antigen transfected cells, but not vector-only transfected cells, are antigen specific.

Therapeutic Uses

[351] Table 1D: In preferred embodiments, the present invention encompasses a method of treating a disease or disorder listed in the "Preferred Indications" column of Table 1D; comprising administering to a patient in which such treatment, prevention, or amelioration is desired a protein, nucleic acid, or antibody of the invention (or fragment or variant thereof) represented by Table 1A and Table 1D (in the same row as the disease or disorder to be treated is listed in the "Preferred Indications" column of Table 1D) in an amount effective to treat, prevent, or ameliorate the disease or disorder.

[352] As indicated in Table 1D, the polynucleotides, polypeptides, agonists, or antagonists of the present invention (including antibodies) can be used in assays to test for one or more biological activities. If these polynucleotides and polypeptides do exhibit activity in a particular assay, it is likely that these molecules may be involved in the diseases associated with the biological activity. Thus, the polynucleotides or polypeptides, or agonists or antagonists thereof (including antibodies) could be used to treat the associated disease.

[353] The present invention encompasses methods of preventing, treating, diagnosing, or ameliorating a disease or disorder. In preferred embodiments, the present invention encompasses a method of treating a disease or disorder listed in the "Preferred Indications" column of Table 1D; comprising administering to a patient in which such treatment, prevention, or amelioration is desired a protein, nucleic acid, or antibody of the invention (or fragment or variant thereof) in an amount effective to treat, prevent, diagnose, or ameliorate the disease or disorder. The first and second columns of Table 1D show the "Gene No." and "cDNA Clone ID No.", respectively, indicating certain nucleic acids and proteins (or antibodies against the same) of the invention (including polynucleotide, polypeptide, and antibody fragments or variants thereof) that may be used in preventing, treating, diagnosing, or ameliorating the disease(s) or disorder(s) indicated in the corresponding row in Column 3 of Table 1D.

[354] In another embodiment, the present invention also encompasses methods of preventing, treating, diagnosing, or ameliorating a disease or disorder listed in the "Preferred Indications" column of Table 1D; comprising administering to a patient combinations of the proteins, nucleic acids, or antibodies of the invention (or fragments or variants thereof), sharing similar indications as shown in the corresponding rows in Column 3 of Table 1D.

[355] The "Preferred Indication" column describes diseases, disorders, and/or conditions that may be treated, prevented, diagnosed, or ameliorated by a protein, nucleic acid, or antibody of the invention (or fragment or variant thereof).

[356] The recitation of "Cancer" in the "Preferred Indication" column indicates that the corresponding nucleic acid and protein, or antibody against the same, of the invention (or fragment or variant thereof) may be used for example, to diagnose, treat, prevent, and/or ameliorate diseases and/or disorders relating to neoplastic diseases (e.g., leukemias, cancers, and/or as described below under "Hyperproliferative Disorders").

[357] In specific embodiments, a protein, nucleic acid, or antibody of the invention (or fragment or variant thereof) having a “Cancer” recitation in the “Preferred Indication” column of Table 1D may be used for example, to diagnose, treat, prevent, and/or ameliorate a neoplasm located in a tissue selected from the group consisting of: colon, abdomen, bone, breast, digestive system, liver, pancreas, prostate, peritoneum, lung, blood (e.g., leukemia), endocrine glands (adrenal, parathyroid, pituitary, testicles, ovary, thymus, thyroid), uterus, eye, head and neck, nervous (central and peripheral), lymphatic system, pelvic, skin, soft tissue, spleen, thoracic, and urogenital.

[358] In specific embodiments, a protein, nucleic acid, or antibody of the invention (or fragment or variant thereof) having a “Cancer” recitation in the “Preferred Indication” column of Table 1D, may be used for example, to diagnose, treat, prevent, and/or ameliorate a pre-neoplastic condition, selected from the group consisting of: hyperplasia (e.g., endometrial hyperplasia and/or as described in the section entitled “Hyperproliferative Disorders”), metaplasia (e.g., connective tissue metaplasia, atypical metaplasia, and/or as described in the section entitled “Hyperproliferative Disorders”), and/or dysplasia (e.g., cervical dysplasia, and bronchopulmonary dysplasia).

[359] In another specific embodiment, a protein, nucleic acid, or antibody of the invention (or fragment or variant thereof) having a “Cancer” recitation in the “Preferred Indication” column of Table 1D, may be used for example, to diagnose, treat, prevent, and/or ameliorate a benign dysproliferative disorder selected from the group consisting of: benign tumors, fibrocystic conditions, tissue hypertrophy, and/or as described in the section entitled “Hyperproliferative Disorders”.

[360] The recitation of “Immune/Hematopoietic” in the “Preferred Indication” column indicates that the corresponding nucleic acid and protein, or antibody against the same, of the invention (or fragment or variant thereof), may be used for example, to diagnose, treat, prevent, and/or ameliorate diseases and/or disorders relating to neoplastic diseases (e.g., as described below under “Hyperproliferative Disorders”), blood disorders (e.g., as described below under “Immune Activity” “Cardiovascular Disorders” and/or “Blood-Related Disorders”), and infections (e.g., as described below under “Infectious Disease”).

[361] In specific embodiments, a protein, nucleic acid, or antibody of the invention (or fragment or variant thereof) having the “Immune/Hematopoietic” recitation in the “Preferred Indication” column of Table 1D, may be used for example, to diagnose, treat, prevent, and/or ameliorate a disease or disorder selected from the group consisting of: anemia, pancytopenia, leukopenia, thrombocytopenia, leukemias, Hodgkin’s disease, non-

Hodgkin's lymphoma, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, asthma, AIDS, autoimmune disease, rheumatoid arthritis, granulomatous disease, immune deficiency, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, immune reactions to transplanted organs and tissues, systemic lupus erythematosus, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, and allergies.

[362] The recitation of "Reproductive" in the "Preferred Indication" column indicates that the corresponding nucleic acid and protein, or antibody against the same, of the invention (or fragment or variant thereof), may be used for example, to diagnose, treat, prevent, and/or ameliorate diseases and/or disorders relating to neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), and disorders of the reproductive system (e.g., as described below under "Reproductive System Disorders").

[363] In specific embodiments, a protein, nucleic acid, or antibody of the invention (or fragment or variant thereof) having a "Reproductive" recitation in the "Preferred Indication" column of Table 1D, may be used for example, to diagnose, treat, prevent, and/or ameliorate a disease or disorder selected from the group consisting of: cryptorchism, prostatitis, inguinal hernia, varicocele, leydig cell tumors, verrucous carcinoma, prostatitis, malacoplakia, Peyronie's disease, penile carcinoma, squamous cell hyperplasia, dysmenorrhea, ovarian adenocarcinoma, Turner's syndrome, mucopurulent cervicitis, Sertoli-leydig tumors, ovarian cancer, uterine cancer, pelvic inflammatory disease, testicular cancer, prostate cancer, Klinefelter's syndrome, Young's syndrome, premature ejaculation, diabetes mellitus, cystic fibrosis, Kartagener's syndrome, testicular atrophy, testicular feminization, anorchia, ectopic testis, epididymitis, orchitis, gonorrhea, syphilis, testicular torsion, vasitis nodosa, germ cell tumors, stromal tumors, dysmenorrhea, retroverted uterus, endometriosis, fibroids, adenomyosis, anovulatory bleeding, amenorrhea, Cushing's syndrome, hydatidiform moles, Asherman's syndrome, premature menopause, precocious puberty, uterine polyps, dysfunctional uterine bleeding, cervicitis, chronic cervicitis, mucopurulent cervicitis, cervical dysplasia, cervical polyps, Nabothian cysts, cervical erosion, cervical incompetence, cervical neoplasms, pseudohermaphroditism, and premenstrual syndrome.

[364] The recitation of "Musculoskeletal" in the "Preferred Indication" column indicates that the corresponding nucleic acid and protein, or antibody against the same, of the invention (or fragment or variant thereof), may be used for example, to diagnose, treat, prevent, and/or ameliorate diseases and/or disorders relating to neoplastic diseases (e.g., as

described below under “Hyperproliferative Disorders”), and disorders of the immune system (e.g., as described below under “Immune Activity”).

[365] In specific embodiments, a protein, nucleic acid, or antibody of the invention (or fragment or variant thereof) having a “Musculoskeletal” recitation in the “Preferred Indication” column of Table 1D, may be used for example, to diagnose, treat, prevent, and/or ameliorate a disease or disorder selected from the group consisting of: bone cancers (e.g., osteochondromas, benign chondromas, chondroblastoma, chondromyxoid fibromas, osteoid osteomas, giant cell tumors, multiple myeloma, osteosarcomas), Paget’s Disease, rheumatoid arthritis, systemic lupus erythematosus, osteomyelitis, Lyme Disease, gout, bursitis, tendonitis, osteoporosis, osteoarthritis, muscular dystrophy, mitochondrial myopathy, cachexia, and multiple sclerosis.

[366] The recitation of “Cardiovascular” in the “Preferred Indication” column indicates that the corresponding nucleic acid and protein, or antibody against the same, of the invention (or fragment or variant thereof), may be used for example, to diagnose, treat, prevent, and/or ameliorate diseases and/or disorders relating to neoplastic diseases (e.g., as described below under “Hyperproliferative Disorders”), and disorders of the cardiovascular system (e.g., as described below under “Cardiovascular Disorders”).

[367] In specific embodiments, a protein, nucleic acid, or antibody of the invention (or fragment or variant thereof) having a “Cardiovascular” recitation in the “Preferred Indication” column of Table 1D, may be used for example, to diagnose, treat, prevent, and/or ameliorate a disease or disorder selected from the group consisting of: myxomas, fibromas, rhabdomyomas, cardiovascular abnormalities (e.g., congenital heart defects, cerebral arteriovenous malformations, septal defects), heart disease (e.g., heart failure, congestive heart disease, arrhythmia, tachycardia, fibrillation, pericardial Disease, endocarditis), cardiac arrest, heart valve disease (e.g., stenosis, regurgitation, prolapse), vascular disease (e.g., hypertension, coronary artery disease, angina, aneurysm, arteriosclerosis, peripheral vascular disease), hyponatremia, hypernatremia, hypokalemia, and hyperkalemia.

[368] The recitation of “Mixed Fetal” in the “Preferred Indication” column indicates that the corresponding nucleic acid and protein, or antibody against the same, of the invention (or fragment or variant thereof), may be used for example, to diagnose, treat, prevent, and/or ameliorate diseases and/or disorders relating to neoplastic diseases (e.g., as described below under “Hyperproliferative Disorders”).

[369] In specific embodiments, a protein, nucleic acid, or antibody of the invention (or fragment or variant thereof) having a “Mixed Fetal” recitation in the “Preferred Indication” column of Table 1D, may be used for example, to diagnose, treat, prevent, and/or ameliorate a disease or disorder selected from the group consisting of: spina bifida, hydranencephaly, neurofibromatosis, fetal alcohol syndrome, diabetes mellitus, PKU, Down’s syndrome, Patau syndrome, Edwards syndrome, Turner syndrome, Apert syndrome, Carpenter syndrome, Conradi syndrome, Crouzon syndrome, cutis laxa, Cornelia de Lange syndrome, Ellis-van Creveld syndrome, Holt-Oram syndrome, Kartagener syndrome, Meckel-Gruber syndrome, Noonan syndrome, Pallister-Hall syndrome, Rubinstein-Taybi syndrome, Scimitar syndrome, Smith-Lemli-Opitz syndrome, thrombocytopenia-absent radius (TAR) syndrome, Treacher Collins syndrome, Williams syndrome, Hirschsprung’s disease, Meckel’s diverticulum, polycystic kidney disease, Turner’s syndrome, and gonadal dysgenesis, Klippel-Feil syndrome, Osteogenesis imperfecta, muscular dystrophy, Tay-Sachs disease, Wilm’s tumor, neuroblastoma, and retinoblastoma.

[370] The recitation of “Excretory” in the “Preferred Indication” column indicates that the corresponding nucleic acid and protein, or antibody against the same, of the invention (or fragment or variant thereof), may be used for example, to diagnose, treat, prevent, and/or ameliorate diseases and/or disorders relating to neoplastic diseases (e.g., as described below under “Hyperproliferative Disorders”) and renal disorders (e.g., as described below under “Renal Disorders”).

[371] In specific embodiments, a protein, nucleic acid, or antibody of the invention (or fragment or variant thereof) having a “Excretory” recitation in the “Preferred Indication” column of Table 1D, may be used for example, to diagnose, treat, prevent, and/or ameliorate a disease or disorder selected from the group consisting of: bladder cancer, prostate cancer, benign prostatic hyperplasia, bladder disorders (e.g., urinary incontinence, urinary retention, urinary obstruction, urinary tract Infections, interstitial cystitis, prostatitis, neurogenic bladder, hematuria), renal disorders (e.g., hydronephrosis, proteinuria, renal failure, pyelonephritis, urolithiasis, reflux nephropathy, and unilateral obstructive uropathy).

[372] The recitation of “Neural/Sensory” in the “Preferred Indication” column indicates that the corresponding nucleic acid and protein, or antibody against the same, of the invention (or fragment or variant thereof), may be used for example, to diagnose, treat, prevent, and/or ameliorate diseases and/or disorders relating to neoplastic diseases (e.g., as

described below under “Hyperproliferative Disorders”) and diseases or disorders of the nervous system (e.g., as described below under “Neural Activity and Neurological Diseases”).

[373] In specific embodiments, a protein, nucleic acid, or antibody of the invention (or fragment or variant thereof) having a “Neural/Sensory” recitation in the “Preferred Indication” column of Table 1D, may be used for example, to diagnose, treat, prevent, and/or ameliorate a disease or disorder selected from the group consisting of: brain cancer (e.g., brain stem glioma, brain tumors, central nervous system (Primary) lymphoma, central nervous system lymphoma, cerebellar astrocytoma, and cerebral astrocytoma, neurodegenerative disorders (e.g., Alzheimer's Disease, Creutzfeldt-Jakob Disease, Parkinson's Disease, and Idiopathic Presenile Dementia), encephalomyelitis, cerebral malaria, meningitis, metabolic brain diseases (e.g., phenylketonuria and pyruvate carboxylase deficiency), cerebellar ataxia, ataxia telangiectasia, and AIDS Dementia Complex, schizophrenia, attention deficit disorder, hyperactive attention deficit disorder, autism, and obsessive compulsive disorders.

[374] The recitation of “Respiratory” in the “Preferred Indication” column indicates that the corresponding nucleic acid and protein, or antibody against the same, of the invention (or fragment or variant thereof), may be used for example, to diagnose, treat, prevent, and/or ameliorate diseases and/or disorders relating to neoplastic diseases (e.g., as described below under “Hyperproliferative Disorders”) and diseases or disorders of the respiratory system (e.g., as described below under “Respiratory Disorders”).

[375] In specific embodiments, a protein, nucleic acid, or antibody of the invention (or fragment or variant thereof) having a “Respiratory” recitation in the “Preferred Indication” column of Table 1D, may be used for example, to diagnose, treat, prevent, and/or ameliorate a disease or disorder selected from the group consisting of: cancers of the respiratory system such as larynx cancer, pharynx cancer, trachea cancer, epiglottis cancer, lung cancer, squamous cell carcinomas, small cell (oat cell) carcinomas, large cell carcinomas, and adenocarcinomas. Allergic reactions, cystic fibrosis, sarcoidosis, histiocytosis X, infiltrative lung diseases (e.g., pulmonary fibrosis and lymphoid interstitial pneumonia), obstructive airway diseases (e.g., asthma, emphysema, chronic or acute bronchitis), occupational lung diseases (e.g., silicosis and asbestosis), pneumonia, and pleurisy.

[376] The recitation of “Endocrine” in the “Preferred Indication” column indicates that the corresponding nucleic acid and protein, or antibody against the same, of the invention

0950083-091201

(or fragment or variant thereof), may be used for example, to diagnose, treat, prevent, and/or ameliorate diseases and/or disorders relating to neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders") and diseases or disorders of the respiratory system (e.g., as described below under "Respiratory Disorders"), renal disorders (e.g., as described below under "Renal Disorders"), and disorders of the endocrine system (e.g., as described below under "Endocrine Disorders").

[377] In specific embodiments, a protein, nucleic acid, or antibody of the invention (or fragment or variant thereof) having an "Endocrine" recitation in the "Preferred Indication" column of Table 1D, may be used for example, to diagnose, treat, prevent, and/or ameliorate a disease or disorder selected from the group consisting of: cancers of endocrine tissues and organs (e.g., cancers of the hypothalamus, pituitary gland, thyroid gland, parathyroid glands, pancreas, adrenal glands, ovaries, and testes), diabetes (e.g., diabetes insipidus, type I and type II diabetes mellitus), obesity, disorders related to pituitary glands (e.g., hyperpituitarism, hypopituitarism, and pituitary dwarfism), hypothyroidism, hyperthyroidism, goiter, reproductive disorders (e.g. male and female infertility), disorders related to adrenal glands (e.g., Addison's Disease, corticosteroid deficiency, and Cushing's Syndrome), kidney cancer (e.g., hypernephroma, transitional cell cancer, and Wilm's tumor), diabetic nephropathy, interstitial nephritis, polycystic kidney disease, glomerulonephritis (e.g., IgM mesangial proliferative glomerulonephritis and glomerulonephritis caused by autoimmune disorders; such as Goodpasture's syndrome), and nephrocalcinosis.

[378] The recitation of "Digestive" in the "Preferred Indication" column indicates that the corresponding nucleic acid and protein, or antibody against the same, of the invention (or fragment or variant thereof), may be used for example, to diagnose, treat, prevent, and/or ameliorate diseases and/or disorders relating to neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders") and diseases or disorders of the gastrointestinal system (e.g., as described below under "Gastrointestinal Disorders").

[379] In specific embodiments, a protein, nucleic acid, or antibody of the invention (or fragment or variant thereof) having a "Digestive" recitation in the "Preferred Indication" column of Table 1D, may be used for example, to diagnose, treat, prevent, and/or ameliorate a disease or disorder selected from the group consisting of: ulcerative colitis, appendicitis, Crohn's disease, hepatitis, hepatic encephalopathy, portal hypertension, cholelithiasis, cancer of the digestive system (e.g., biliary tract cancer, stomach cancer, colon cancer, gastric cancer, pancreatic cancer, cancer of the bile duct, tumors of the colon

(e.g., polyps or cancers), and cirrhosis), pancreatitis, ulcerative disease, pyloric stenosis, gastroenteritis, gastritis, gastric atrophy, benign tumors of the duodenum, distension, irritable bowel syndrome, malabsorption, congenital disorders of the small intestine, bacterial and parasitic infection, megacolon, Hirschsprung's disease, aganglionic megacolon, acquired megacolon, colitis, anorectal disorders (e.g., anal fistulas, hemorrhoids), congenital disorders of the liver (e.g., Wilson's disease, hemochromatosis, cystic fibrosis, biliary atresia, and alpha1-antitrypsin deficiency), portal hypertension, cholelithiasis, and jaundice.

[380] The recitation of "Connective/Epithelial" in the "Preferred Indication" column indicates that the corresponding nucleic acid and protein, or antibody against the same, of the invention (or fragment or variant thereof), may be used for example, to diagnose, treat, prevent, and/or ameliorate diseases and/or disorders relating to neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), cellular and genetic abnormalities (e.g., as described below under "Diseases at the Cellular Level"), angiogenesis (e.g., as described below under "Anti-Angiogenesis Activity"), and or to promote or inhibit regeneration (e.g., as described below under "Regeneration"), and wound healing (e.g., as described below under "Wound Healing and Epithelial Cell Proliferation").

In specific embodiments, a protein, nucleic acid, or antibody of the invention (or fragment or variant thereof) having a "Connective/Epithelial" recitation in the "Preferred Indication" column of Table 1D, may be used for example, to diagnose, treat, prevent, and/or ameliorate a disease or disorder selected from the group consisting of: connective tissue metaplasia, mixed connective tissue disease, focal epithelial hyperplasia, epithelial metaplasia, mucoepithelial dysplasia, graft v. host disease, polymyositis, cystic hyperplasia, cerebral dysplasia, tissue hypertrophy, Alzheimer's disease, lymphoproliferative disorder, Waldenstrom's macroglobulinemia, Crohn's disease, pernicious anemia, idiopathic Addison's disease, glomerulonephritis, bullous pemphigoid, Sjogren's syndrome, diabetes mellitus, cystic fibrosis, osteoblastoma, osteoclastoma, osteosarcoma, chondrosarcoma, osteoporosis, osteoarthritis, periodontal disease, wound healing, relapsing polychondritis, vasculitis, polyarteritis nodosa, Wegener's granulomatosis, cellulitis, rheumatoid arthritis, psoriatic arthritis, discoid lupus erythematosus, systemic lupus erythematosus, scleroderma, CREST syndrome, Sjogren's syndrome, polymyositis, dermatomyositis, mixed connective tissue disease, relapsing polychondritis, vasculitis, Henoch-Schonlein syndrome, erythema nodosum, polyarteritis nodosa, temporal (giant cell) arteritis, Takayasu's arteritis, Wegener's granulomatosis, Reiter's syndrome, Behcet's syndrome, ankylosing spondylitis, cellulitis, keloids, Ehler Danlos syndrome, Marfan syndrome, pseudoxantoma elasticum,

osteogenese imperfecta, chondrodysplasias, epidermolysis bullosa, Alport syndrome, and cutis laxa.

[381] The present invention is further directed to antibody-based therapies which involve administering antibodies of the invention to an animal, preferably a mammal, and most preferably a human, patient for treating one or more of the disclosed diseases, disorders, or conditions. Therapeutic compounds of the invention include, but are not limited to, antibodies of the invention (including fragments, analogs and derivatives thereof as described herein) and nucleic acids encoding antibodies of the invention (including fragments, analogs and derivatives thereof and anti-idiotypic antibodies as described herein). The antibodies of the invention can be used to treat, inhibit or prevent diseases, disorders or conditions associated with aberrant expression and/or activity of a polypeptide of the invention, including, but not limited to, any one or more of the diseases, disorders, or conditions described herein. The treatment and/or prevention of diseases, disorders, or conditions associated with aberrant expression and/or activity of a polypeptide of the invention includes, but is not limited to, alleviating symptoms associated with those diseases, disorders or conditions. Antibodies of the invention may be provided in pharmaceutically acceptable compositions as known in the art or as described herein.

[382] In a specific and preferred embodiment, the present invention is directed to antibody-based therapies which involve administering antibodies of the invention to an animal, preferably a mammal, and most preferably a human, patient for treating one or more diseases, disorders, or conditions, including but not limited to: neural disorders, immune system disorders, muscular disorders, reproductive disorders, gastrointestinal disorders, pulmonary disorders, cardiovascular disorders, renal disorders, proliferative disorders, and/or cancerous diseases and conditions., and/or as described elsewhere herein. Therapeutic compounds of the invention include, but are not limited to, antibodies of the invention (e.g., antibodies directed to the full length protein expressed on the cell surface of a mammalian cell; antibodies directed to an epitope of a polypeptide of the invention (such as, for example, a predicted linear epitope shown in column 7 of Table 1B; or a conformational epitope, including fragments, analogs and derivatives thereof as described herein) and nucleic acids encoding antibodies of the invention (including fragments, analogs and derivatives thereof and anti-idiotypic antibodies as described herein). The antibodies of the invention can be used to treat, inhibit or prevent diseases, disorders or conditions associated with aberrant expression and/or activity of a polypeptide of the

invention, including, but not limited to, any one or more of the diseases, disorders, or conditions described herein. The treatment and/or prevention of diseases, disorders, or conditions associated with aberrant expression and/or activity of a polypeptide of the invention includes, but is not limited to, alleviating symptoms associated with those diseases, disorders or conditions. Antibodies of the invention may be provided in pharmaceutically acceptable compositions as known in the art or as described herein.

[383] A summary of the ways in which the antibodies of the present invention may be used therapeutically includes binding polynucleotides or polypeptides of the present invention locally or systemically in the body or by direct cytotoxicity of the antibody, e.g. as mediated by complement (CDC) or by effector cells (ADCC). Some of these approaches are described in more detail below. Armed with the teachings provided herein, one of ordinary skill in the art will know how to use the antibodies of the present invention for diagnostic, monitoring or therapeutic purposes without undue experimentation.

[384] The antibodies of this invention may be advantageously utilized in combination with other monoclonal or chimeric antibodies, or with lymphokines or hematopoietic growth factors (such as, e.g., IL-2, IL-3 and IL-7), for example, which serve to increase the number or activity of effector cells which interact with the antibodies.

[385] The antibodies of the invention may be administered alone or in combination with other types of treatments (e.g., radiation therapy, chemotherapy, hormonal therapy, immunotherapy and anti-tumor agents). Generally, administration of products of a species origin or species reactivity (in the case of antibodies) that is the same species as that of the patient is preferred. Thus, in a preferred embodiment, human antibodies, fragments derivatives, analogs, or nucleic acids, are administered to a human patient for therapy or prophylaxis.

[386] It is preferred to use high affinity and/or potent *in vivo* inhibiting and/or neutralizing antibodies against polypeptides or polynucleotides of the present invention, fragments or regions thereof, for both immunoassays directed to and therapy of disorders related to polynucleotides or polypeptides, including fragments thereof, of the present invention. Such antibodies, fragments, or regions, will preferably have an affinity for polynucleotides or polypeptides of the invention, including fragments thereof. Preferred binding affinities include those with a dissociation constant or K_d less than 5×10^{-2} M, 10^{-2} M, 5×10^{-3} M, 10^{-3} M, 5×10^{-4} M, 10^{-4} M, 5×10^{-5} M, 10^{-5} M, 5×10^{-6} M, 10^{-6} M, 5×10^{-7} M, 10^{-7} M, 5×10^{-8} M, 10^{-8} M, 5×10^{-9} M, 10^{-9} M, 5×10^{-10} M, 10^{-10} M, 5×10^{-11} M,

10^{-11} M, 5×10^{-12} M, 10^{-12} M, 5×10^{-13} M, 10^{-13} M, 5×10^{-14} M, 10^{-14} M, 5×10^{-15} M, and 10^{-15} M.

Gene Therapy

[387] In a specific embodiment, nucleic acids comprising sequences encoding antibodies or functional derivatives thereof, are administered to treat, inhibit or prevent a disease or disorder associated with aberrant expression and/or activity of a polypeptide of the invention, by way of gene therapy. Gene therapy refers to therapy performed by the administration to a subject of an expressed or expressible nucleic acid. In this embodiment of the invention, the nucleic acids produce their encoded protein that mediates a therapeutic effect.

[388] Any of the methods for gene therapy available in the art can be used according to the present invention. Exemplary methods are described below.

[389] For general reviews of the methods of gene therapy, see Goldspiel et al., *Clinical Pharmacy* 12:488-505 (1993); Wu and Wu, *Biotherapy* 3:87-95 (1991); Tolstoshev, *Ann. Rev. Pharmacol. Toxicol.* 32:573-596 (1993); Mulligan, *Science* 260:926-932 (1993); and Morgan and Anderson, *Ann. Rev. Biochem.* 62:191-217 (1993); May, *TIBTECH* 11(5):155-215 (1993). Methods commonly known in the art of recombinant DNA technology which can be used are described in Ausubel et al. (eds.), *Current Protocols in Molecular Biology*, John Wiley & Sons, NY (1993); and Kriegler, *Gene Transfer and Expression, A Laboratory Manual*, Stockton Press, NY (1990).

[390] In a preferred embodiment, the compound comprises nucleic acid sequences encoding an antibody, said nucleic acid sequences being part of expression vectors that express the antibody or fragments or chimeric proteins or heavy or light chains thereof in a suitable host. In particular, such nucleic acid sequences have promoters operably linked to the antibody coding region, said promoter being inducible or constitutive, and, optionally, tissue-specific. In another particular embodiment, nucleic acid molecules are used in which the antibody coding sequences and any other desired sequences are flanked by regions that promote homologous recombination at a desired site in the genome, thus providing for intrachromosomal expression of the antibody encoding nucleic acids (Koller and Smithies, *Proc. Natl. Acad. Sci. USA* 86:8932-8935 (1989); Zijlstra et al., *Nature* 342:435-438 (1989). In specific embodiments, the expressed antibody molecule is a single chain

09950083-091201

antibody; alternatively, the nucleic acid sequences include sequences encoding both the heavy and light chains, or fragments thereof, of the antibody.

[391] Delivery of the nucleic acids into a patient may be either direct, in which case the patient is directly exposed to the nucleic acid or nucleic acid- carrying vectors, or indirect, in which case, cells are first transformed with the nucleic acids *in vitro*, then transplanted into the patient. These two approaches are known, respectively, as *in vivo* or *ex vivo* gene therapy.

[392] In a specific embodiment, the nucleic acid sequences are directly administered *in vivo*, where it is expressed to produce the encoded product. This can be accomplished by any of numerous methods known in the art, e.g., by constructing them as part of an appropriate nucleic acid expression vector and administering it so that they become intracellular, e.g., by infection using defective or attenuated retrovirals or other viral vectors (see U.S. Patent No. 4,980,286), or by direct injection of naked DNA, or by use of microparticle bombardment (e.g., a gene gun; Biolistic, Dupont), or coating with lipids or cell-surface receptors or transfecting agents, encapsulation in liposomes, microparticles, or microcapsules, or by administering them in linkage to a peptide which is known to enter the nucleus, by administering it in linkage to a ligand subject to receptor-mediated endocytosis (see, e.g., Wu and Wu, J. Biol. Chem. 262:4429-4432 (1987)) (which can be used to target cell types specifically expressing the receptors), etc. In another embodiment, nucleic acid-ligand complexes can be formed in which the ligand comprises a fusogenic viral peptide to disrupt endosomes, allowing the nucleic acid to avoid lysosomal degradation. In yet another embodiment, the nucleic acid can be targeted *in vivo* for cell specific uptake and expression, by targeting a specific receptor (see, e.g., PCT Publications WO 92/06180; WO 92/22635; WO92/20316; WO93/14188, WO 93/20221). Alternatively, the nucleic acid can be introduced intracellularly and incorporated within host cell DNA for expression, by homologous recombination (Koller and Smithies, Proc. Natl. Acad. Sci. USA 86:8932-8935 (1989); Zijlstra et al., Nature 342:435-438 (1989)).

[393] In a specific embodiment, viral vectors that contains nucleic acid sequences encoding an antibody of the invention are used. For example, a retroviral vector can be used (see Miller et al., Meth. Enzymol. 217:581-599 (1993)). These retroviral vectors contain the components necessary for the correct packaging of the viral genome and integration into the host cell DNA. The nucleic acid sequences encoding the antibody to be used in gene therapy are cloned into one or more vectors, which facilitates delivery of the

gene into a patient. More detail about retroviral vectors can be found in Boesen et al., Biotherapy 6:291-302 (1994), which describes the use of a retroviral vector to deliver the *mdr1* gene to hematopoietic stem cells in order to make the stem cells more resistant to chemotherapy. Other references illustrating the use of retroviral vectors in gene therapy are: Clowes et al., J. Clin. Invest. 93:644-651 (1994); Kiem et al., Blood 83:1467-1473 (1994); Salmons and Gunzberg, Human Gene Therapy 4:129-141 (1993); and Grossman and Wilson, Curr. Opin. in Genetics and Devel. 3:110-114 (1993).

[394] Adenoviruses are other viral vectors that can be used in gene therapy. Adenoviruses are especially attractive vehicles for delivering genes to respiratory epithelia. Adenoviruses naturally infect respiratory epithelia where they cause a mild disease. Other targets for adenovirus-based delivery systems are liver, the central nervous system, endothelial cells, and muscle. Adenoviruses have the advantage of being capable of infecting non-dividing cells. Kozarsky and Wilson, Current Opinion in Genetics and Development 3:499-503 (1993) present a review of adenovirus-based gene therapy. Bout et al., Human Gene Therapy 5:3-10 (1994) demonstrated the use of adenovirus vectors to transfer genes to the respiratory epithelia of rhesus monkeys. Other instances of the use of adenoviruses in gene therapy can be found in Rosenfeld et al., Science 252:431-434 (1991); Rosenfeld et al., Cell 68:143- 155 (1992); Mastrangeli et al., J. Clin. Invest. 91:225-234 (1993); PCT Publication WO94/12649; and Wang, et al., Gene Therapy 2:775-783 (1995). In a preferred embodiment, adenovirus vectors are used.

[395] Adeno-associated virus (AAV) has also been proposed for use in gene therapy (Walsh et al., Proc. Soc. Exp. Biol. Med. 204:289-300 (1993); U.S. Patent No. 5,436,146).

[396] Another approach to gene therapy involves transferring a gene to cells in tissue culture by such methods as electroporation, lipofection, calcium phosphate mediated transfection, or viral infection. Usually, the method of transfer includes the transfer of a selectable marker to the cells. The cells are then placed under selection to isolate those cells that have taken up and are expressing the transferred gene. Those cells are then delivered to a patient.

[397] In this embodiment, the nucleic acid is introduced into a cell prior to administration *in vivo* of the resulting recombinant cell. Such introduction can be carried out by any method known in the art, including but not limited to transfection, electroporation, microinjection, infection with a viral or bacteriophage vector containing the nucleic acid sequences, cell fusion, chromosome-mediated gene transfer, microcell-

mediated gene transfer, spheroplast fusion, etc. Numerous techniques are known in the art for the introduction of foreign genes into cells (see, e.g., Loeffler and Behr, Meth. Enzymol. 217:599-618 (1993); Cohen et al., Meth. Enzymol. 217:618-644 (1993); Cline, Pharmac. Ther. 29:69-92m (1985) and may be used in accordance with the present invention, provided that the necessary developmental and physiological functions of the recipient cells are not disrupted. The technique should provide for the stable transfer of the nucleic acid to the cell, so that the nucleic acid is expressible by the cell and preferably heritable and expressible by its cell progeny.

[398] The resulting recombinant cells can be delivered to a patient by various methods known in the art. Recombinant blood cells (e.g., hematopoietic stem or progenitor cells) are preferably administered intravenously. The amount of cells envisioned for use depends on the desired effect, patient state, etc., and can be determined by one skilled in the art.

[399] Cells into which a nucleic acid can be introduced for purposes of gene therapy encompass any desired, available cell type, and include but are not limited to epithelial cells, endothelial cells, keratinocytes, fibroblasts, muscle cells, hepatocytes; blood cells such as T lymphocytes, B lymphocytes, monocytes, macrophages, neutrophils, eosinophils, megakaryocytes, granulocytes; various stem or progenitor cells, in particular hematopoietic stem or progenitor cells, e.g., as obtained from bone marrow, umbilical cord blood, peripheral blood, fetal liver, etc.

[400] In a preferred embodiment, the cell used for gene therapy is autologous to the patient.

[401] In an embodiment in which recombinant cells are used in gene therapy, nucleic acid sequences encoding an antibody are introduced into the cells such that they are expressible by the cells or their progeny, and the recombinant cells are then administered *in vivo* for therapeutic effect. In a specific embodiment, stem or progenitor cells are used. Any stem and/or progenitor cells which can be isolated and maintained *in vitro* can potentially be used in accordance with this embodiment of the present invention (see e.g. PCT Publication WO 94/08598; Stemple and Anderson, Cell 71:973-985 (1992); Rheinwald, Meth. Cell Bio. 21A:229 (1980); and Pittelkow and Scott, Mayo Clinic Proc. 61:771 (1986)).

[402] In a specific embodiment, the nucleic acid to be introduced for purposes of gene therapy comprises an inducible promoter operably linked to the coding region, such that

0950083-091201

expression of the nucleic acid is controllable by the presence or absence of an appropriate inducer of transcription.

Demonstration of Therapeutic or Prophylactic Activity

[403] The compounds or pharmaceutical compositions of the invention are preferably tested *in vitro*, and then *in vivo* for the desired therapeutic or prophylactic activity, prior to use in humans. For example, *in vitro* assays to demonstrate the therapeutic or prophylactic utility of a compound or pharmaceutical composition include, the effect of a compound on a cell line or a patient tissue sample. The effect of the compound or composition on the cell line and/or tissue sample can be determined utilizing techniques known to those of skill in the art including, but not limited to, rosette formation assays and cell lysis assays. In accordance with the invention, *in vitro* assays which can be used to determine whether administration of a specific compound is indicated, include *in vitro* cell culture assays in which a patient tissue sample is grown in culture, and exposed to or otherwise administered a compound, and the effect of such compound upon the tissue sample is observed.

Therapeutic/Prophylactic Administration and Composition

[404] The invention provides methods of treatment, inhibition and prophylaxis by administration to a subject of an effective amount of a compound or pharmaceutical composition of the invention, preferably a polypeptide or antibody of the invention. In a preferred embodiment, the compound is substantially purified (e.g., substantially free from substances that limit its effect or produce undesired side-effects). The subject is preferably an animal, including but not limited to animals such as cows, pigs, horses, chickens, cats, dogs, etc., and is preferably a mammal, and most preferably human.

[405] Formulations and methods of administration that can be employed when the compound comprises a nucleic acid or an immunoglobulin are described above; additional appropriate formulations and routes of administration can be selected from among those described herein below.

[406] Various delivery systems are known and can be used to administer a compound of the invention, e.g., encapsulation in liposomes, microparticles, microcapsules, recombinant cells capable of expressing the compound, receptor-mediated endocytosis (see, e.g., Wu and Wu, J. Biol. Chem. 262:4429-4432 (1987)), construction of a nucleic acid as part of a retroviral or other vector, etc. Methods of introduction include but are not limited

09950087, 091201

to intradermal, intramuscular, intraperitoneal, intravenous, subcutaneous, intranasal, epidural, and oral routes. The compounds or compositions may be administered by any convenient route, for example by infusion or bolus injection, by absorption through epithelial or mucocutaneous linings (e.g., oral mucosa, rectal and intestinal mucosa, etc.) and may be administered together with other biologically active agents. Administration can be systemic or local. In addition, it may be desirable to introduce the pharmaceutical compounds or compositions of the invention into the central nervous system by any suitable route, including intraventricular and intrathecal injection; intraventricular injection may be facilitated by an intraventricular catheter, for example, attached to a reservoir, such as an Ommaya reservoir. Pulmonary administration can also be employed, e.g., by use of an inhaler or nebulizer, and formulation with an aerosolizing agent.

[407] In a specific embodiment, it may be desirable to administer the pharmaceutical compounds or compositions of the invention locally to the area in need of treatment; this may be achieved by, for example, and not by way of limitation, local infusion during surgery, topical application, e.g., in conjunction with a wound dressing after surgery, by injection, by means of a catheter, by means of a suppository, or by means of an implant, said implant being of a porous, non-porous, or gelatinous material, including membranes, such as sialastic membranes, or fibers. Preferably, when administering a protein, including an antibody, of the invention, care must be taken to use materials to which the protein does not absorb.

[408] In another embodiment, the compound or composition can be delivered in a vesicle, in particular a liposome (see Langer, *Science* 249:1527-1533 (1990); Treat et al., in *Liposomes in the Therapy of Infectious Disease and Cancer*, Lopez-Berestein and Fidler (eds.), Liss, New York, pp. 353- 365 (1989); Lopez-Berestein, *ibid.*, pp. 317-327; see generally *ibid.*)

[409] In yet another embodiment, the compound or composition can be delivered in a controlled release system. In one embodiment, a pump may be used (see Langer, *supra*; Sefton, *CRC Crit. Ref. Biomed. Eng.* 14:201 (1987); Buchwald et al., *Surgery* 88:507 (1980); Saudek et al., *N. Engl. J. Med.* 321:574 (1989)). In another embodiment, polymeric materials can be used (see *Medical Applications of Controlled Release*, Langer and Wise (eds.), CRC Pres., Boca Raton, Florida (1974); *Controlled Drug Bioavailability, Drug Product Design and Performance*, Smolen and Ball (eds.), Wiley, New York (1984); Ranger and Peppas, J., *Macromol. Sci. Rev. Macromol. Chem.* 23:61 (1983); see also Levy

et al., Science 228:190 (1985); During et al., Ann. Neurol. 25:351 (1989); Howard et al., J.Neurosurg. 71:105 (1989)). In yet another embodiment, a controlled release system can be placed in proximity of the therapeutic target, e.g., the brain, thus requiring only a fraction of the systemic dose (see, e.g., Goodson, in Medical Applications of Controlled Release, *supra*, vol. 2, pp. 115-138 (1984)).

[410] Other controlled release systems are discussed in the review by Langer (Science 249:1527-1533 (1990)).

[411] In a specific embodiment where the compound of the invention is a nucleic acid encoding a protein, the nucleic acid can be administered *in vivo* to promote expression of its encoded protein, by constructing it as part of an appropriate nucleic acid expression vector and administering it so that it becomes intracellular, e.g., by use of a retroviral vector (see U.S. Patent No. 4,980,286), or by direct injection, or by use of microparticle bombardment (e.g., a gene gun; Biolistic, Dupont), or coating with lipids or cell-surface receptors or transfecting agents, or by administering it in linkage to a homeobox- like peptide which is known to enter the nucleus (see e.g., Joliot et al., Proc. Natl. Acad. Sci. USA 88:1864-1868 (1991)), etc. Alternatively, a nucleic acid can be introduced intracellularly and incorporated within host cell DNA for expression, by homologous recombination.

[412] The present invention also provides pharmaceutical compositions. Such compositions comprise a therapeutically effective amount of a compound, and a pharmaceutically acceptable carrier. In a specific embodiment, the term "pharmaceutically acceptable" means approved by a regulatory agency of the Federal or a state government or listed in the U.S. Pharmacopeia or other generally recognized pharmacopeia for use in animals, and more particularly in humans. The term "carrier" refers to a diluent, adjuvant, excipient, or vehicle with which the therapeutic is administered. Such pharmaceutical carriers can be sterile liquids, such as water and oils, including those of petroleum, animal, vegetable or synthetic origin, such as peanut oil, soybean oil, mineral oil, sesame oil and the like. Water is a preferred carrier when the pharmaceutical composition is administered intravenously. Saline solutions and aqueous dextrose and glycerol solutions can also be employed as liquid carriers, particularly for injectable solutions. Suitable pharmaceutical excipients include starch, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, sodium stearate, glycerol monostearate, talc, sodium chloride, dried skim milk, glycerol, propylene, glycol, water, ethanol and the like. The composition, if desired, can also contain minor amounts of wetting or emulsifying agents, or pH buffering agents.

These compositions can take the form of solutions, suspensions, emulsion, tablets, pills, capsules, powders, sustained-release formulations and the like. The composition can be formulated as a suppository, with traditional binders and carriers such as triglycerides. Oral formulation can include standard carriers such as pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate, etc. Examples of suitable pharmaceutical carriers are described in "Remington's Pharmaceutical Sciences" by E.W. Martin. Such compositions will contain a therapeutically effective amount of the compound, preferably in purified form, together with a suitable amount of carrier so as to provide the form for proper administration to the patient. The formulation should suit the mode of administration.

[413] In a preferred embodiment, the composition is formulated in accordance with routine procedures as a pharmaceutical composition adapted for intravenous administration to human beings. Typically, compositions for intravenous administration are solutions in sterile isotonic aqueous buffer. Where necessary, the composition may also include a solubilizing agent and a local anesthetic such as lignocaine to ease pain at the site of the injection. Generally, the ingredients are supplied either separately or mixed together in unit dosage form, for example, as a dry lyophilized powder or water free concentrate in a hermetically sealed container such as an ampoule or sachette indicating the quantity of active agent. Where the composition is to be administered by infusion, it can be dispensed with an infusion bottle containing sterile pharmaceutical grade water or saline. Where the composition is administered by injection, an ampoule of sterile water for injection or saline can be provided so that the ingredients may be mixed prior to administration.

[414] The compounds of the invention can be formulated as neutral or salt forms. Pharmaceutically acceptable salts include those formed with anions such as those derived from hydrochloric, phosphoric, acetic, oxalic, tartaric acids, etc., and those formed with cations such as those derived from sodium, potassium, ammonium, calcium, ferric hydroxides, isopropylamine, triethylamine, 2-ethylamino ethanol, histidine, procaine, etc.

[415] The amount of the compound of the invention which will be effective in the treatment, inhibition and prevention of a disease or disorder associated with aberrant expression and/or activity of a polypeptide of the invention can be determined by standard clinical techniques. In addition, in vitro assays may optionally be employed to help identify optimal dosage ranges. The precise dose to be employed in the formulation will also depend on the route of administration, and the seriousness of the disease or disorder,

and should be decided according to the judgment of the practitioner and each patient's circumstances. Effective doses may be extrapolated from dose-response curves derived from in vitro or animal model test systems.

[416] For antibodies, the dosage administered to a patient is typically 0.1 mg/kg to 100 mg/kg of the patient's body weight. Preferably, the dosage administered to a patient is between 0.1 mg/kg and 20 mg/kg of the patient's body weight, more preferably 1 mg/kg to 10 mg/kg of the patient's body weight. Generally, human antibodies have a longer half-life within the human body than antibodies from other species due to the immune response to the foreign polypeptides. Thus, lower dosages of human antibodies and less frequent administration is often possible. Further, the dosage and frequency of administration of antibodies of the invention may be reduced by enhancing uptake and tissue penetration (e.g., into the brain) of the antibodies by modifications such as, for example, lipidation.

[417] The invention also provides a pharmaceutical pack or kit comprising one or more containers filled with one or more of the ingredients of the pharmaceutical compositions of the invention. Optionally associated with such container(s) can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, which notice reflects approval by the agency of manufacture, use or sale for human administration.

Diagnosis and Imaging

[418] Labeled antibodies, and derivatives and analogs thereof, which specifically bind to a polypeptide of interest can be used for diagnostic purposes to detect, diagnose, or monitor diseases, disorders, and/or conditions associated with the aberrant expression and/or activity of a polypeptide of the invention. The invention provides for the detection of aberrant expression of a polypeptide of interest, comprising (a) assaying the expression of the polypeptide of interest in cells or body fluid of an individual using one or more antibodies specific to the polypeptide interest and (b) comparing the level of gene expression with a standard gene expression level, whereby an increase or decrease in the assayed polypeptide gene expression level compared to the standard expression level is indicative of aberrant expression.

[419] The invention provides a diagnostic assay for diagnosing a disorder, comprising (a) assaying the expression of the polypeptide of interest in cells or body fluid of an individual using one or more antibodies specific to the polypeptide interest and (b)

comparing the level of gene expression with a standard gene expression level, whereby an increase or decrease in the assayed polypeptide gene expression level compared to the standard expression level is indicative of a particular disorder. With respect to cancer, the presence of a relatively high amount of transcript in biopsied tissue from an individual may indicate a predisposition for the development of the disease, or may provide a means for detecting the disease prior to the appearance of actual clinical symptoms. A more definitive diagnosis of this type may allow health professionals to employ preventative measures or aggressive treatment earlier thereby preventing the development or further progression of the cancer.

[420] Antibodies of the invention can be used to assay protein levels in a biological sample using classical immunohistological methods known to those of skill in the art (e.g., see Jalkanen et al., J. Cell. Biol. 101:976-985 (1985); Jalkanen et al., J. Cell. Biol. 105:3087-3096 (1987)). Other antibody-based methods useful for detecting protein gene expression include immunoassays, such as the enzyme linked immunosorbent assay (ELISA) and the radioimmunoassay (RIA). Suitable antibody assay labels are known in the art and include enzyme labels, such as, glucose oxidase; radioisotopes, such as iodine (125I, 121I), carbon (14C), sulfur (35S), tritium (3H), indium (112In), and technetium (99Tc); luminescent labels, such as luminol; and fluorescent labels, such as fluorescein and rhodamine, and biotin.

[421] One facet of the invention is the detection and diagnosis of a disease or disorder associated with aberrant expression of a polypeptide of interest in an animal, preferably a mammal and most preferably a human. In one embodiment, diagnosis comprises: a) administering (for example, parenterally, subcutaneously, or intraperitoneally) to a subject an effective amount of a labeled molecule which specifically binds to the polypeptide of interest; b) waiting for a time interval following the administering for permitting the labeled molecule to preferentially concentrate at sites in the subject where the polypeptide is expressed (and for unbound labeled molecule to be cleared to background level); c) determining background level; and d) detecting the labeled molecule in the subject, such that detection of labeled molecule above the background level indicates that the subject has a particular disease or disorder associated with aberrant expression of the polypeptide of interest. Background level can be determined by various methods including, comparing the amount of labeled molecule detected to a standard value previously determined for a particular system.

[422] It will be understood in the art that the size of the subject and the imaging system used will determine the quantity of imaging moiety needed to produce diagnostic images. In the case of a radioisotope moiety, for a human subject, the quantity of radioactivity injected will normally range from about 5 to 20 millicuries of ^{99m}Tc . The labeled antibody or antibody fragment will then preferentially accumulate at the location of cells which contain the specific protein. *In vivo* tumor imaging is described in S.W. Burchiel et al., "Immunopharmacokinetics of Radiolabeled Antibodies and Their Fragments." (Chapter 13 in *Tumor Imaging: The Radiochemical Detection of Cancer*, S.W. Burchiel and B. A. Rhodes, eds., Masson Publishing Inc. (1982)).

[423] Depending on several variables, including the type of label used and the mode of administration, the time interval following the administration for permitting the labeled molecule to preferentially concentrate at sites in the subject and for unbound labeled molecule to be cleared to background level is 6 to 48 hours or 6 to 24 hours or 6 to 12 hours. In another embodiment the time interval following administration is 5 to 20 days or 5 to 10 days.

[424] In an embodiment, monitoring of the disease or disorder is carried out by repeating the method for diagnosing the disease or disease, for example, one month after initial diagnosis, six months after initial diagnosis, one year after initial diagnosis, etc.

[425] Presence of the labeled molecule can be detected in the patient using methods known in the art for *in vivo* scanning. These methods depend upon the type of label used. Skilled artisans will be able to determine the appropriate method for detecting a particular label. Methods and devices that may be used in the diagnostic methods of the invention include, but are not limited to, computed tomography (CT), whole body scan such as position emission tomography (PET), magnetic resonance imaging (MRI), and sonography.

[426] In a specific embodiment, the molecule is labeled with a radioisotope and is detected in the patient using a radiation responsive surgical instrument (Thurston et al., U.S. Patent No. 5,441,050). In another embodiment, the molecule is labeled with a fluorescent compound and is detected in the patient using a fluorescence responsive scanning instrument. In another embodiment, the molecule is labeled with a positron emitting metal and is detected in the patient using positron emission-tomography. In yet another embodiment, the molecule is labeled with a paramagnetic label and is detected in a patient using magnetic resonance imaging (MRI).

Kits

[427] The present invention provides kits that can be used in the above methods. In one embodiment, a kit comprises an antibody of the invention, preferably a purified antibody, in one or more containers. In a specific embodiment, the kits of the present invention contain a substantially isolated polypeptide comprising an epitope which is specifically immunoreactive with an antibody included in the kit. Preferably, the kits of the present invention further comprise a control antibody which does not react with the polypeptide of interest. In another specific embodiment, the kits of the present invention contain a means for detecting the binding of an antibody to a polypeptide of interest (e.g., the antibody may be conjugated to a detectable substrate such as a fluorescent compound, an enzymatic substrate, a radioactive compound or a luminescent compound, or a second antibody which recognizes the first antibody may be conjugated to a detectable substrate).

[428] In another specific embodiment of the present invention, the kit is a diagnostic kit for use in screening serum containing antibodies specific against proliferative and/or cancerous polynucleotides and polypeptides. Such a kit may include a control antibody that does not react with the polypeptide of interest. Such a kit may include a substantially isolated polypeptide antigen comprising an epitope which is specifically immunoreactive with at least one anti-polypeptide antigen antibody. Further, such a kit includes means for detecting the binding of said antibody to the antigen (e.g., the antibody may be conjugated to a fluorescent compound such as fluorescein or rhodamine which can be detected by flow cytometry). In specific embodiments, the kit may include a recombinantly produced or chemically synthesized polypeptide antigen. The polypeptide antigen of the kit may also be attached to a solid support.

[429] In a more specific embodiment the detecting means of the above-described kit includes a solid support to which said polypeptide antigen is attached. Such a kit may also include a non-attached reporter-labeled anti-human antibody. In this embodiment, binding of the antibody to the polypeptide antigen can be detected by binding of the said reporter-labeled antibody.

[430] In an additional embodiment, the invention includes a diagnostic kit for use in screening serum containing antigens of the polypeptide of the invention. The diagnostic kit includes a substantially isolated antibody specifically immunoreactive with polypeptide or polynucleotide antigens, and means for detecting the binding of the polynucleotide or polypeptide antigen to the antibody. In one embodiment, the antibody is attached to a solid

support. In a specific embodiment, the antibody may be a monoclonal antibody. The detecting means of the kit may include a second, labeled monoclonal antibody. Alternatively, or in addition, the detecting means may include a labeled, competing antigen.

[431] In one diagnostic configuration, test serum is reacted with a solid phase reagent having a surface-bound antigen obtained by the methods of the present invention. After binding with specific antigen antibody to the reagent and removing unbound serum components by washing, the reagent is reacted with reporter-labeled anti-human antibody to bind reporter to the reagent in proportion to the amount of bound anti-antigen antibody on the solid support. The reagent is again washed to remove unbound labeled antibody, and the amount of reporter associated with the reagent is determined. Typically, the reporter is an enzyme which is detected by incubating the solid phase in the presence of a suitable fluorometric, luminescent or colorimetric substrate (Sigma, St. Louis, MO).

[432] The solid surface reagent in the above assay is prepared by known techniques for attaching protein material to solid support material, such as polymeric beads, dip sticks, 96-well plate or filter material. These attachment methods generally include non-specific adsorption of the protein to the support or covalent attachment of the protein, typically through a free amine group, to a chemically reactive group on the solid support, such as an activated carboxyl, hydroxyl, or aldehyde group. Alternatively, streptavidin coated plates can be used in conjunction with biotinylated antigen(s).

[433] Thus, the invention provides an assay system or kit for carrying out this diagnostic method. The kit generally includes a support with surface-bound recombinant antigens, and a reporter-labeled anti-human antibody for detecting surface-bound anti-antigen antibody.

Uses of the Polynucleotides

[434] Each of the polynucleotides identified herein can be used in numerous ways as reagents. The following description should be considered exemplary and utilizes known techniques.

[435] The polynucleotides of the present invention are useful for chromosome identification. There exists an ongoing need to identify new chromosome markers, since few chromosome marking reagents, based on actual sequence data (repeat polymorphisms), are presently available. Each sequence is specifically targeted to and can hybridize with a

particular location on an individual human chromosome, thus each polynucleotide of the present invention can routinely be used as a chromosome marker using techniques known in the art. Table 1B, column 9 provides the chromosome location of some of the polynucleotides of the invention.

[436] Briefly, sequences can be mapped to chromosomes by preparing PCR primers (preferably at least 15 bp (e.g., 15-25 bp) from the sequences shown in SEQ ID NO:X. Primers can optionally be selected using computer analysis so that primers do not span more than one predicted exon in the genomic DNA. These primers are then used for PCR screening of somatic cell hybrids containing individual human chromosomes. Only those hybrids containing the human gene corresponding to SEQ ID NO:X will yield an amplified fragment.

[437] Similarly, somatic hybrids provide a rapid method of PCR mapping the polynucleotides to particular chromosomes. Three or more clones can be assigned per day using a single thermal cycler. Moreover, sublocalization of the polynucleotides can be achieved with panels of specific chromosome fragments. Other gene mapping strategies that can be used include in situ hybridization, prescreening with labeled flow-sorted chromosomes, preselection by hybridization to construct chromosome specific-cDNA libraries, and computer mapping techniques (See, e.g., Shuler, Trends Biotechnol 16:456-459 (1998) which is hereby incorporated by reference in its entirety).

[438] Precise chromosomal location of the polynucleotides can also be achieved using fluorescence in situ hybridization (FISH) of a metaphase chromosomal spread. This technique uses polynucleotides as short as 500 or 600 bases; however, polynucleotides 2,000-4,000 bp are preferred. For a review of this technique, see Verma et al., "Human Chromosomes: a Manual of Basic Techniques," Pergamon Press, New York (1988).

[439] For chromosome mapping, the polynucleotides can be used individually (to mark a single chromosome or a single site on that chromosome) or in panels (for marking multiple sites and/or multiple chromosomes).

[440] Thus, the present invention also provides a method for chromosomal localization which involves (a) preparing PCR primers from the polynucleotide sequences in Table 1B and/or Table 2 and SEQ ID NO:X and (b) screening somatic cell hybrids containing individual chromosomes.

[441] The polynucleotides of the present invention would likewise be useful for radiation hybrid mapping, HAPPY mapping, and long range restriction mapping. For a

review of these techniques and others known in the art, see, e.g. Dear, "Genome Mapping: A Practical Approach," IRL Press at Oxford University Press, London (1997); Aydin, J. Mol. Med. 77:691-694 (1999); Hacia et al., Mol. Psychiatry 3:483-492 (1998); Herrick et al., Chromosome Res. 7:409-423 (1999); Hamilton et al., Methods Cell Biol. 62:265-280 (2000); and/or Ott, J. Hered. 90:68-70 (1999) each of which is hereby incorporated by reference in its entirety.

[442] Once a polynucleotide has been mapped to a precise chromosomal location, the physical position of the polynucleotide can be used in linkage analysis. Linkage analysis establishes coinheritance between a chromosomal location and presentation of a particular disease. (Disease mapping data are found, for example, in V. McKusick, Mendelian Inheritance in Man (available on line through Johns Hopkins University Welch Medical Library)). Column 10 of Table 1B provides an OMIM reference identification number of diseases associated with the cytologic band disclosed in column 9 of Table 1B, as determined using techniques described herein and by reference to Table 5. Assuming 1 megabase mapping resolution and one gene per 20 kb, a cDNA precisely localized to a chromosomal region associated with the disease could be one of 50-500 potential causative genes.

[443] Thus, once coinheritance is established, differences in a polynucleotide of the invention and the corresponding gene between affected and unaffected individuals can be examined. First, visible structural alterations in the chromosomes, such as deletions or translocations, are examined in chromosome spreads or by PCR. If no structural alterations exist, the presence of point mutations are ascertained. Mutations observed in some or all affected individuals, but not in normal individuals, indicates that the mutation may cause the disease. However, complete sequencing of the polypeptide and the corresponding gene from several normal individuals is required to distinguish the mutation from a polymorphism. If a new polymorphism is identified, this polymorphic polypeptide can be used for further linkage analysis.

[444] Furthermore, increased or decreased expression of the gene in affected individuals as compared to unaffected individuals can be assessed using the polynucleotides of the invention. Any of these alterations (altered expression, chromosomal rearrangement, or mutation) can be used as a diagnostic or prognostic marker. Diagnostic and prognostic methods, kits and reagents encompassed by the present invention are briefly described

below and more thoroughly elsewhere herein (see e.g., the sections labeled "Antibodies", "Diagnostic Assays", and "Methods for Detecting Diseases").

[445] Thus, the invention also provides a diagnostic method useful during diagnosis of a disorder, involving measuring the expression level of polynucleotides of the present invention in cells or body fluid from an individual and comparing the measured gene expression level with a standard level of polynucleotide expression level, whereby an increase or decrease in the gene expression level compared to the standard is indicative of a disorder. Additional non-limiting examples of diagnostic methods encompassed by the present invention are more thoroughly described elsewhere herein (see, e.g., Example 12).

[446] In still another embodiment, the invention includes a kit for analyzing samples for the presence of proliferative and/or cancerous polynucleotides derived from a test subject. In a general embodiment, the kit includes at least one polynucleotide probe containing a nucleotide sequence that will specifically hybridize with a polynucleotide of the invention and a suitable container. In a specific embodiment, the kit includes two polynucleotide probes defining an internal region of the polynucleotide of the invention, where each probe has one strand containing a 31' mer-end internal to the region. In a further embodiment, the probes may be useful as primers for polymerase chain reaction amplification.

[447] Where a diagnosis of a related disorder, including, for example, diagnosis of a tumor, has already been made according to conventional methods, the present invention is useful as a prognostic indicator, whereby patients exhibiting enhanced or depressed polynucleotide of the invention expression will experience a worse clinical outcome relative to patients expressing the gene at a level nearer the standard level.

[448] By "measuring the expression level of polynucleotides of the invention" is intended qualitatively or quantitatively measuring or estimating the level of the polypeptide of the invention or the level of the mRNA encoding the polypeptide of the invention in a first biological sample either directly (e.g., by determining or estimating absolute protein level or mRNA level) or relatively (e.g., by comparing to the polypeptide level or mRNA level in a second biological sample). Preferably, the polypeptide level or mRNA level in the first biological sample is measured or estimated and compared to a standard polypeptide level or mRNA level, the standard being taken from a second biological sample obtained from an individual not having the related disorder or being determined by averaging levels from a population of individuals not having a related disorder. As will be appreciated in the

art, once a standard polypeptide level or mRNA level is known, it can be used repeatedly as a standard for comparison.

[449] By "biological sample" is intended any biological sample obtained from an individual, body fluid, cell line, tissue culture, or other source which contains polypeptide of the present invention or the corresponding mRNA. As indicated, biological samples include body fluids (such as semen, lymph, vaginal pool, sera, plasma, urine, synovial fluid and spinal fluid) which contain the polypeptide of the present invention, and tissue sources found to express the polypeptide of the present invention. Methods for obtaining tissue biopsies and body fluids from mammals are well known in the art. Where the biological sample is to include mRNA, a tissue biopsy is the preferred source.

[450] The method(s) provided above may preferably be applied in a diagnostic method and/or kits in which polynucleotides and/or polypeptides of the invention are attached to a solid support. In one exemplary method, the support may be a "gene chip" or a "biological chip" as described in US Patents 5,837,832, 5,874,219, and 5,856,174. Further, such a gene chip with polynucleotides of the invention attached may be used to identify polymorphisms between the isolated polynucleotide sequences of the invention, with polynucleotides isolated from a test subject. The knowledge of such polymorphisms (i.e. their location, as well as, their existence) would be beneficial in identifying disease loci for many disorders, such as for example, in neural disorders, immune system disorders, muscular disorders, reproductive disorders, gastrointestinal disorders, pulmonary disorders, digestive disorders, metabolic disorders, cardiovascular disorders, renal disorders, proliferative disorders, and/or cancerous diseases and conditions. Such a method is described in US Patents 5,858,659 and 5,856,104. The US Patents referenced *supra* are hereby incorporated by reference in their entirety herein.

[451] The present invention encompasses polynucleotides of the present invention that are chemically synthesized, or reproduced as peptide nucleic acids (PNA), or according to other methods known in the art. The use of PNAs would serve as the preferred form if the polynucleotides of the invention are incorporated onto a solid support, or gene chip. For the purposes of the present invention, a peptide nucleic acid (PNA) is a polyamide type of DNA analog and the monomeric units for adenine, guanine, thymine and cytosine are available commercially (Perceptive Biosystems). Certain components of DNA, such as phosphorus, phosphorus oxides, or deoxyribose derivatives, are not present in PNAs. As disclosed by Nielsen et al., Science 254, 1497 (1991); and Egholm et al., Nature 365, 666 (1993), PNAs

bind specifically and tightly to complementary DNA strands and are not degraded by nucleases. In fact, PNA binds more strongly to DNA than DNA itself does. This is probably because there is no electrostatic repulsion between the two strands, and also the polyamide backbone is more flexible. Because of this, PNA/DNA duplexes bind under a wider range of stringency conditions than DNA/DNA duplexes, making it easier to perform multiplex hybridization. Smaller probes can be used than with DNA due to the strong binding. In addition, it is more likely that single base mismatches can be determined with PNA/DNA hybridization because a single mismatch in a PNA/DNA 15-mer lowers the melting point (T_{sub.m}) by 8°-20° C, vs. 4°-16° C for the DNA/DNA 15-mer duplex. Also, the absence of charge groups in PNA means that hybridization can be done at low ionic strengths and reduce possible interference by salt during the analysis.

[452] The compounds of the present invention have uses which include, but are not limited to, detecting cancer in mammals. In particular the invention is useful during diagnosis of pathological cell proliferative neoplasias which include, but are not limited to: acute myelogenous leukemias including acute monocytic leukemia, acute myeloblastic leukemia, acute promyelocytic leukemia, acute myelomonocytic leukemia, acute erythroleukemia, acute megakaryocytic leukemia, and acute undifferentiated leukemia, etc.; and chronic myelogenous leukemias including chronic myelomonocytic leukemia, chronic granulocytic leukemia, etc. Preferred mammals include monkeys, apes, cats, dogs, cows, pigs, horses, rabbits and humans. Particularly preferred are humans.

[453] Pathological cell proliferative disorders are often associated with inappropriate activation of proto-oncogenes. (Germann, E. P. et al., "The Etiology of Acute Leukemia: Molecular Genetics and Viral Oncology," in Neoplastic Diseases of the Blood, Vol 1., Wiernik, P. H. et al. eds., 161-182 (1985)). Neoplasias are now believed to result from the qualitative alteration of a normal cellular gene product, or from the quantitative modification of gene expression by insertion into the chromosome of a viral sequence, by chromosomal translocation of a gene to a more actively transcribed region, or by some other mechanism. (Germann et al., *supra*) It is likely that mutated or altered expression of specific genes is involved in the pathogenesis of some leukemias, among other tissues and cell types. (Germann et al., *supra*) Indeed, the human counterparts of the oncogenes involved in some animal neoplasias have been amplified or translocated in some cases of human leukemia and carcinoma. (Germann et al., *supra*)

09950083-091201

[454] For example, c-myc expression is highly amplified in the non-lymphocytic leukemia cell line HL-60. When HL-60 cells are chemically induced to stop proliferation, the level of c-myc is found to be downregulated. (International Publication Number WO 91/15580). However, it has been shown that exposure of HL-60 cells to a DNA construct that is complementary to the 5' end of c-myc or c-myb blocks translation of the corresponding mRNAs which downregulates expression of the c-myc or c-myb proteins and causes arrest of cell proliferation and differentiation of the treated cells. (International Publication Number WO 91/15580; Wickstrom et al., Proc. Natl. Acad. Sci. 85:1028 (1988); Anfossi et al., Proc. Natl. Acad. Sci. 86:3379 (1989)). However, the skilled artisan would appreciate the present invention's usefulness is not be limited to treatment, prevention, and/or prognosis of proliferative disorders of cells and tissues of hematopoietic origin, in light of the numerous cells and cell types of varying origins which are known to exhibit proliferative phenotypes.

[455] In addition to the foregoing, a polynucleotide of the present invention can be used to control gene expression through triple helix formation or through antisense DNA or RNA. Antisense techniques are discussed, for example, in Okano, J. Neurochem. 56: 560 (1991); "Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988). Triple helix formation is discussed in, for instance Lee et al., Nucleic Acids Research 6: 3073 (1979); Cooney et al., Science 241: 456 (1988); and Dervan et al., Science 251: 1360 (1991). Both methods rely on binding of the polynucleotide to a complementary DNA or RNA. For these techniques, preferred polynucleotides are usually oligonucleotides 20 to 40 bases in length and complementary to either the region of the gene involved in transcription (triple helix - see Lee et al., Nucl. Acids Res. 6:3073 (1979); Cooney et al., Science 241:456 (1988); and Dervan et al., Science 251:1360 (1991)) or to the mRNA itself (antisense - Okano, J. Neurochem. 56:560 (1991); Oligodeoxy-nucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988)). Triple helix formation optimally results in a shut-off of RNA transcription from DNA, while antisense RNA hybridization blocks translation of an mRNA molecule into polypeptide. The oligonucleotide described above can also be delivered to cells such that the antisense RNA or DNA may be expressed *in vivo* to inhibit production of polypeptide of the present invention antigens. Both techniques are effective in model systems, and the information disclosed herein can be used to design antisense or triple helix polynucleotides in an effort to treat disease, and in particular, for the treatment

of proliferative diseases and/or conditions. Non-limiting antisense and triple helix methods encompassed by the present invention are more thoroughly described elsewhere herein (see, e.g., the section labeled "Antisense and Ribozyme (Antagonists)").

[456] Polynucleotides of the present invention are also useful in gene therapy. One goal of gene therapy is to insert a normal gene into an organism having a defective gene, in an effort to correct the genetic defect. The polynucleotides disclosed in the present invention offer a means of targeting such genetic defects in a highly accurate manner. Another goal is to insert a new gene that was not present in the host genome, thereby producing a new trait in the host cell. Additional non-limiting examples of gene therapy methods encompassed by the present invention are more thoroughly described elsewhere herein (see, e.g., the sections labeled "Gene Therapy Methods", and Examples 16, 17 and 18).

[457] The polynucleotides are also useful for identifying individuals from minute biological samples. The United States military, for example, is considering the use of restriction fragment length polymorphism (RFLP) for identification of its personnel. In this technique, an individual's genomic DNA is digested with one or more restriction enzymes, and probed on a Southern blot to yield unique bands for identifying personnel. This method does not suffer from the current limitations of "Dog Tags" which can be lost, switched, or stolen, making positive identification difficult. The polynucleotides of the present invention can be used as additional DNA markers for RFLP.

[458] The polynucleotides of the present invention can also be used as an alternative to RFLP, by determining the actual base-by-base DNA sequence of selected portions of an individual's genome. These sequences can be used to prepare PCR primers for amplifying and isolating such selected DNA, which can then be sequenced. Using this technique, individuals can be identified because each individual will have a unique set of DNA sequences. Once an unique ID database is established for an individual, positive identification of that individual, living or dead, can be made from extremely small tissue samples.

[459] Forensic biology also benefits from using DNA-based identification techniques as disclosed herein. DNA sequences taken from very small biological samples such as tissues, e.g., hair or skin, or body fluids, e.g., blood, saliva, semen, synovial fluid, amniotic fluid, breast milk, lymph, pulmonary sputum or surfactant, urine, fecal matter, etc., can be amplified using PCR. In one prior art technique, gene sequences amplified from

polymorphic loci, such as DQa class II HLA gene, are used in forensic biology to identify individuals. (Erlich, H., PCR Technology, Freeman and Co. (1992)). Once these specific polymorphic loci are amplified, they are digested with one or more restriction enzymes, yielding an identifying set of bands on a Southern blot probed with DNA corresponding to the DQa class II HLA gene. Similarly, polynucleotides of the present invention can be used as polymorphic markers for forensic purposes.

[460] There is also a need for reagents capable of identifying the source of a particular tissue. Such need arises, for example, in forensics when presented with tissue of unknown origin. Appropriate reagents can comprise, for example, DNA probes or primers prepared from the sequences of the present invention, specific to tissues, including but not limited to those shown in Table 1B. Panels of such reagents can identify tissue by species and/or by organ type. In a similar fashion, these reagents can be used to screen tissue cultures for contamination. Additional non-limiting examples of such uses are further described herein.

[461] The polynucleotides of the present invention are also useful as hybridization probes for differential identification of the tissue(s) or cell type(s) present in a biological sample. Similarly, polypeptides and antibodies directed to polypeptides of the present invention are useful to provide immunological probes for differential identification of the tissue(s) (e.g., immunohistochemistry assays) or cell type(s) (e.g., immunocytochemistry assays). In addition, for a number of disorders of the above tissues or cells, significantly higher or lower levels of gene expression of the polynucleotides/polypeptides of the present invention may be detected in certain tissues (e.g., tissues expressing polypeptides and/or polynucleotides of the present invention, for example, those disclosed in column 8 of Table 1B, and/or cancerous and/or wounded tissues) or bodily fluids (e.g., semen, lymph, vaginal pool, serum, plasma, urine, synovial fluid or spinal fluid) taken from an individual having such a disorder, relative to a "standard" gene expression level, i.e., the expression level in healthy tissue from an individual not having the disorder.

[462] Thus, the invention provides a diagnostic method of a disorder, which involves: (a) assaying gene expression level in cells or body fluid of an individual; (b) comparing the gene expression level with a standard gene expression level, whereby an increase or decrease in the assayed gene expression level compared to the standard expression level is indicative of a disorder.

[463] In the very least, the polynucleotides of the present invention can be used as molecular weight markers on Southern gels, as diagnostic probes for the presence of a

specific mRNA in a particular cell type, as a probe to "subtract-out" known sequences in the process of discovering novel polynucleotides, for selecting and making oligomers for attachment to a "gene chip" or other support, to raise anti-DNA antibodies using DNA immunization techniques, and as an antigen to elicit an immune response.

Uses of the Polypeptides

[464] Each of the polypeptides identified herein can be used in numerous ways. The following description should be considered exemplary and utilizes known techniques.

[465] Polypeptides and antibodies directed to polypeptides of the present invention are useful to provide immunological probes for differential identification of the tissue(s) (e.g., immunohistochemistry assays such as, for example, ABC immunoperoxidase (Hsu et al., J. Histochem. Cytochem. 29:577-580 (1981)) or cell type(s) (e.g., immunocytochemistry assays).

[466] Antibodies can be used to assay levels of polypeptides encoded by polynucleotides of the invention in a biological sample using classical immunohistological methods known to those of skill in the art (e.g., see Jalkanen, et al., J. Cell. Biol. 101:976-985 (1985); Jalkanen, et al., J. Cell. Biol. 105:3087-3096 (1987)). Other antibody-based methods useful for detecting protein gene expression include immunoassays, such as the enzyme linked immunosorbent assay (ELISA) and the radioimmunoassay (RIA). Suitable antibody assay labels are known in the art and include enzyme labels, such as, glucose oxidase; radioisotopes, such as iodine (^{131}I , ^{125}I , ^{123}I , ^{121}I), carbon (^{14}C), sulfur (^{35}S), tritium (^3H), indium ($^{115\text{m}}\text{In}$, $^{113\text{m}}\text{In}$, ^{112}In , ^{111}In), and technetium (^{99}Tc , $^{99\text{m}}\text{Tc}$), thallium (^{201}Tl), gallium (^{68}Ga , ^{67}Ga), palladium (^{103}Pd), molybdenum (^{99}Mo), xenon (^{133}Xe), fluorine (^{18}F), ^{153}Sm , ^{177}Lu , ^{159}Gd , ^{149}Pm , ^{140}La , ^{175}Yb , ^{166}Ho , ^{90}Y , ^{47}Sc , ^{186}Re , ^{188}Re , ^{142}Pr , ^{105}Rh , ^{97}Ru ; luminescent labels, such as luminol; and fluorescent labels, such as fluorescein and rhodamine, and biotin.

[467] In addition to assaying levels of polypeptide of the present invention in a biological sample, proteins can also be detected *in vivo* by imaging. Antibody labels or markers for *in vivo* imaging of protein include those detectable by X-radiography, NMR or ESR. For X-radiography, suitable labels include radioisotopes such as barium or cesium, which emit detectable radiation but are not overtly harmful to the subject. Suitable markers for NMR and ESR include those with a detectable characteristic spin, such as deuterium,

which may be incorporated into the antibody by labeling of nutrients for the relevant hybridoma.

[468] A protein-specific antibody or antibody fragment which has been labeled with an appropriate detectable imaging moiety, such as a radioisotope (for example, ^{131}I , ^{112}In , $^{99\text{m}}\text{Tc}$, (^{131}I , ^{125}I , ^{123}I , ^{121}I), carbon (^{14}C), sulfur (^{35}S), tritium (^3H), indium ($^{115\text{m}}\text{In}$, $^{113\text{m}}\text{In}$, ^{112}In , ^{111}In), and technetium (^{99}Tc , $^{99\text{m}}\text{Tc}$), thallium (^{201}Tl), gallium (^{68}Ga , ^{67}Ga), palladium (^{103}Pd), molybdenum (^{99}Mo), xenon (^{133}Xe), fluorine (^{18}F , ^{153}Sm , ^{177}Lu , ^{159}Gd , ^{149}Pm , ^{140}La , ^{175}Yb , ^{166}Ho , ^{90}Y , ^{47}Sc , ^{186}Re , ^{188}Re , ^{142}Pr , ^{105}Rh , ^{97}Ru), a radio-opaque substance, or a material detectable by nuclear magnetic resonance, is introduced (for example, parenterally, subcutaneously or intraperitoneally) into the mammal to be examined for immune system disorder. It will be understood in the art that the size of the subject and the imaging system used will determine the quantity of imaging moiety needed to produce diagnostic images. In the case of a radioisotope moiety, for a human subject, the quantity of radioactivity injected will normally range from about 5 to 20 millicuries of $^{99\text{m}}\text{Tc}$. The labeled antibody or antibody fragment will then preferentially accumulate at the location of cells which express the polypeptide encoded by a polynucleotide of the invention. *In vivo* tumor imaging is described in S.W. Burchiel et al., "Immunopharmacokinetics of Radiolabeled Antibodies and Their Fragments" (Chapter 13 in *Tumor Imaging: The Radiochemical Detection of Cancer*, S.W. Burchiel and B. A. Rhodes, eds., Masson Publishing Inc. (1982)).

[469] In one embodiment, the invention provides a method for the specific delivery of compositions of the invention to cells by administering polypeptides of the invention (e.g., polypeptides encoded by polynucleotides of the invention and/or antibodies) that are associated with heterologous polypeptides or nucleic acids. In one example, the invention provides a method for delivering a therapeutic protein into the targeted cell. In another example, the invention provides a method for delivering a single stranded nucleic acid (e.g., antisense or ribozymes) or double stranded nucleic acid (e.g., DNA that can integrate into the cell's genome or replicate episomally and that can be transcribed) into the targeted cell.

[470] In another embodiment, the invention provides a method for the specific destruction of cells (e.g., the destruction of tumor cells) by administering polypeptides of the invention in association with toxins or cytotoxic prodrugs.

[471] By "toxin" is meant one or more compounds that bind and activate endogenous cytotoxic effector systems, radioisotopes, holotoxins, modified toxins, catalytic subunits of

toxins, or any molecules or enzymes not normally present in or on the surface of a cell that under defined conditions cause the cell's death. Toxins that may be used according to the methods of the invention include, but are not limited to, radioisotopes known in the art, compounds such as, for example, antibodies (or complement fixing containing portions thereof) that bind an inherent or induced endogenous cytotoxic effector system, thymidine kinase, endonuclease, RNase, alpha toxin, ricin, abrin, *Pseudomonas* exotoxin A, diphtheria toxin, saporin, momordin, gelonin, pokeweed antiviral protein, alpha-sarcin and cholera toxin. "Toxin" also includes a cytostatic or cytocidal agent, a therapeutic agent or a radioactive metal ion, e.g., alpha-emitters such as, for example, ^{213}Bi , or other radioisotopes such as, for example, ^{103}Pd , ^{133}Xe , ^{131}I , ^{68}Ge , ^{57}Co , ^{65}Zn , ^{85}Sr , ^{32}P , ^{35}S , ^{90}Y , ^{153}Sm , ^{153}Gd , ^{169}Yb , ^{51}Cr , ^{54}Mn , ^{75}Se , ^{113}Sn , $^{90}\text{Yttrium}$, ^{117}Tin , $^{186}\text{Rhenium}$, $^{166}\text{Holmium}$, and $^{188}\text{Rhenium}$; luminescent labels, such as luminol; and fluorescent labels, such as fluorescein and rhodamine, and biotin. In a specific embodiment, the invention provides a method for the specific destruction of cells (e.g., the destruction of tumor cells) by administering polypeptides of the invention or antibodies of the invention in association with the radioisotope ^{90}Y . In another specific embodiment, the invention provides a method for the specific destruction of cells (e.g., the destruction of tumor cells) by administering polypeptides of the invention or antibodies of the invention in association with the radioisotope ^{111}In . In a further specific embodiment, the invention provides a method for the specific destruction of cells (e.g., the destruction of tumor cells) by administering polypeptides of the invention or antibodies of the invention in association with the radioisotope ^{131}I .

[472] Techniques known in the art may be applied to label polypeptides of the invention (including antibodies). Such techniques include, but are not limited to, the use of bifunctional conjugating agents (see e.g., U.S. Patent Nos. 5,756,065; 5,714,631; 5,696,239; 5,652,361; 5,505,931; 5,489,425; 5,435,990; 5,428,139; 5,342,604; 5,274,119; 4,994,560; and 5,808,003; the contents of each of which are hereby incorporated by reference in its entirety).

[473] Thus, the invention provides a diagnostic method of a disorder, which involves (a) assaying the expression level of a polypeptide of the present invention in cells or body fluid of an individual; and (b) comparing the assayed polypeptide expression level with a standard polypeptide expression level, whereby an increase or decrease in the assayed polypeptide expression level compared to the standard expression level is indicative of a

disorder. With respect to cancer, the presence of a relatively high amount of transcript in biopsied tissue from an individual may indicate a predisposition for the development of the disease, or may provide a means for detecting the disease prior to the appearance of actual clinical symptoms. A more definitive diagnosis of this type may allow health professionals to employ preventative measures or aggressive treatment earlier thereby preventing the development or further progression of the cancer.

[474] Moreover, polypeptides of the present invention can be used to treat or prevent diseases or conditions such as, for example, neural disorders, immune system disorders, muscular disorders, reproductive disorders, gastrointestinal disorders, pulmonary disorders, cardiovascular disorders, renal disorders, proliferative disorders, and/or cancerous diseases and conditions. For example, patients can be administered a polypeptide of the present invention in an effort to replace absent or decreased levels of the polypeptide (e.g., insulin), to supplement absent or decreased levels of a different polypeptide (e.g., hemoglobin S for hemoglobin B, SOD, catalase, DNA repair proteins), to inhibit the activity of a polypeptide (e.g., an oncogene or tumor suppressor), to activate the activity of a polypeptide (e.g., by binding to a receptor), to reduce the activity of a membrane bound receptor by competing with it for free ligand (e.g., soluble TNF receptors used in reducing inflammation), or to bring about a desired response (e.g., blood vessel growth inhibition, enhancement of the immune response to proliferative cells or tissues).

[475] Similarly, antibodies directed to a polypeptide of the present invention can also be used to treat disease (as described *supra*, and elsewhere herein). For example, administration of an antibody directed to a polypeptide of the present invention can bind, and/or neutralize the polypeptide, and/or reduce overproduction of the polypeptide. Similarly, administration of an antibody can activate the polypeptide, such as by binding to a polypeptide bound to a membrane (receptor).

[476] At the very least, the polypeptides of the present invention can be used as molecular weight markers on SDS-PAGE gels or on molecular sieve gel filtration columns using methods well known to those of skill in the art. Polypeptides can also be used to raise antibodies, which in turn are used to measure protein expression from a recombinant cell, as a way of assessing transformation of the host cell. Moreover, the polypeptides of the present invention can be used to test the biological activities described herein.

Diagnostic Assays

[477] The compounds of the present invention are useful for diagnosis, treatment, prevention and/or prognosis of various disorders in mammals, preferably humans. Such disorders include, but are not limited to, those described for Table 1D and as indicated in the "Preferred Indications" column of Table 1D, and also herein under the section heading "Biological Activities".

[478] For a number of disorders, substantially altered (increased or decreased) levels of gene expression can be detected in tissues, cells or bodily fluids (e.g., sera, plasma, urine, semen, synovial fluid or spinal fluid) taken from an individual having such a disorder, relative to a "standard" gene expression level, that is, the expression level in tissues or bodily fluids from an individual not having the disorder. Thus, the invention provides a diagnostic method useful during diagnosis of a disorder, which involves measuring the expression level of the gene encoding the polypeptide in tissues, cells or body fluid from an individual and comparing the measured gene expression level with a standard gene expression level, whereby an increase or decrease in the gene expression level(s) compared to the standard is indicative of a disorder. These diagnostic assays may be performed *in vivo* or *in vitro*, such as, for example, on blood samples, biopsy tissue or autopsy tissue.

[479] The present invention is also useful as a prognostic indicator, whereby patients exhibiting enhanced or depressed gene expression will experience a worse clinical outcome relative to patients expressing the gene at a level nearer the standard level.

[480] In certain embodiments, a polypeptide of the invention, or polynucleotides, antibodies, agonists, or antagonists corresponding to that polypeptide, may be used to diagnose and/or prognose diseases and/or disorders associated with the tissue(s) in which the polypeptide of the invention is expressed, including one, two, three, four, five, or more tissues disclosed in Table 1B, column 8 (Tissue Distribution Library Code).

[481] By "assaying the expression level of the gene encoding the polypeptide" is intended qualitatively or quantitatively measuring or estimating the level of the polypeptide of the invention or the level of the mRNA encoding the polypeptide of the invention in a first biological sample either directly (e.g., by determining or estimating absolute protein level or mRNA level) or relatively (e.g., by comparing to the polypeptide level or mRNA level in a second biological sample). Preferably, the polypeptide expression level or mRNA level in the first biological sample is measured or estimated and compared to a standard polypeptide level or mRNA level, the standard being taken from a second biological sample

obtained from an individual not having the disorder or being determined by averaging levels from a population of individuals not having the disorder. As will be appreciated in the art, once a standard polypeptide level or mRNA level is known, it can be used repeatedly as a standard for comparison.

[482] By "biological sample" is intended any biological sample obtained from an individual, cell line, tissue culture, or other source containing polypeptides of the invention (including portions thereof) or mRNA. As indicated, biological samples include body fluids (such as sera, plasma, urine, synovial fluid and spinal fluid) and tissue sources found to express the full length or fragments thereof of a polypeptide or mRNA. Methods for obtaining tissue biopsies and body fluids from mammals are well known in the art. Where the biological sample is to include mRNA, a tissue biopsy is the preferred source.

[483] Total cellular RNA can be isolated from a biological sample using any suitable technique such as the single-step guanidinium-thiocyanate-phenol-chloroform method described in Chomczynski and Sacchi, *Anal. Biochem.* 162:156-159 (1987). Levels of mRNA encoding the polypeptides of the invention are then assayed using any appropriate method. These include Northern blot analysis, S1 nuclease mapping, the polymerase chain reaction (PCR), reverse transcription in combination with the polymerase chain reaction (RT-PCR), and reverse transcription in combination with the ligase chain reaction (RT-LCR).

[484] The present invention also relates to diagnostic assays such as quantitative and diagnostic assays for detecting levels of polypeptides of the invention, in a biological sample (e.g., cells and tissues), including determination of normal and abnormal levels of polypeptides. Thus, for instance, a diagnostic assay in accordance with the invention for detecting over-expression of polypeptides of the invention compared to normal control tissue samples may be used to detect the presence of tumors. Assay techniques that can be used to determine levels of a polypeptide, such as a polypeptide of the present invention in a sample derived from a host are well-known to those of skill in the art. Such assay methods include radioimmunoassays, competitive-binding assays, Western Blot analysis and ELISA assays. Assaying polypeptide levels in a biological sample can occur using any art-known method.

[485] Assaying polypeptide levels in a biological sample can occur using antibody-based techniques. For example, polypeptide expression in tissues can be studied with classical immunohistological methods (Jalkanen et al., *J. Cell. Biol.* 101:976-985

(1985); Jalkanen, M., et al., J. Cell . Biol. 105:3087-3096 (1987)). Other antibody-based methods useful for detecting polypeptide gene expression include immunoassays, such as the enzyme linked immunosorbent assay (ELISA) and the radioimmunoassay (RIA). Suitable antibody assay labels are known in the art and include enzyme labels, such as, glucose oxidase, and radioisotopes, such as iodine (^{125}I , ^{121}I), carbon (^{14}C), sulfur (^{35}S), tritium (^3H), indium (^{112}In), and technetium ($^{99\text{m}}\text{Tc}$), and fluorescent labels, such as fluorescein and rhodamine, and biotin.

[486] The tissue or cell type to be analyzed will generally include those which are known, or suspected, to express the gene of interest (such as, for example, cancer). The protein isolation methods employed herein may, for example, be such as those described in Harlow and Lane (Harlow, E. and Lane, D., 1988, "Antibodies: A Laboratory Manual", Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York), which is incorporated herein by reference in its entirety. The isolated cells can be derived from cell culture or from a patient. The analysis of cells taken from culture may be a necessary step in the assessment of cells that could be used as part of a cell-based gene therapy technique or, alternatively, to test the effect of compounds on the expression of the gene.

[487] For example, antibodies, or fragments of antibodies, such as those described herein, may be used to quantitatively or qualitatively detect the presence of gene products or conserved variants or peptide fragments thereof. This can be accomplished, for example, by immunofluorescence techniques employing a fluorescently labeled antibody coupled with light microscopic, flow cytometric, or fluorimetric detection.

[488] In a preferred embodiment, antibodies, or fragments of antibodies directed to any one or all of the predicted epitope domains of the polypeptides of the invention (shown in column 7 of Table 1B) may be used to quantitatively or qualitatively detect the presence of gene products or conserved variants or peptide fragments thereof. This can be accomplished, for example, by immunofluorescence techniques employing a fluorescently labeled antibody coupled with light microscopic, flow cytometric, or fluorimetric detection.

[489] In an additional preferred embodiment, antibodies, or fragments of antibodies directed to a conformational epitope of a polypeptide of the invention may be used to quantitatively or qualitatively detect the presence of gene products or conserved variants or peptide fragments thereof. This can be accomplished, for example, by immunofluorescence techniques employing a fluorescently labeled antibody coupled with light microscopic, flow cytometric, or fluorimetric detection.

[490] The antibodies (or fragments thereof), and/or polypeptides of the present invention may, additionally, be employed histologically, as in immunofluorescence, immunoelectron microscopy or non-immunological assays, for in situ detection of gene products or conserved variants or peptide fragments thereof. In situ detection may be accomplished by removing a histological specimen from a patient, and applying thereto a labeled antibody or polypeptide of the present invention. The antibody (or fragment thereof) or polypeptide is preferably applied by overlaying the labeled antibody (or fragment) onto a biological sample. Through the use of such a procedure, it is possible to determine not only the presence of the gene product, or conserved variants or peptide fragments, or polypeptide binding, but also its distribution in the examined tissue. Using the present invention, those of ordinary skill will readily perceive that any of a wide variety of histological methods (such as staining procedures) can be modified in order to achieve such in situ detection.

[491] Immunoassays and non-immunoassays for gene products or conserved variants or peptide fragments thereof will typically comprise incubating a sample, such as a biological fluid, a tissue extract, freshly harvested cells, or lysates of cells which have been incubated in cell culture, in the presence of a detectably labeled antibody capable of binding gene products or conserved variants or peptide fragments thereof, and detecting the bound antibody by any of a number of techniques well-known in the art.

[492] The biological sample may be brought in contact with and immobilized onto a solid phase support or carrier such as nitrocellulose, or other solid support which is capable of immobilizing cells, cell particles or soluble proteins. The support may then be washed with suitable buffers followed by treatment with the detectably labeled antibody or detectable polypeptide of the invention. The solid phase support may then be washed with the buffer a second time to remove unbound antibody or polypeptide. Optionally the antibody is subsequently labeled. The amount of bound label on solid support may then be detected by conventional means.

[493] By "solid phase support or carrier" is intended any support capable of binding an antigen or an antibody. Well-known supports or carriers include glass, polystyrene, polypropylene, polyethylene, dextran, nylon, amylases, natural and modified celluloses, polyacrylamides, gabbros, and magnetite. The nature of the carrier can be either soluble to some extent or insoluble for the purposes of the present invention. The support material may have virtually any possible structural configuration so long as the coupled molecule is

capable of binding to an antigen or antibody. Thus, the support configuration may be spherical, as in a bead, or cylindrical, as in the inside surface of a test tube, or the external surface of a rod. Alternatively, the surface may be flat such as a sheet, test strip, etc. Preferred supports include polystyrene beads. Those skilled in the art will know many other suitable carriers for binding antibody or antigen, or will be able to ascertain the same by use of routine experimentation.

[494] The binding activity of a given lot of antibody or antigen polypeptide may be determined according to well known methods. Those skilled in the art will be able to determine operative and optimal assay conditions for each determination by employing routine experimentation.

[495] In addition to assaying polypeptide levels or polynucleotide levels in a biological sample obtained from an individual, polypeptide or polynucleotide can also be detected *in vivo* by imaging. For example, in one embodiment of the invention, polypeptides and/or antibodies of the invention are used to image diseased cells, such as neoplasms. In another embodiment, polynucleotides of the invention (e.g., polynucleotides complementary to all or a portion of an mRNA) and/or antibodies (e.g., antibodies directed to any one or a combination of the epitopes of a polypeptide of the invention, antibodies directed to a conformational epitope of a polypeptide of the invention, or antibodies directed to the full length polypeptide expressed on the cell surface of a mammalian cell) are used to image diseased or neoplastic cells.

[496] Antibody labels or markers for *in vivo* imaging of polypeptides of the invention include those detectable by X-radiography, NMR, MRI, CAT-scans or ESR. For X-radiography, suitable labels include radioisotopes such as barium or cesium, which emit detectable radiation but are not overtly harmful to the subject. Suitable markers for NMR and ESR include those with a detectable characteristic spin, such as deuterium, which may be incorporated into the antibody by labeling of nutrients for the relevant hybridoma. Where *in vivo* imaging is used to detect enhanced levels of polypeptides for diagnosis in humans, it may be preferable to use human antibodies or "humanized" chimeric monoclonal antibodies. Such antibodies can be produced using techniques described herein or otherwise known in the art. For example methods for producing chimeric antibodies are known in the art. See, for review, Morrison, *Science* 229:1202 (1985); Oi et al., *BioTechniques* 4:214 (1986); Cabilly et al., U.S. Patent No. 4,816,567; Taniguchi et al., EP 171496; Morrison et al., EP 173494; Neuberger et al., WO 8601533; Robinson et al., WO

8702671; Boulianne *et al.*, *Nature* 312:643 (1984); Neuberger *et al.*, *Nature* 314:268 (1985).

[497] Additionally, any polypeptides of the invention whose presence can be detected, can be administered. For example, polypeptides of the invention labeled with a radio-opaque or other appropriate compound can be administered and visualized *in vivo*, as discussed, above for labeled antibodies. Further, such polypeptides can be utilized for *in vitro* diagnostic procedures.

[498] A polypeptide-specific antibody or antibody fragment which has been labeled with an appropriate detectable imaging moiety, such as a radioisotope (for example, ^{131}I , ^{112}In , $^{99\text{m}}\text{Tc}$), a radio-opaque substance, or a material detectable by nuclear magnetic resonance, is introduced (for example, parenterally, subcutaneously or intraperitoneally) into the mammal to be examined for a disorder. It will be understood in the art that the size of the subject and the imaging system used will determine the quantity of imaging moiety needed to produce diagnostic images. In the case of a radioisotope moiety, for a human subject, the quantity of radioactivity injected will normally range from about 5 to 20 millicuries of $^{99\text{m}}\text{Tc}$. The labeled antibody or antibody fragment will then preferentially accumulate at the location of cells which contain the antigenic protein. *In vivo* tumor imaging is described in S.W. Burchiel *et al.*, "Immunopharmacokinetics of Radiolabeled Antibodies and Their Fragments" (Chapter 13 in *Tumor Imaging: The Radiochemical Detection of Cancer*, S.W. Burchiel and B. A. Rhodes, eds., Masson Publishing Inc. (1982)).

[499] With respect to antibodies, one of the ways in which an antibody of the present invention can be detectably labeled is by linking the same to a reporter enzyme and using the linked product in an enzyme immunoassay (EIA) (Voller, A., "The Enzyme Linked Immunosorbent Assay (ELISA)", 1978, Diagnostic Horizons 2:1-7, Microbiological Associates Quarterly Publication, Walkersville, MD); Voller *et al.*, *J. Clin. Pathol.* 31:507-520 (1978); Butler, J.E., *Meth. Enzymol.* 73:482-523 (1981); Maggio, E. (ed.), 1980, Enzyme Immunoassay, CRC Press, Boca Raton, FL.; Ishikawa, E. *et al.*, (eds.), 1981, Enzyme Immunoassay, Kigaku Shoin, Tokyo). The reporter enzyme which is bound to the antibody will react with an appropriate substrate, preferably a chromogenic substrate, in such a manner as to produce a chemical moiety which can be detected, for example, by spectrophotometric, fluorimetric or by visual means. Reporter enzymes which can be used to detectably label the antibody include, but are not limited to, malate dehydrogenase,

staphylococcal nuclease, delta-5-steroid isomerase, yeast alcohol dehydrogenase, alpha-glycerophosphate, dehydrogenase, triose phosphate isomerase, horseradish peroxidase, alkaline phosphatase, asparaginase, glucose oxidase, beta-galactosidase, ribonuclease, urease, catalase, glucose-6-phosphate dehydrogenase, glucoamylase and acetylcholinesterase. Additionally, the detection can be accomplished by colorimetric methods which employ a chromogenic substrate for the reporter enzyme. Detection may also be accomplished by visual comparison of the extent of enzymatic reaction of a substrate in comparison with similarly prepared standards.

[500] Detection may also be accomplished using any of a variety of other immunoassays. For example, by radioactively labeling the antibodies or antibody fragments, it is possible to detect polypeptides through the use of a radioimmunoassay (RIA) (see, for example, Weintraub, B., Principles of Radioimmunoassays, Seventh Training Course on Radioligand Assay Techniques, The Endocrine Society, March, 1986, which is incorporated by reference herein). The radioactive isotope can be detected by means including, but not limited to, a gamma counter, a scintillation counter, or autoradiography.

[501] It is also possible to label the antibody with a fluorescent compound. When the fluorescently labeled antibody is exposed to light of the proper wave length, its presence can then be detected due to fluorescence. Among the most commonly used fluorescent labeling compounds are fluorescein isothiocyanate, rhodamine, phycoerythrin, phycocyanin, allophycocyanin, ophthaldehyde and fluorescamine.

[502] The antibody can also be detectably labeled using fluorescence emitting metals such as ^{152}Eu , or others of the lanthanide series. These metals can be attached to the antibody using such metal chelating groups as diethylenetriaminepentacetic acid (DTPA) or ethylenediaminetetraacetic acid (EDTA).

[503] The antibody also can be detectably labeled by coupling it to a chemiluminescent compound. The presence of the chemiluminescent-tagged antibody is then determined by detecting the presence of luminescence that arises during the course of a chemical reaction. Examples of particularly useful chemiluminescent labeling compounds are luminol, isoluminol, theromatic acridinium ester, imidazole, acridinium salt and oxalate ester.

[504] Likewise, a bioluminescent compound may be used to label the antibody of the present invention. Bioluminescence is a type of chemiluminescence found in biological systems in, which a catalytic protein increases the efficiency of the chemiluminescent

reaction. The presence of a bioluminescent protein is determined by detecting the presence of luminescence. Important bioluminescent compounds for purposes of labeling are luciferin, luciferase and aequorin.

Methods for Detecting Diseases

[505] In general, a disease may be detected in a patient based on the presence of one or more proteins of the invention and/or polynucleotides encoding such proteins in a biological sample (for example, blood, sera, urine, and/or tumor biopsies) obtained from the patient. In other words, such proteins may be used as markers to indicate the presence or absence of a disease or disorder, including cancer and/or as described elsewhere herein. In addition, such proteins may be useful for the detection of other diseases and cancers. The binding agents provided herein generally permit detection of the level of antigen that binds to the agent in the biological sample. Polynucleotide primers and probes may be used to detect the level of mRNA encoding polypeptides of the invention, which is also indicative of the presence or absence of a disease or disorder, including cancer. In general, polypeptides of the invention should be present at a level that is at least three fold higher in diseased tissue than in normal tissue.

[506] There are a variety of assay formats known to those of ordinary skill in the art for using a binding agent to detect polypeptide markers in a sample. See, e.g., Harlow and Lane, *supra*. In general, the presence or absence of a disease in a patient may be determined by (a) contacting a biological sample obtained from a patient with a binding agent; (b) detecting in the sample a level of polypeptide that binds to the binding agent; and (c) comparing the level of polypeptide with a predetermined cut-off value.

[507] In a preferred embodiment, the assay involves the use of a binding agent(s) immobilized on a solid support to bind to and remove the polypeptide of the invention from the remainder of the sample. The bound polypeptide may then be detected using a detection reagent that contains a reporter group and specifically binds to the binding agent/polypeptide complex. Such detection reagents may comprise, for example, a binding agent that specifically binds to the polypeptide or an antibody or other agent that specifically binds to the binding agent, such as an anti-immunoglobulin, protein G, protein A or a lectin. Alternatively, a competitive assay may be utilized, in which a polypeptide is labeled with a reporter group and allowed to bind to the immobilized binding agent after incubation of the binding agent with the sample. The extent to which components of the

sample inhibit the binding of the labeled polypeptide to the binding agent is indicative of the reactivity of the sample with the immobilized binding agent. Suitable polypeptides for use within such assays include polypeptides of the invention and portions thereof, or antibodies, to which the binding agent binds, as described above.

[508] The solid support may be any material known to those of skill in the art to which polypeptides of the invention may be attached. For example, the solid support may be a test well in a microtiter plate or a nitrocellulose or other suitable membrane. Alternatively, the support may be a bead or disc, such as glass fiberglass, latex or a plastic material such as polystyrene or polyvinylchloride. The support may also be a magnetic particle or a fiber optic sensor, such as those disclosed, for example, in U.S. Patent No. 5,359,681. The binding agent may be immobilized on the solid support using a variety of techniques known to those of skill in the art, which are amply described in the patent and scientific literature. In the context of the present invention, the term "immobilization" refers to both noncovalent association, such as adsorption, and covalent attachment (which may be a direct linkage between the agent and functional groups on the support or may be a linkage by way of a cross-linking agent). Immobilization by adsorption to a well in a microtiter plate or to a membrane is preferred. In such cases, adsorption may be achieved by contacting the binding agent, in a suitable buffer, with the solid support for the suitable amount of time. The contact time varies with temperature, but is typically between about 1 hour and about 1 day. In general, contacting a well of plastic microtiter plate (such as polystyrene or polyvinylchloride) with an amount of binding agent ranging from about 10 ng to about 10 ug, and preferably about 100 ng to about 1 ug, is sufficient to immobilize an adequate amount of binding agent.

[509] Covalent attachment of binding agent to a solid support may generally be achieved by first reacting the support with a bifunctional reagent that will react with both the support and a functional group, such as a hydroxyl or amino group, on the binding agent. For example, the binding agent may be covalently attached to supports having an appropriate polymer coating using benzoquinone or by condensation of an aldehyde group on the support with an amine and an active hydrogen on the binding partner (see, e.g., Pierce Immunotechnology Catalog and Handbook, 1991, at A12-A13).

Gene Therapy Methods

[510] Also encompassed by the invention are gene therapy methods for treating or preventing disorders, diseases and conditions. The gene therapy methods relate to the introduction of nucleic acid (DNA, RNA and antisense DNA or RNA) sequences into an animal to achieve expression of the polypeptide of the present invention. This method requires a polynucleotide which codes for a polypeptide of the present invention operatively linked to a promoter and any other genetic elements necessary for the expression of the polypeptide by the target tissue. Such gene therapy and delivery techniques are known in the art, see, for example, WO90/11092, which is herein incorporated by reference.

[511] Thus, for example, cells from a patient may be engineered with a polynucleotide (DNA or RNA) comprising a promoter operably linked to a polynucleotide of the present invention *ex vivo*, with the engineered cells then being provided to a patient to be treated with the polypeptide of the present invention. Such methods are well-known in the art. For example, see Belldegrün, A., et al., J. Natl. Cancer Inst. 85: 207-216 (1993); Ferrantini, M. et al., Cancer Research 53: 1107-1112 (1993); Ferrantini, M. et al., J. Immunology 153: 4604-4615 (1994); Kaido, T., et al., Int. J. Cancer 60: 221-229 (1995); Ogura, H., et al., Cancer Research 50: 5102-5106 (1990); Santodonato, L., et al., Human Gene Therapy 7:1-10 (1996); Santodonato, L., et al., Gene Therapy 4:1246-1255 (1997); and Zhang, J.-F. et al., Cancer Gene Therapy 3: 31-38 (1996)), which are herein incorporated by reference. In one embodiment, the cells which are engineered are arterial cells. The arterial cells may be reintroduced into the patient through direct injection to the artery, the tissues surrounding the artery, or through catheter injection.

[512] As discussed in more detail below, the polynucleotide constructs can be delivered by any method that delivers injectable materials to the cells of an animal, such as, injection into the interstitial space of tissues (heart, muscle, skin, lung, liver, and the like). The polynucleotide constructs may be delivered in a pharmaceutically acceptable liquid or aqueous carrier.

[513] In one embodiment, the polynucleotide of the present invention is delivered as a naked polynucleotide. The term "naked" polynucleotide, DNA or RNA refers to sequences that are free from any delivery vehicle that acts to assist, promote or facilitate entry into the cell, including viral sequences, viral particles, liposome formulations, lipofectin or precipitating agents and the like. However, the polynucleotide of the present invention can also be delivered in liposome formulations and lipofectin formulations and the like can be

prepared by methods well known to those skilled in the art. Such methods are described, for example, in U.S. Patent Nos. 5,593,972, 5,589,466, and 5,580,859, which are herein incorporated by reference.

[514] The polynucleotide vector constructs used in the gene therapy method are preferably constructs that will not integrate into the host genome nor will they contain sequences that allow for replication. Appropriate vectors include pWLNEO, pSV2CAT, pOG44, pXT1 and pSG available from Stratagene; pSVK3, pBPV, pMSG and pSVL available from Pharmacia; and pEF1/V5, pcDNA3.1, and pRc/CMV2 available from Invitrogen. Other suitable vectors will be readily apparent to the skilled artisan.

[515] Any strong promoter known to those skilled in the art can be used for driving the expression of the polynucleotide sequence. Suitable promoters include adenoviral promoters, such as the adenoviral major late promoter; or heterologous promoters, such as the cytomegalovirus (CMV) promoter; the respiratory syncytial virus (RSV) promoter; inducible promoters, such as the MMT promoter, the metallothionein promoter; heat shock promoters; the albumin promoter; the ApoAI promoter; human globin promoters; viral thymidine kinase promoters, such as the Herpes Simplex thymidine kinase promoter; retroviral LTRs; the b-actin promoter; and human growth hormone promoters. The promoter also may be the native promoter for the polynucleotide of the present invention.

[516] Unlike other gene therapy techniques, one major advantage of introducing naked nucleic acid sequences into target cells is the transitory nature of the polynucleotide synthesis in the cells. Studies have shown that non-replicating DNA sequences can be introduced into cells to provide production of the desired polypeptide for periods of up to six months.

[517] The polynucleotide construct can be delivered to the interstitial space of tissues within the an animal, including of muscle, skin, brain, lung, liver, spleen, bone marrow, thymus, heart, lymph, blood, bone, cartilage, pancreas, kidney, gall bladder, stomach, intestine, testis, ovary, uterus, rectum, nervous system, eye, gland, and connective tissue. Interstitial space of the tissues comprises the intercellular, fluid, mucopolysaccharide matrix among the reticular fibers of organ tissues, elastic fibers in the walls of vessels or chambers, collagen fibers of fibrous tissues, or that same matrix within connective tissue ensheathing muscle cells or in the lacunae of bone. It is similarly the space occupied by the plasma of the circulation and the lymph fluid of the lymphatic channels. Delivery to the interstitial space of muscle tissue is preferred for the reasons discussed below. They may be conveniently

delivered by injection into the tissues comprising these cells. They are preferably delivered to and expressed in persistent, non-dividing cells which are differentiated, although delivery and expression may be achieved in non-differentiated or less completely differentiated cells, such as, for example, stem cells of blood or skin fibroblasts. *In vivo* muscle cells are particularly competent in their ability to take up and express polynucleotides.

[518] For the naked nucleic acid sequence injection, an effective dosage amount of DNA or RNA will be in the range of from about 0.05 mg/kg body weight to about 50 mg/kg body weight. Preferably the dosage will be from about 0.005 mg/kg to about 20 mg/kg and more preferably from about 0.05 mg/kg to about 5 mg/kg. Of course, as the artisan of ordinary skill will appreciate, this dosage will vary according to the tissue site of injection. The appropriate and effective dosage of nucleic acid sequence can readily be determined by those of ordinary skill in the art and may depend on the condition being treated and the route of administration.

[519] The preferred route of administration is by the parenteral route of injection into the interstitial space of tissues. However, other parenteral routes may also be used, such as, inhalation of an aerosol formulation particularly for delivery to lungs or bronchial tissues, throat or mucous membranes of the nose. In addition, naked DNA constructs can be delivered to arteries during angioplasty by the catheter used in the procedure.

[520] The naked polynucleotides are delivered by any method known in the art, including, but not limited to, direct needle injection at the delivery site, intravenous injection, topical administration, catheter infusion, and so-called "gene guns". These delivery methods are known in the art.

[521] The constructs may also be delivered with delivery vehicles such as viral sequences, viral particles, liposome formulations, lipofectin, precipitating agents, etc. Such methods of delivery are known in the art.

[522] In certain embodiments, the polynucleotide constructs are complexed in a liposome preparation. Liposomal preparations for use in the instant invention include cationic (positively charged), anionic (negatively charged) and neutral preparations. However, cationic liposomes are particularly preferred because a tight charge complex can be formed between the cationic liposome and the polyanionic nucleic acid. Cationic liposomes have been shown to mediate intracellular delivery of plasmid DNA (Felgner et al., Proc. Natl. Acad. Sci. USA (1987) 84:7413-7416, which is herein incorporated by reference); mRNA (Malone et al., Proc. Natl. Acad. Sci. USA (1989) 86:6077-6081, which

is herein incorporated by reference); and purified transcription factors (Debs et al., J. Biol. Chem. (1990) 265:10189-10192, which is herein incorporated by reference), in functional form.

[523] Cationic liposomes are readily available. For example, N[1-2,3-dioleoyloxy)propyl]-N,N,N-triethylammonium (DOTMA) liposomes are particularly useful and are available under the trademark Lipofectin, from GIBCO BRL, Grand Island, N.Y. (See, also, Felgner et al., Proc. Natl Acad. Sci. USA (1987) 84:7413-7416, which is herein incorporated by reference). Other commercially available liposomes include transfectace (DDAB/DOPE) and DOTAP/DOPE (Boehringer).

[524] Other cationic liposomes can be prepared from readily available materials using techniques well known in the art. See, e.g. PCT Publication No. WO 90/11092 (which is herein incorporated by reference) for a description of the synthesis of DOTAP (1,2-bis(oleoyloxy)-3-(trimethylammonio)propane) liposomes. Preparation of DOTMA liposomes is explained in the literature, see, e.g., P. Felgner et al., Proc. Natl. Acad. Sci. USA 84:7413-7417, which is herein incorporated by reference. Similar methods can be used to prepare liposomes from other cationic lipid materials.

[525] Similarly, anionic and neutral liposomes are readily available, such as from Avanti Polar Lipids (Birmingham, Ala.), or can be easily prepared using readily available materials. Such materials include phosphatidyl, choline, cholesterol, phosphatidyl ethanolamine, dioleoylphosphatidyl choline (DOPC), dioleoylphosphatidyl glycerol (DOPG), dioleoylphosphatidyl ethanolamine (DOPE), among others. These materials can also be mixed with the DOTMA and DOTAP starting materials in appropriate ratios. Methods for making liposomes using these materials are well known in the art.

[526] For example, commercially dioleoylphosphatidyl choline (DOPC), dioleoylphosphatidyl glycerol (DOPG), and dioleoylphosphatidyl ethanolamine (DOPE) can be used in various combinations to make conventional liposomes, with or without the addition of cholesterol. Thus, for example, DOPG/DOPC vesicles can be prepared by drying 50 mg each of DOPG and DOPC under a stream of nitrogen gas into a sonication vial. The sample is placed under a vacuum pump overnight and is hydrated the following day with deionized water. The sample is then sonicated for 2 hours in a capped vial, using a Heat Systems model 350 sonicator equipped with an inverted cup (bath type) probe at the maximum setting while the bath is circulated at 15EC. Alternatively, negatively charged vesicles can be prepared without sonication to produce multilamellar vesicles or by

09950083-091201

extrusion through nucleopore membranes to produce unilamellar vesicles of discrete size. Other methods are known and available to those of skill in the art.

[527] The liposomes can comprise multilamellar vesicles (MLVs), small unilamellar vesicles (SUVs), or large unilamellar vesicles (LUVs), with SUVs being preferred. The various liposome-nucleic acid complexes are prepared using methods well known in the art. See, e.g., Straubinger et al., *Methods of Immunology* (1983), 101:512-527, which is herein incorporated by reference. For example, MLVs containing nucleic acid can be prepared by depositing a thin film of phospholipid on the walls of a glass tube and subsequently hydrating with a solution of the material to be encapsulated. SUVs are prepared by extended sonication of MLVs to produce a homogeneous population of unilamellar liposomes. The material to be entrapped is added to a suspension of preformed MLVs and then sonicated. When using liposomes containing cationic lipids, the dried lipid film is resuspended in an appropriate solution such as sterile water or an isotonic buffer solution such as 10 mM Tris/NaCl, sonicated, and then the preformed liposomes are mixed directly with the DNA. The liposome and DNA form a very stable complex due to binding of the positively charged liposomes to the cationic DNA. SUVs find use with small nucleic acid fragments. LUVs are prepared by a number of methods, well known in the art. Commonly used methods include Ca^{2+} -EDTA chelation (Papahadjopoulos et al., *Biochim. Biophys. Acta* (1975) 394:483; Wilson et al., *Cell* 17:77 (1979)); ether injection (Deamer, D. and Bangham, A., *Biochim. Biophys. Acta* 443:629 (1976); Ostro et al., *Biochem. Biophys. Res. Commun.* 76:836 (1977); Fraley et al., *Proc. Natl. Acad. Sci. USA* 76:3348 (1979)); detergent dialysis (Enoch, H. and Strittmatter, P., *Proc. Natl. Acad. Sci. USA* 76:145 (1979)); and reverse-phase evaporation (REV) (Fraley et al., *J. Biol. Chem.* 255:10431 (1980); Szoka, F. and Papahadjopoulos, D., *Proc. Natl. Acad. Sci. USA* 75:145 (1978); Schaefer-Ridder et al., *Science* 215:166 (1982)), which are herein incorporated by reference.

[528] Generally, the ratio of DNA to liposomes will be from about 10:1 to about 1:10. Preferably, the ration will be from about 5:1 to about 1:5. More preferably, the ration will be about 3:1 to about 1:3. Still more preferably, the ratio will be about 1:1.

[529] U.S. Patent No. 5,676,954 (which is herein incorporated by reference) reports on the injection of genetic material, complexed with cationic liposomes carriers, into mice. U.S. Patent Nos. 4,897,355, 4,946,787, 5,049,386, 5,459,127, 5,589,466, 5,693,622, 5,580,859, 5,703,055, and international publication no. WO 94/9469 (which are herein

incorporated by reference) provide cationic lipids for use in transfecting DNA into cells and mammals. U.S. Patent Nos. 5,589,466, 5,693,622, 5,580,859, 5,703,055, and international publication no. WO 94/9469 provide methods for delivering DNA-cationic lipid complexes to mammals.

[530] In certain embodiments, cells are engineered, *ex vivo* or *in vivo*, using a retroviral particle containing RNA which comprises a sequence encoding a polypeptide of the present invention. Retroviruses from which the retroviral plasmid vectors may be derived include, but are not limited to, Moloney Murine Leukemia Virus, spleen necrosis virus, Rous sarcoma Virus, Harvey Sarcoma Virus, avian leukosis virus, gibbon ape leukemia virus, human immunodeficiency virus, Myeloproliferative Sarcoma Virus, and mammary tumor virus.

[531] The retroviral plasmid vector is employed to transduce packaging cell lines to form producer cell lines. Examples of packaging cells which may be transfected include, but are not limited to, the PE501, PA317, R-2, R-AM, PA12, T19-14X, VT-19-17-H2, RCRE, RCRIP, GP+E-86, GP+envAm12, and DAN cell lines as described in Miller, Human Gene Therapy 1:5-14 (1990), which is incorporated herein by reference in its entirety. The vector may transduce the packaging cells through any means known in the art. Such means include, but are not limited to, electroporation, the use of liposomes, and CaPO₄ precipitation. In one alternative, the retroviral plasmid vector may be encapsulated into a liposome, or coupled to a lipid, and then administered to a host.

[532] The producer cell line generates infectious retroviral vector particles which include polynucleotide encoding a polypeptide of the present invention. Such retroviral vector particles then may be employed, to transduce eukaryotic cells, either *in vitro* or *in vivo*. The transduced eukaryotic cells will express a polypeptide of the present invention.

[533] In certain other embodiments, cells are engineered, *ex vivo* or *in vivo*, with polynucleotide contained in an adenovirus vector. Adenovirus can be manipulated such that it encodes and expresses a polypeptide of the present invention, and at the same time is inactivated in terms of its ability to replicate in a normal lytic viral life cycle. Adenovirus expression is achieved without integration of the viral DNA into the host cell chromosome, thereby alleviating concerns about insertional mutagenesis. Furthermore, adenoviruses have been used as live enteric vaccines for many years with an excellent safety profile (Schwartz et al. Am. Rev. Respir. Dis.109:233-238 (1974)). Finally, adenovirus mediated gene transfer has been demonstrated in a number of instances including transfer of

alpha-1-antitrypsin and CFTR to the lungs of cotton rats (Rosenfeld, M. A. et al. (1991) Science 252:431-434; Rosenfeld et al., (1992) Cell 68:143-155). Furthermore, extensive studies to attempt to establish adenovirus as a causative agent in human cancer were uniformly negative (Green, M. et al. (1979) Proc. Natl. Acad. Sci. USA 76:6606).

[534] Suitable adenoviral vectors useful in the present invention are described, for example, in Kozarsky and Wilson, Curr. Opin. Genet. Devel. 3:499-503 (1993); Rosenfeld et al., Cell 68:143-155 (1992); Engelhardt et al., Human Genet. Ther. 4:759-769 (1993); Yang et al., Nature Genet. 7:362-369 (1994); Wilson et al., Nature 365:691-692 (1993); and U.S. Patent No. 5,652,224, which are herein incorporated by reference. For example, the adenovirus vector Ad2 is useful and can be grown in human 293 cells. These cells contain the E1 region of adenovirus and constitutively express Ela and Elb, which complement the defective adenoviruses by providing the products of the genes deleted from the vector. In addition to Ad2, other varieties of adenovirus (e.g., Ad3, Ad5, and Ad7) are also useful in the present invention.

[535] Preferably, the adenoviruses used in the present invention are replication deficient. Replication deficient adenoviruses require the aid of a helper virus and/or packaging cell line to form infectious particles. The resulting virus is capable of infecting cells and can express a polynucleotide of interest which is operably linked to a promoter, but cannot replicate in most cells. Replication deficient adenoviruses may be deleted in one or more of all or a portion of the following genes: E1a, E1b, E3, E4, E2a, or L1 through L5.

[536] In certain other embodiments, the cells are engineered, ex vivo or *in vivo*, using an adeno-associated virus (AAV). AAVs are naturally occurring defective viruses that require helper viruses to produce infectious particles (Muzyczka, N., Curr. Topics in Microbiol. Immunol. 158:97 (1992)). It is also one of the few viruses that may integrate its DNA into non-dividing cells. Vectors containing as little as 300 base pairs of AAV can be packaged and can integrate, but space for exogenous DNA is limited to about 4.5 kb. Methods for producing and using such AAVs are known in the art. See, for example, U.S. Patent Nos. 5,139,941, 5,173,414, 5,354,678, 5,436,146, 5,474,935, 5,478,745, and 5,589,377.

[537] For example, an appropriate AAV vector for use in the present invention will include all the sequences necessary for DNA replication, encapsidation, and host-cell integration. The polynucleotide construct is inserted into the AAV vector using standard cloning methods, such as those found in Sambrook et al., Molecular Cloning: A Laboratory

Manual, Cold Spring Harbor Press (1989). The recombinant AAV vector is then transfected into packaging cells which are infected with a helper virus, using any standard technique, including lipofection, electroporation, calcium phosphate precipitation, etc. Appropriate helper viruses include adenoviruses, cytomegaloviruses, vaccinia viruses, or herpes viruses. Once the packaging cells are transfected and infected, they will produce infectious AAV viral particles which contain the polynucleotide construct. These viral particles are then used to transduce eukaryotic cells, either *ex vivo* or *in vivo*. The transduced cells will contain the polynucleotide construct integrated into its genome, and will express a polypeptide of the invention.

[538] Another method of gene therapy involves operably associating heterologous control regions and endogenous polynucleotide sequences (e.g. encoding a polypeptide of the present invention) via homologous recombination (see, e.g., U.S. Patent No. 5,641,670, issued June 24, 1997; International Publication No. WO 96/29411, published September 26, 1996; International Publication No. WO 94/12650, published August 4, 1994; Koller et al., Proc. Natl. Acad. Sci. USA 86:8932-8935 (1989); and Zijlstra et al., Nature 342:435-438 (1989), which are herein incorporated by reference. This method involves the activation of a gene which is present in the target cells, but which is not normally expressed in the cells, or is expressed at a lower level than desired.

[539] Polynucleotide constructs are made, using standard techniques known in the art, which contain the promoter with targeting sequences flanking the promoter. Suitable promoters are described herein. The targeting sequence is sufficiently complementary to an endogenous sequence to permit homologous recombination of the promoter-targeting sequence with the endogenous sequence. The targeting sequence will be sufficiently near the 5' end of the desired endogenous polynucleotide sequence so the promoter will be operably linked to the endogenous sequence upon homologous recombination.

[540] The promoter and the targeting sequences can be amplified using PCR. Preferably, the amplified promoter contains distinct restriction enzyme sites on the 5' and 3' ends. Preferably, the 3' end of the first targeting sequence contains the same restriction enzyme site as the 5' end of the amplified promoter and the 5' end of the second targeting sequence contains the same restriction site as the 3' end of the amplified promoter. The amplified promoter and targeting sequences are digested and ligated together.

[541] The promoter-targeting sequence construct is delivered to the cells, either as naked polynucleotide, or in conjunction with transfection-facilitating agents, such as

liposomes, viral sequences, viral particles, whole viruses, lipofection, precipitating agents, etc., described in more detail above. The P promoter-targeting sequence can be delivered by any method, included direct needle injection, intravenous injection, topical administration, catheter infusion, particle accelerators, etc. The methods are described in more detail below.

[542] The promoter-targeting sequence construct is taken up by cells. Homologous recombination between the construct and the endogenous sequence takes place, such that an endogenous sequence is placed under the control of the promoter. The promoter then drives the expression of the endogenous sequence.

[543] The polynucleotide encoding a polypeptide of the present invention may contain a secretory signal sequence that facilitates secretion of the protein. Typically, the signal sequence is positioned in the coding region of the polynucleotide to be expressed towards or at the 5' end of the coding region. The signal sequence may be homologous or heterologous to the polynucleotide of interest and may be homologous or heterologous to the cells to be transfected. Additionally, the signal sequence may be chemically synthesized using methods known in the art.

[544] Any mode of administration of any of the above-described polynucleotides constructs can be used so long as the mode results in the expression of one or more molecules in an amount sufficient to provide a therapeutic effect. This includes direct needle injection, systemic injection, catheter infusion, biolistic injectors, particle accelerators (i.e., "gene guns"), gelfoam sponge depots, other commercially available depot materials, osmotic pumps (e.g., Alza minipumps), oral or suppositorial solid (tablet or pill) pharmaceutical formulations, and decanting or topical applications during surgery. For example, direct injection of naked calcium phosphate-precipitated plasmid into rat liver and rat spleen or a protein-coated plasmid into the portal vein has resulted in gene expression of the foreign gene in the rat livers (Kaneda et al., Science 243:375 (1989)).

[545] A preferred method of local administration is by direct injection. Preferably, a recombinant molecule of the present invention complexed with a delivery vehicle is administered by direct injection into or locally within the area of arteries. Administration of a composition locally within the area of arteries refers to injecting the composition centimeters and preferably, millimeters within arteries.

[546] Another method of local administration is to contact a polynucleotide construct of the present invention in or around a surgical wound. For example, a patient can undergo

0950083-091201

surgery and the polynucleotide construct can be coated on the surface of tissue inside the wound or the construct can be injected into areas of tissue inside the wound.

[547] Therapeutic compositions useful in systemic administration, include recombinant molecules of the present invention complexed to a targeted delivery vehicle of the present invention. Suitable delivery vehicles for use with systemic administration comprise liposomes comprising ligands for targeting the vehicle to a particular site. In specific embodiments, suitable delivery vehicles for use with systemic administration comprise liposomes comprising polypeptides of the invention for targeting the vehicle to a particular site.

[548] Preferred methods of systemic administration, include intravenous injection, aerosol, oral and percutaneous (topical) delivery. Intravenous injections can be performed using methods standard in the art. Aerosol delivery can also be performed using methods standard in the art (see, for example, Stribling et al., Proc. Natl. Acad. Sci. USA 189:11277-11281, 1992, which is incorporated herein by reference). Oral delivery can be performed by complexing a polynucleotide construct of the present invention to a carrier capable of withstanding degradation by digestive enzymes in the gut of an animal. Examples of such carriers, include plastic capsules or tablets, such as those known in the art. Topical delivery can be performed by mixing a polynucleotide construct of the present invention with a lipophilic reagent (e.g., DMSO) that is capable of passing into the skin.

[549] Determining an effective amount of substance to be delivered can depend upon a number of factors including, for example, the chemical structure and biological activity of the substance, the age and weight of the animal, the precise condition requiring treatment and its severity, and the route of administration. The frequency of treatments depends upon a number of factors, such as the amount of polynucleotide constructs administered per dose, as well as the health and history of the subject. The precise amount, number of doses, and timing of doses will be determined by the attending physician or veterinarian.

[550] Therapeutic compositions of the present invention can be administered to any animal, preferably to mammals and birds. Preferred mammals include humans, dogs, cats, mice, rats, rabbits sheep, cattle, horses and pigs, with humans being particularly preferred.

Biological Activities

[551] Polynucleotides or polypeptides, or agonists or antagonists of the present invention, can be used in assays to test for one or more biological activities. If these

polynucleotides or polypeptides, or agonists or antagonists of the present invention, do exhibit activity in a particular assay, it is likely that these molecules may be involved in the diseases associated with the biological activity. Thus, the polynucleotides and polypeptides, and agonists or antagonists could be used to treat the associated disease.

[552] Members of the secreted family of proteins are believed to be involved in biological activities associated with, for example, cellular signaling. Accordingly, compositions of the invention (including polynucleotides, polypeptides and antibodies of the invention, and fragments and variants thereof) may be used in diagnosis, prognosis, prevention and/or treatment of diseases and/or disorders associated with aberrant activity of secreted polypeptides.

[553] In preferred embodiments, compositions of the invention (including polynucleotides, polypeptides and antibodies of the invention, and fragments and variants thereof) may be used in the diagnosis, prognosis, prevention and/or treatment of diseases and/or disorders relating to diseases and disorders of the endocrine system, the nervous system (See, for example, "Neurological Disorders" section below), and the immune system (See, for example, "Immune Activity" section below).

[554] In certain embodiments, a polypeptide of the invention, or polynucleotides, antibodies, agonists, or antagonists corresponding to that polypeptide, may be used to diagnose and/or prognose diseases and/or disorders associated with the tissue(s) in which the polypeptide of the invention is expressed including one, two, three, four, five, or more tissues disclosed in Table 1B, column 8 (Tissue Distribution Library Code).

[555] Thus, polynucleotides, translation products and antibodies of the invention are useful in the diagnosis, detection and/or treatment of diseases and/or disorders associated with activities that include, but are not limited to, prohormone activation, neurotransmitter activity, cellular signaling, cellular proliferation, cellular differentiation, and cell migration.

[556] More generally, polynucleotides, translation products and antibodies corresponding to this gene may be useful for the diagnosis, prognosis, prevention and/or treatment of diseases and/or disorders associated with the following systems.

Immune Activity

[557] Polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be useful in treating, preventing, diagnosing and/or prognosing

immunodeficiency (CVID), common variable immunodeficiency (CVI) (acquired), and transient hypogammaglobulinemia of infancy.

[560] In specific embodiments, ataxia-telangiectasia or conditions associated with ataxia-telangiectasia are treated, prevented, diagnosed, and/or prognosing using the polypeptides or polynucleotides of the invention, and/or agonists or antagonists thereof.

[561] Examples of congenital immunodeficiencies in which T cell and/or B cell function and/or number is decreased include, but are not limited to: DiGeorge anomaly, severe combined immunodeficiencies (SCID) (including, but not limited to, X-linked SCID, autosomal recessive SCID, adenosine deaminase deficiency, purine nucleoside phosphorylase (PNP) deficiency, Class II MHC deficiency (Bare lymphocyte syndrome), Wiskott-Aldrich syndrome, and ataxia telangiectasia), thymic hypoplasia, third and fourth pharyngeal pouch syndrome, 22q11.2 deletion, chronic mucocutaneous candidiasis, natural killer cell deficiency (NK), idiopathic CD4+ T-lymphocytopenia, immunodeficiency with predominant T cell defect (unspecified), and unspecified immunodeficiency of cell mediated immunity.

[562] In specific embodiments, DiGeorge anomaly or conditions associated with DiGeorge anomaly are treated, prevented, diagnosed, and/or prognosed using polypeptides or polynucleotides of the invention, or antagonists or agonists thereof.

[563] Other immunodeficiencies that may be treated, prevented, diagnosed, and/or prognosed using polypeptides or polynucleotides of the invention, and/or agonists or antagonists thereof, include, but are not limited to, chronic granulomatous disease, Chédiak-Higashi syndrome, myeloperoxidase deficiency, leukocyte glucose-6-phosphate dehydrogenase deficiency, X-linked lymphoproliferative syndrome (XLP), leukocyte adhesion deficiency, complement component deficiencies (including C1, C2, C3, C4, C5, C6, C7, C8 and/or C9 deficiencies), reticular dysgenesis, thymic aplasia, immunodeficiency with thymoma, severe congenital leukopenia, dysplasia with immunodeficiency, neonatal neutropenia, short limbed dwarfism, and Nezelof syndrome-combined immunodeficiency with Igs.

[564] In a preferred embodiment, the immunodeficiencies and/or conditions associated with the immunodeficiencies recited above are treated, prevented, diagnosed and/or

prognosed using polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention.

[565] In a preferred embodiment polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention could be used as an agent to boost immunoresponsiveness among immunodeficient individuals. In specific embodiments, polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention could be used as an agent to boost immunoresponsiveness among B cell and/or T cell immunodeficient individuals.

[566] The polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be useful in treating, preventing, diagnosing and/or prognosing autoimmune disorders. Many autoimmune disorders result from inappropriate recognition of self as foreign material by immune cells. This inappropriate recognition results in an immune response leading to the destruction of the host tissue. Therefore, the administration of polynucleotides and polypeptides of the invention that can inhibit an immune response, particularly the proliferation, differentiation, or chemotaxis of T-cells, may be an effective therapy in preventing autoimmune disorders.

[567] Autoimmune diseases or disorders that may be treated, prevented, diagnosed and/or prognosed by polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention include, but are not limited to, one or more of the following: systemic lupus erythematosus, rheumatoid arthritis, ankylosing spondylitis, multiple sclerosis, autoimmune thyroiditis, Hashimoto's thyroiditis, autoimmune hemolytic anemia, hemolytic anemia, thrombocytopenia, autoimmune thrombocytopenia purpura, autoimmune neonatal thrombocytopenia, idiopathic thrombocytopenia purpura, purpura (e.g., Henloch-Scoenlein purpura), autoimmunocytopenia, Goodpasture's syndrome, Pemphigus vulgaris, myasthenia gravis, Grave's disease (hyperthyroidism), and insulin-resistant diabetes mellitus.

[568] Additional disorders that are likely to have an autoimmune component that may be treated, prevented, and/or diagnosed with the compositions of the invention include, but are not limited to, type II collagen-induced arthritis, antiphospholipid syndrome, dermatitis, allergic encephalomyelitis, myocarditis, relapsing polychondritis, rheumatic heart disease, neuritis, uveitis ophthalmia, polyendocrinopathies, Reiter's Disease, Stiff-Man Syndrome,

autoimmune pulmonary inflammation, autism, Guillain-Barre Syndrome, insulin dependent diabetes mellitus, and autoimmune inflammatory eye disorders.

[569] Additional disorders that are likely to have an autoimmune component that may be treated, prevented, diagnosed and/or prognosed with the compositions of the invention include, but are not limited to, scleroderma with anti-collagen antibodies (often characterized, e.g., by nucleolar and other nuclear antibodies), mixed connective tissue disease (often characterized, e.g., by antibodies to extractable nuclear antigens (e.g., ribonucleoprotein)), polymyositis (often characterized, e.g., by nonhistone ANA), pernicious anemia (often characterized, e.g., by antiparietal cell, microsomes, and intrinsic factor antibodies), idiopathic Addison's disease (often characterized, e.g., by humoral and cell-mediated adrenal cytotoxicity, infertility (often characterized, e.g., by antispermatozoal antibodies), glomerulonephritis (often characterized, e.g., by glomerular basement membrane antibodies or immune complexes), bullous pemphigoid (often characterized, e.g., by IgG and complement in basement membrane), Sjogren's syndrome (often characterized, e.g., by multiple tissue antibodies, and/or a specific nonhistone ANA (SS-B)), diabetes mellitus (often characterized, e.g., by cell-mediated and humoral islet cell antibodies), and adrenergic drug resistance (including adrenergic drug resistance with asthma or cystic fibrosis) (often characterized, e.g., by beta-adrenergic receptor antibodies).

[570] Additional disorders that may have an autoimmune component that may be treated, prevented, diagnosed and/or prognosed with the compositions of the invention include, but are not limited to, chronic active hepatitis (often characterized, e.g., by smooth muscle antibodies), primary biliary cirrhosis (often characterized, e.g., by mitochondria antibodies), other endocrine gland failure (often characterized, e.g., by specific tissue antibodies in some cases), vitiligo (often characterized, e.g., by melanocyte antibodies), vasculitis (often characterized, e.g., by Ig and complement in vessel walls and/or low serum complement), post-MI (often characterized, e.g., by myocardial antibodies), cardiomy syndrome (often characterized, e.g., by myocardial antibodies), urticaria (often characterized, e.g., by IgG and IgM antibodies to IgE), atopic dermatitis (often characterized, e.g., by IgG and IgM antibodies to IgE), asthma (often characterized, e.g., by IgG and IgM antibodies to IgE), and many other inflammatory, granulomatous, degenerative, and atrophic disorders.

[571] In a preferred embodiment, the autoimmune diseases and disorders and/or conditions associated with the diseases and disorders recited above are treated, prevented, diagnosed and/or prognosed using for example, antagonists or agonists, polypeptides or polynucleotides, or antibodies of the present invention. In a specific preferred embodiment, rheumatoid arthritis is treated, prevented, and/or diagnosed using polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention.

[572] In another specific preferred embodiment, systemic lupus erythematosus is treated, prevented, and/or diagnosed using polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention. In another specific preferred embodiment, idiopathic thrombocytopenia purpura is treated, prevented, and/or diagnosed using polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention.

[573] In another specific preferred embodiment IgA nephropathy is treated, prevented, and/or diagnosed using polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention.

[574] In a preferred embodiment, the autoimmune diseases and disorders and/or conditions associated with the diseases and disorders recited above are treated, prevented, diagnosed and/or prognosed using polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention

[575] In preferred embodiments, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are used as a immunosuppressive agent(s).

[576] Polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be useful in treating, preventing, prognosing, and/or diagnosing diseases, disorders, and/or conditions of hematopoietic cells. Polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention could be used to increase differentiation and proliferation of hematopoietic cells, including the pluripotent stem cells, in an effort to treat or prevent those diseases, disorders, and/or conditions associated with a decrease in certain (or many) types hematopoietic cells, including but not limited to, leukopenia, neutropenia, anemia, and thrombocytopenia. Alternatively, Polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention could be used to increase differentiation and proliferation of hematopoietic cells, including the

pluripotent stem cells, in an effort to treat or prevent those diseases, disorders, and/or conditions associated with an increase in certain (or many) types of hematopoietic cells, including but not limited to, histiocytosis.

[577] Allergic reactions and conditions, such as asthma (particularly allergic asthma) or other respiratory problems, may also be treated, prevented, diagnosed and/or prognosed using polypeptides, antibodies, or polynucleotides of the invention, and/or agonists or antagonists thereof. Moreover, these molecules can be used to treat, prevent, prognose, and/or diagnose anaphylaxis, hypersensitivity to an antigenic molecule, or blood group incompatibility.

[578] Additionally, polypeptides or polynucleotides of the invention, and/or agonists or antagonists thereof, may be used to treat, prevent, diagnose and/or prognose IgE-mediated allergic reactions. Such allergic reactions include, but are not limited to, asthma, rhinitis, and eczema. In specific embodiments, polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be used to modulate IgE concentrations in vitro or in vivo.

[579] Moreover, polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention have uses in the diagnosis, prognosis, prevention, and/or treatment of inflammatory conditions. For example, since polypeptides, antibodies, or polynucleotides of the invention, and/or agonists or antagonists of the invention may inhibit the activation, proliferation and/or differentiation of cells involved in an inflammatory response, these molecules can be used to prevent and/or treat chronic and acute inflammatory conditions. Such inflammatory conditions include, but are not limited to, for example, inflammation associated with infection (e.g., septic shock, sepsis, or systemic inflammatory response syndrome), ischemia-reperfusion injury, endotoxin lethality, complement-mediated hyperacute rejection, nephritis, cytokine or chemokine induced lung injury, inflammatory bowel disease, Crohn's disease, over production of cytokines (e.g., TNF or IL-1.), respiratory disorders (e.g., asthma and allergy); gastrointestinal disorders (e.g., inflammatory bowel disease); cancers (e.g., gastric, ovarian, lung, bladder, liver, and breast); CNS disorders (e.g., multiple sclerosis; ischemic brain injury and/or stroke, traumatic brain injury, neurodegenerative disorders (e.g., Parkinson's disease and Alzheimer's disease); AIDS-related dementia; and prion disease);

cardiovascular disorders (e.g., atherosclerosis, myocarditis, cardiovascular disease, and cardiopulmonary bypass complications); as well as many additional diseases, conditions, and disorders that are characterized by inflammation (e.g., hepatitis, rheumatoid arthritis, gout, trauma, pancreatitis, sarcoidosis, dermatitis, renal ischemia-reperfusion injury, Grave's disease, systemic lupus erythematosus, diabetes mellitus, and allogenic transplant rejection).

[580] Because inflammation is a fundamental defense mechanism, inflammatory disorders can effect virtually any tissue of the body. Accordingly, polynucleotides, polypeptides, and antibodies of the invention, as well as agonists or antagonists thereof, have uses in the treatment of tissue-specific inflammatory disorders, including, but not limited to, adrenalitis, alveolitis, angiocholecystitis, appendicitis, balanitis, blepharitis, bronchitis, bursitis, carditis, cellulitis, cervicitis, cholecystitis, chondritis, cochlitis, colitis, conjunctivitis, cystitis, dermatitis, diverticulitis, encephalitis, endocarditis, esophagitis, eustachitis, fibrositis, folliculitis, gastritis, gastroenteritis, gingivitis, glossitis, hepatosplenitis, keratitis, labyrinthitis, laryngitis, lymphangitis, mastitis, media otitis, meningitis, metritis, mucitis, myocarditis, myositis, myringitis, nephritis, neuritis, orchitis, osteochondritis, otitis, pericarditis, peritendonitis, peritonitis, pharyngitis, phlebitis, poliomyelitis, prostatitis, pulpitis, retinitis, rhinitis, salpingitis, scleritis, sclerochoroiditis, scrotitis, sinusitis, spondylitis, steatitis, stomatitis, synovitis, syringitis, tendonitis, tonsillitis, urethritis, and vaginitis.

[581] In specific embodiments, polypeptides, antibodies, or polynucleotides of the invention, and/or agonists or antagonists thereof, are useful to diagnose, prognose, prevent, and/or treat organ transplant rejections and graft-versus-host disease. Organ rejection occurs by host immune cell destruction of the transplanted tissue through an immune response. Similarly, an immune response is also involved in GVHD, but, in this case, the foreign transplanted immune cells destroy the host tissues. Polypeptides, antibodies, or polynucleotides of the invention, and/or agonists or antagonists thereof, that inhibit an immune response, particularly the activation, proliferation, differentiation, or chemotaxis of T-cells, may be an effective therapy in preventing organ rejection or GVHD. In specific embodiments, polypeptides, antibodies, or polynucleotides of the invention, and/or agonists or antagonists thereof, that inhibit an immune response, particularly the activation,

proliferation, differentiation, or chemotaxis of T-cells, may be an effective therapy in preventing experimental allergic and hyperacute xenograft rejection.

[582] In other embodiments, polypeptides, antibodies, or polynucleotides of the invention, and/or agonists or antagonists thereof, are useful to diagnose, prognose, prevent, and/or treat immune complex diseases, including, but not limited to, serum sickness, post streptococcal glomerulonephritis, polyarteritis nodosa, and immune complex-induced vasculitis.

[583] Polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the invention can be used to treat, detect, and/or prevent infectious agents. For example, by increasing the immune response, particularly increasing the proliferation activation and/or differentiation of B and/or T cells, infectious diseases may be treated, detected, and/or prevented. The immune response may be increased by either enhancing an existing immune response, or by initiating a new immune response. Alternatively, polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may also directly inhibit the infectious agent (refer to section of application listing infectious agents, etc), without necessarily eliciting an immune response.

[584] In another embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are used as a vaccine adjuvant that enhances immune responsiveness to an antigen. In a specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are used as an adjuvant to enhance tumor-specific immune responses.

[585] In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are used as an adjuvant to enhance anti-viral immune responses. Anti-viral immune responses that may be enhanced using the compositions of the invention as an adjuvant, include virus and virus associated diseases or symptoms described herein or otherwise known in the art. In specific embodiments, the compositions of the invention are used as an adjuvant to enhance an immune response to a virus, disease, or symptom selected from the group consisting of: AIDS, meningitis, Dengue, EBV, and hepatitis (e.g., hepatitis B). In another specific embodiment, the compositions of the invention are used as an adjuvant to enhance an immune response to a virus, disease, or symptom selected from the group consisting of: HIV/AIDS, respiratory

syncytial virus, Dengue, rotavirus, Japanese B encephalitis, influenza A and B, parainfluenza, measles, cytomegalovirus, rabies, Junin, Chikungunya, Rift Valley Fever, herpes simplex, and yellow fever.

[586] In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are used as an adjuvant to enhance anti-bacterial or anti-fungal immune responses. Anti-bacterial or anti-fungal immune responses that may be enhanced using the compositions of the invention as an adjuvant, include bacteria or fungus and bacteria or fungus associated diseases or symptoms described herein or otherwise known in the art. In specific embodiments, the compositions of the invention are used as an adjuvant to enhance an immune response to a bacteria or fungus, disease, or symptom selected from the group consisting of: tetanus, Diphtheria, botulism, and meningitis type B.

[587] In another specific embodiment, the compositions of the invention are used as an adjuvant to enhance an immune response to a bacteria or fungus, disease, or symptom selected from the group consisting of: *Vibrio cholerae*, *Mycobacterium leprae*, *Salmonella typhi*, *Salmonella paratyphi*, *Meisseria meningitidis*, *Streptococcus pneumoniae*, Group B streptococcus, *Shigella spp.*, Enterotoxigenic *Escherichia coli*, Enterohemorrhagic *E. coli*, and *Borrelia burgdorferi*.

[588] In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are used as an adjuvant to enhance anti-parasitic immune responses. Anti-parasitic immune responses that may be enhanced using the compositions of the invention as an adjuvant, include parasite and parasite associated diseases or symptoms described herein or otherwise known in the art. In specific embodiments, the compositions of the invention are used as an adjuvant to enhance an immune response to a parasite. In another specific embodiment, the compositions of the invention are used as an adjuvant to enhance an immune response to Plasmodium (malaria) or Leishmania.

[589] In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention may also be employed to treat infectious diseases including silicosis, sarcoidosis, and idiopathic pulmonary fibrosis; for example, by preventing the recruitment and activation of mononuclear phagocytes.

vivo. Moreover, in related embodiments, said enhancement or antagonism of antigen presentation may be useful as an anti-tumor treatment or to modulate the immune system.

[603] In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are used as an agent to direct an individual's immune system towards development of a humoral response (i.e. TH2) as opposed to a TH1 cellular response.

[604] In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are used as a means to induce tumor proliferation and thus make it more susceptible to anti-neoplastic agents. For example, multiple myeloma is a slowly dividing disease and is thus refractory to virtually all anti-neoplastic regimens. If these cells were forced to proliferate more rapidly their susceptibility profile would likely change.

[605] In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are used as a stimulator of B cell production in pathologies such as AIDS, chronic lymphocyte disorder and/or Common Variable Immunodeficiency.

[606] In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are used as a therapy for generation and/or regeneration of lymphoid tissues following surgery, trauma or genetic defect. In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are used in the pretreatment of bone marrow samples prior to transplant.

[607] In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are used as a gene-based therapy for genetically inherited disorders resulting in immuno-incompetence/immunodeficiency such as observed among SCID patients.

[608] In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are used as a means of activating monocytes/macrophages to defend against parasitic diseases that effect monocytes such as Leishmania.

[609] In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are used as a means of regulating secreted cytokines that are elicited by polypeptides of the invention.

[610] In another embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are used in one or more of the applications described herein, as they may apply to veterinary medicine.

[611] In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are used as a means of blocking various aspects of immune responses to foreign agents or self. Examples of diseases or conditions in which blocking of certain aspects of immune responses may be desired include autoimmune disorders such as lupus, and arthritis, as well as immunoresponsiveness to skin allergies, inflammation, bowel disease, injury and diseases/disorders associated with pathogens.

[612] In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are used as a therapy for preventing the B cell proliferation and Ig secretion associated with autoimmune diseases such as idiopathic thrombocytopenic purpura, systemic lupus erythematosus and multiple sclerosis.

[613] In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are used as an inhibitor of B and/or T cell migration in endothelial cells. This activity disrupts tissue architecture or cognate responses and is useful, for example in disrupting immune responses, and blocking sepsis.

[614] In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are used as a therapy for chronic hypergammaglobulinemia evident in such diseases as monoclonal gammopathy of undetermined significance (MGUS), Waldenstrom's disease, related idiopathic monoclonal gammopathies, and plasmacytomas.

[615] In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention may be employed for instance to inhibit polypeptide chemotaxis and activation of macrophages and their precursors, and of neutrophils, basophils, B lymphocytes and some T-cell subsets, e.g., activated and CD8 cytotoxic T cells and natural killer cells, in certain autoimmune and chronic inflammatory

and infective diseases. Examples of autoimmune diseases are described herein and include multiple sclerosis, and insulin-dependent diabetes.

[616] The polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention may also be employed to treat idiopathic hyper-eosinophilic syndrome by, for example, preventing eosinophil production and migration.

[617] In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are used to enhance or inhibit complement mediated cell lysis.

[618] In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are used to enhance or inhibit antibody dependent cellular cytotoxicity.

[619] In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention may also be employed for treating atherosclerosis, for example, by preventing monocyte infiltration in the artery wall.

[620] In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention may be employed to treat adult respiratory distress syndrome (ARDS).

[621] In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention may be useful for stimulating wound and tissue repair, stimulating angiogenesis, and/or stimulating the repair of vascular or lymphatic diseases or disorders. Additionally, agonists and antagonists of the invention may be used to stimulate the regeneration of mucosal surfaces.

[622] In a specific embodiment, polynucleotides or polypeptides, and/or agonists thereof are used to diagnose, prognose, treat, and/or prevent a disorder characterized by primary or acquired immunodeficiency, deficient serum immunoglobulin production, recurrent infections, and/or immune system dysfunction. Moreover, polynucleotides or polypeptides, and/or agonists thereof may be used to treat or prevent infections of the joints, bones, skin, and/or parotid glands, blood-borne infections (e.g., sepsis, meningitis, septic arthritis, and/or osteomyelitis), autoimmune diseases (e.g., those disclosed herein), inflammatory disorders, and malignancies, and/or any disease or disorder or condition associated with these infections, diseases, disorders and/or malignancies) including, but not

limited to, CVID, other primary immune deficiencies, HIV disease, CLL, recurrent bronchitis, sinusitis, otitis media, conjunctivitis, pneumonia, hepatitis, meningitis, herpes zoster (e.g., severe herpes zoster), and/or pneumocystis carinii. Other diseases and disorders that may be prevented, diagnosed, prognosed, and/or treated with polynucleotides or polypeptides, and/or agonists of the present invention include, but are not limited to, HIV infection, HTLV-BLV infection, lymphopenia, phagocyte bactericidal dysfunction anemia, thrombocytopenia, and hemoglobinuria.

[623] In another embodiment, polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention are used to treat, and/or diagnose an individual having common variable immunodeficiency disease ("CVID"; also known as "acquired agammaglobulinemia" and "acquired hypogammaglobulinemia") or a subset of this disease.

[624] In a specific embodiment, polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be used to diagnose, prognose, prevent, and/or treat cancers or neoplasms including immune cell or immune tissue-related cancers or neoplasms. Examples of cancers or neoplasms that may be prevented, diagnosed, or treated by polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention include, but are not limited to, acute myelogenous leukemia, chronic myelogenous leukemia, Hodgkin's disease, non-Hodgkin's lymphoma, acute lymphocytic anemia (ALL) Chronic lymphocyte leukemia, plasmacytomas, multiple myeloma, Burkitt's lymphoma, EBV-transformed diseases, and/or diseases and disorders described in the section entitled "Hyperproliferative Disorders" elsewhere herein.

[625] In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are used as a therapy for decreasing cellular proliferation of Large B-cell Lymphomas.

[626] In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are used as a means of decreasing the involvement of B cells and Ig associated with Chronic Myelogenous Leukemia.

[627] In specific embodiments, the compositions of the invention are used as an agent to boost immunoresponsiveness among B cell immunodeficient individuals, such as, for example, an individual who has undergone a partial or complete splenectomy.

[628] Antagonists of the invention include, for example, binding and/or inhibitory antibodies, antisense nucleic acids, ribozymes or soluble forms of the polypeptides of the present invention (e.g., Fc fusion protein; see, e.g., Example 9). Agonists of the invention include, for example, binding or stimulatory antibodies, and soluble forms of the polypeptides (e.g., Fc fusion proteins; see, e.g., Example 9). polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention may be employed in a composition with a pharmaceutically acceptable carrier, e.g., as described herein.

[629] In another embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are administered to an animal (including, but not limited to, those listed above, and also including transgenic animals) incapable of producing functional endogenous antibody molecules or having an otherwise compromised endogenous immune system, but which is capable of producing human immunoglobulin molecules by means of a reconstituted or partially reconstituted immune system from another animal (see, e.g., published PCT Application Nos. WO98/24893, WO/9634096, WO/9633735, and WO/9110741). Administration of polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention to such animals is useful for the generation of monoclonal antibodies against the polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention.

Blood-Related Disorders

[630] The polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be used to modulate hemostatic (the stopping of bleeding) or thrombolytic (clot dissolving) activity. For example, by increasing hemostatic or thrombolytic activity, polynucleotides or polypeptides, and/or agonists or antagonists of the present invention could be used to treat or prevent blood coagulation diseases, disorders, and/or conditions (e.g., afibrinogenemia, factor deficiencies, hemophilia), blood platelet diseases, disorders, and/or conditions (e.g., thrombocytopenia), or wounds resulting from trauma, surgery, or other causes. Alternatively, polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention that can decrease hemostatic or thrombolytic activity could be used to inhibit or dissolve clotting. These molecules could be important in the treatment or prevention of heart attacks (infarction), strokes, or scarring.

[631] In specific embodiments, the polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be used to prevent, diagnose, prognose, and/or treat thrombosis, arterial thrombosis, venous thrombosis, thromboembolism, pulmonary embolism, atherosclerosis, myocardial infarction, transient ischemic attack, unstable angina. In specific embodiments, the polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be used for the prevention of occlusion of saphenous grafts, for reducing the risk of periprocedural thrombosis as might accompany angioplasty procedures, for reducing the risk of stroke in patients with atrial fibrillation including nonrheumatic atrial fibrillation, for reducing the risk of embolism associated with mechanical heart valves and or mitral valves disease. Other uses for the polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention, include, but are not limited to, the prevention of occlusions in extracorporeal devices (e.g., intravascular canulas, vascular access shunts in hemodialysis patients, hemodialysis machines, and cardiopulmonary bypass machines).

[632] In another embodiment, a polypeptide of the invention, or polynucleotides, antibodies, agonists, or antagonists corresponding to that polypeptide, may be used to prevent, diagnose, prognose, and/or treat diseases and disorders of the blood and/or blood forming organs associated with the tissue(s) in which the polypeptide of the invention is expressed, including one, two, three, four, five, or more tissues disclosed in Table 1B, column 8 (Tissue Distribution Library Code).

[633] The polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be used to modulate hematopoietic activity (the formation of blood cells). For example, the polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be used to increase the quantity of all or subsets of blood cells, such as, for example, erythrocytes, lymphocytes (B or T cells), myeloid cells (e.g., basophils, eosinophils, neutrophils, mast cells, macrophages) and platelets. The ability to decrease the quantity of blood cells or subsets of blood cells may be useful in the prevention, detection, diagnosis and/or treatment of anemias and leukopenias described below. Alternatively, the polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be used to decrease the quantity of all or subsets of blood cells, such as, for example, erythrocytes, lymphocytes (B or T cells), myeloid cells (e.g., basophils, eosinophils, neutrophils, mast cells, macrophages) and platelets.. The ability to decrease the quantity of blood cells or subsets of blood cells may be useful in the

prevention, detection, diagnosis and/or treatment of leukocytoses, such as, for example eosinophilia.

[634] The polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be used to prevent, treat, or diagnose blood dyscrasia.

[635] Anemias are conditions in which the number of red blood cells or amount of hemoglobin (the protein that carries oxygen) in them is below normal. Anemia may be caused by excessive bleeding, decreased red blood cell production, or increased red blood cell destruction (hemolysis). The polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be useful in treating, preventing, and/or diagnosing anemias. Anemias that may be treated prevented or diagnosed by the polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention include iron deficiency anemia, hypochromic anemia, microcytic anemia, chlorosis, hereditary sideroblastic anemia, idiopathic acquired sideroblastic anemia, red cell aplasia, megaloblastic anemia (e.g., pernicious anemia, (vitamin B12 deficiency) and folic acid deficiency anemia), aplastic anemia, hemolytic anemias (e.g., autoimmune hemolytic anemia, microangiopathic hemolytic anemia, and paroxysmal nocturnal hemoglobinuria). The polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be useful in treating, preventing, and/or diagnosing anemias associated with diseases including but not limited to, anemias associated with systemic lupus erythematosus, cancers, lymphomas, chronic renal disease, and enlarged spleens. The polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be useful in treating, preventing, and/or diagnosing anemias arising from drug treatments such as anemias associated with methyldopa, dapsone, and/or sulfadiazine. Additionally, the polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be useful in treating, preventing, and/or diagnosing anemias associated with abnormal red blood cell architecture including but not limited to, hereditary spherocytosis, hereditary elliptocytosis, glucose-6-phosphate dehydrogenase deficiency, and sickle cell anemia.

[636] The polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be useful in treating, preventing, and/or diagnosing hemoglobin abnormalities, (e.g., those associated with sickle cell anemia, hemoglobin C disease, hemoglobin S-C disease, and hemoglobin E disease). Additionally, the polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be

0950083-091201

useful in diagnosing, prognosing, preventing, and/or treating thalassemias, including, but not limited to major and minor forms of alpha-thalassemia and beta-thalassemia.

[637] In another embodiment, the polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be useful in diagnosing, prognosing, preventing, and/or treating bleeding disorders including, but not limited to, thrombocytopenia (e.g., idiopathic thrombocytopenic purpura, and thrombotic thrombocytopenic purpura), Von Willebrand's disease, hereditary platelet disorders (e.g., storage pool disease such as Chediak-Higashi and Hermansky-Pudlak syndromes, thromboxane A2 dysfunction, thromboasthenia, and Bernard-Soulier syndrome), hemolytic-uremic syndrome, hemophelias such as hemophilia A or Factor VII deficiency and Christmas disease or Factor IX deficiency, Hereditary Hemorrhagic Telangiectasia, also known as Rendu-Osler-Weber syndrome, allergic purpura (Henoch Schonlein purpura) and disseminated intravascular coagulation.

[638] The effect of the polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention on the clotting time of blood may be monitored using any of the clotting tests known in the art including, but not limited to, whole blood partial thromboplastin time (PTT), the activated partial thromboplastin time (aPTT), the activated clotting time (ACT), the recalcified activated clotting time, or the Lee-White Clotting time.

[639] Several diseases and a variety of drugs can cause platelet dysfunction. Thus, in a specific embodiment, the polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be useful in diagnosing, prognosing, preventing, and/or treating acquired platelet dysfunction such as platelet dysfunction accompanying kidney failure, leukemia, multiple myeloma, cirrhosis of the liver, and systemic lupus erythematosus as well as platelet dysfunction associated with drug treatments, including treatment with aspirin, ticlopidine, nonsteroidal anti-inflammatory drugs (used for arthritis, pain, and sprains), and penicillin in high doses.

[640] In another embodiment, the polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be useful in diagnosing, prognosing, preventing, and/or treating diseases and disorders characterized by or associated with increased or decreased numbers of white blood cells. Leukopenia occurs when the number of white blood cells decreases below normal. Leukopenias include, but are not limited to, neutropenia and lymphocytopenia. An increase in the number of white blood cells compared to normal is known as leukocytosis. The body generates increased numbers of

white blood cells during infection. Thus, leukocytosis may simply be a normal physiological parameter that reflects infection. Alternatively, leukocytosis may be an indicator of injury or other disease such as cancer. Leukocytoses, include but are not limited to, eosinophilia, and accumulations of macrophages. In specific embodiments, the polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be useful in diagnosing, prognosing, preventing, and/or treating leukopenia. In other specific embodiments, the polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be useful in diagnosing, prognosing, preventing, and/or treating leukocytosis.

[641] Leukopenia may be a generalized decrease in all types of white blood cells, or may be a specific depletion of particular types of white blood cells. Thus, in specific embodiments, the polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be useful in diagnosing, prognosing, preventing, and/or treating decreases in neutrophil numbers, known as neutropenia. Neutropenias that may be diagnosed, prognosed, prevented, and/or treated by the polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention include, but are not limited to, infantile genetic agranulocytosis, familial neutropenia, cyclic neutropenia, neutropenias resulting from or associated with dietary deficiencies (e.g., vitamin B 12 deficiency or folic acid deficiency), neutropenias resulting from or associated with drug treatments (e.g., antibiotic regimens such as penicillin treatment, sulfonamide treatment, anticoagulant treatment, anticonvulsant drugs, anti-thyroid drugs, and cancer chemotherapy), and neutropenias resulting from increased neutrophil destruction that may occur in association with some bacterial or viral infections, allergic disorders, autoimmune diseases, conditions in which an individual has an enlarged spleen (e.g., Felty syndrome, malaria and sarcoidosis), and some drug treatment regimens.

[642] The polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be useful in diagnosing, prognosing, preventing, and/or treating lymphocytopenias (decreased numbers of B and/or T lymphocytes), including, but not limited to lymphocytopenias resulting from or associated with stress, drug treatments (e.g., drug treatment with corticosteroids, cancer chemotherapies, and/or radiation therapies), AIDS infection and/or other diseases such as, for example, cancer, rheumatoid arthritis, systemic lupus erythematosus, chronic infections, some viral infections and/or hereditary

[648] In other embodiments, the polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be useful as a treatment prior to surgery, to increase blood cell production.

[649] In other embodiments, the polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be useful as an agent to enhance the migration, phagocytosis, superoxide production, antibody dependent cellular cytotoxicity of neutrophils, eosinophils and macrophages.

[650] In other embodiments, the polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be useful as an agent to increase the number of stem cells in circulation prior to stem cells pheresis. In another specific embodiment, the polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be useful as an agent to increase the number of stem cells in circulation prior to platelet pheresis.

[651] In other embodiments, the polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be useful as an agent to increase cytokine production.

[652] In other embodiments, the polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be useful in preventing, diagnosing, and/or treating primary hematopoietic disorders.

Hyperproliferative Disorders

[653] In certain embodiments, polynucleotides or polypeptides, or agonists or antagonists of the present invention can be used to treat or detect hyperproliferative disorders, including neoplasms. Polynucleotides or polypeptides, or agonists or antagonists of the present invention may inhibit the proliferation of the disorder through direct or indirect interactions. Alternatively, Polynucleotides or polypeptides, or agonists or antagonists of the present invention may proliferate other cells which can inhibit the hyperproliferative disorder.

[654] For example, by increasing an immune response, particularly increasing antigenic qualities of the hyperproliferative disorder or by proliferating, differentiating, or mobilizing T-cells, hyperproliferative disorders can be treated. This immune response may be increased by either enhancing an existing immune response, or by initiating a new immune

response. Alternatively, decreasing an immune response may also be a method of treating hyperproliferative disorders, such as a chemotherapeutic agent.

[655] Examples of hyperproliferative disorders that can be treated or detected by polynucleotides or polypeptides, or agonists or antagonists of the present invention include, but are not limited to neoplasms located in the: colon, abdomen, bone, breast, digestive system, liver, pancreas, peritoneum, endocrine glands (adrenal, parathyroid, pituitary, testicles, ovary, thymus, thyroid), eye, head and neck, nervous (central and peripheral), lymphatic system, pelvis, skin, soft tissue, spleen, thorax, and urogenital tract.

[656] Similarly, other hyperproliferative disorders can also be treated or detected by polynucleotides or polypeptides, or agonists or antagonists of the present invention. Examples of such hyperproliferative disorders include, but are not limited to: Acute Childhood Lymphoblastic Leukemia, Acute Lymphoblastic Leukemia, Acute Lymphocytic Leukemia, Acute Myeloid Leukemia, Adrenocortical Carcinoma, Adult (Primary) Hepatocellular Cancer, Adult (Primary) Liver Cancer, Adult Acute Lymphocytic Leukemia, Adult Acute Myeloid Leukemia, Adult Hodgkin's Disease, Adult Hodgkin's Lymphoma, Adult Lymphocytic Leukemia, Adult Non-Hodgkin's Lymphoma, Adult Primary Liver Cancer, Adult Soft Tissue Sarcoma, AIDS-Related Lymphoma, AIDS-Related Malignancies, Anal Cancer, Astrocytoma, Bile Duct Cancer, Bladder Cancer, Bone Cancer, Brain Stem Glioma, Brain Tumors, Breast Cancer, Cancer of the Renal Pelvis and Ureter, Central Nervous System (Primary) Lymphoma, Central Nervous System Lymphoma, Cerebellar Astrocytoma, Cerebral Astrocytoma, Cervical Cancer, Childhood (Primary) Hepatocellular Cancer, Childhood (Primary) Liver Cancer, Childhood Acute Lymphoblastic Leukemia, Childhood Acute Myeloid Leukemia, Childhood Brain Stem Glioma, Childhood Cerebellar Astrocytoma, Childhood Cerebral Astrocytoma, Childhood Extracranial Germ Cell Tumors, Childhood Hodgkin's Disease, Childhood Hodgkin's Lymphoma, Childhood Hypothalamic and Visual Pathway Glioma, Childhood Lymphoblastic Leukemia, Childhood Medulloblastoma, Childhood Non-Hodgkin's Lymphoma, Childhood Pineal and Supratentorial Primitive Neuroectodermal Tumors, Childhood Primary Liver Cancer, Childhood Rhabdomyosarcoma, Childhood Soft Tissue Sarcoma, Childhood Visual Pathway and Hypothalamic Glioma, Chronic Lymphocytic Leukemia, Chronic Myelogenous Leukemia, Colon Cancer, Cutaneous T-Cell Lymphoma, Endocrine Pancreas Islet Cell Carcinoma, Endometrial Cancer, Ependymoma, Epithelial Cancer, Esophageal Cancer, Ewing's Sarcoma and Related Tumors, Exocrine Pancreatic Cancer, Extracranial

[657] In another preferred embodiment, polynucleotides or polypeptides, or agonists or antagonists of the present invention are used to diagnose, prognose, prevent, and/or treat premalignant conditions and to prevent progression to a neoplastic or malignant state, including but not limited to those disorders described above. Such uses are indicated in conditions known or suspected of preceding progression to neoplasia or cancer, in particular, where non-neoplastic cell growth consisting of hyperplasia, metaplasia, or most particularly, dysplasia has occurred (for review of such abnormal growth conditions, see Robbins and Angell, 1976, Basic Pathology, 2d Ed., W. B. Saunders Co., Philadelphia, pp. 68-79.)

[658] Hyperplasia is a form of controlled cell proliferation, involving an increase in cell number in a tissue or organ, without significant alteration in structure or function. Hyperplastic disorders which can be diagnosed, prognosed, prevented, and/or treated with compositions of the invention (including polynucleotides, polypeptides, agonists or antagonists) include, but are not limited to, angiofollicular mediastinal lymph node hyperplasia, angiolymphoid hyperplasia with eosinophilia, atypical melanocytic hyperplasia, basal cell hyperplasia, benign giant lymph node hyperplasia, cementum hyperplasia, congenital adrenal hyperplasia, congenital sebaceous hyperplasia, cystic hyperplasia, cystic hyperplasia of the breast, denture hyperplasia, ductal hyperplasia, endometrial hyperplasia, fibromuscular hyperplasia, focal epithelial hyperplasia, gingival hyperplasia, inflammatory fibrous hyperplasia, inflammatory papillary hyperplasia, intravascular papillary endothelial hyperplasia, nodular hyperplasia of prostate, nodular regenerative hyperplasia, pseudoepitheliomatous hyperplasia, senile sebaceous hyperplasia, and verrucous hyperplasia.

[659] Metaplasia is a form of controlled cell growth in which one type of adult or fully differentiated cell substitutes for another type of adult cell. Metaplastic disorders which can be diagnosed, prognosed, prevented, and/or treated with compositions of the invention (including polynucleotides, polypeptides, agonists or antagonists) include, but are not limited to, agnogenic myeloid metaplasia, apocrine metaplasia, atypical metaplasia, autoparenchymatous metaplasia, connective tissue metaplasia, epithelial metaplasia, intestinal metaplasia, metaplastic anemia, metaplastic ossification, metaplastic polyps, myeloid metaplasia, primary myeloid metaplasia, secondary myeloid metaplasia, squamous metaplasia, squamous metaplasia of amnion, and symptomatic myeloid metaplasia.

[660] Dysplasia is frequently a forerunner of cancer, and is found mainly in the epithelia; it is the most disorderly form of non-neoplastic cell growth, involving a loss in individual cell uniformity and in the architectural orientation of cells. Dysplastic cells often have abnormally large, deeply stained nuclei, and exhibit pleomorphism. Dysplasia characteristically occurs where there exists chronic irritation or inflammation. Dysplastic disorders which can be diagnosed, prognosed, prevented, and/or treated with compositions of the invention (including polynucleotides, polypeptides, agonists or antagonists) include, but are not limited to, anhidrotic ectodermal dysplasia, anterofacial dysplasia, asphyxiating thoracic dysplasia, atriadigital dysplasia, bronchopulmonary dysplasia, cerebral dysplasia, cervical dysplasia, chondroectodermal dysplasia, cleidocranial dysplasia, congenital ectodermal dysplasia, craniodiaphysial dysplasia, craniocarpotarsal dysplasia, craniometaphysial dysplasia, dentin dysplasia, diaphysial dysplasia, ectodermal dysplasia, enamel dysplasia, encephalo-ophthalmic dysplasia, dysplasia epiphysialis hemimelia, dysplasia epiphysialis multiplex, dysplasia epiphysialis punctata, epithelial dysplasia, faciogradigital dysplasia, familial fibrous dysplasia of jaws, familial white folded dysplasia, fibromuscular dysplasia, fibrous dysplasia of bone, florid osseous dysplasia, hereditary renal-retinal dysplasia, hidrotic ectodermal dysplasia, hypohidrotic ectodermal dysplasia, lymphopenic thymic dysplasia, mammary dysplasia, mandibulofacial dysplasia, metaphysial dysplasia, Mondini dysplasia, monostotic fibrous dysplasia, mucoepithelial dysplasia, multiple epiphysial dysplasia, oculoauriculovertbral dysplasia, oculodentodigital dysplasia, oculovertbral dysplasia, odontogenic dysplasia, ophthalmomandibulomelic dysplasia, periapical cemental dysplasia, polyostotic fibrous dysplasia, pseudoachondroplastic spondyloepiphysial dysplasia, retinal dysplasia, septo-optic dysplasia, spondyloepiphysial dysplasia, and ventriculoradial dysplasia.

[661] Additional pre-neoplastic disorders which can be diagnosed, prognosed, prevented, and/or treated with compositions of the invention (including polynucleotides, polypeptides, agonists or antagonists) include, but are not limited to, benign dysproliferative disorders (e.g., benign tumors, fibrocystic conditions, tissue hypertrophy, intestinal polyps, colon polyps, and esophageal dysplasia), leukoplakia, keratoses, Bowen's disease, Farmer's Skin, solar cheilitis, and solar keratosis.

[662] In another embodiment, a polypeptide of the invention, or polynucleotides, antibodies, agonists, or antagonists corresponding to that polypeptide, may be used to diagnose and/or prognose disorders associated with the tissue(s) in which the polypeptide of

the invention is expressed, including one, two, three, four, five, or more tissues disclosed in Table 1B, column 8 (Tissue Distribution Library Code).

[663] In another embodiment, polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention conjugated to a toxin or a radioactive isotope, as described herein, may be used to treat cancers and neoplasms, including, but not limited to those described herein. In a further preferred embodiment, polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention conjugated to a toxin or a radioactive isotope, as described herein, may be used to treat acute myelogenous leukemia.

[664] Additionally, polynucleotides, polypeptides, and/or agonists or antagonists of the invention may affect apoptosis, and therefore, would be useful in treating a number of diseases associated with increased cell survival or the inhibition of apoptosis. For example, diseases associated with increased cell survival or the inhibition of apoptosis that could be diagnosed, prognosed, prevented, and/or treated by polynucleotides, polypeptides, and/or agonists or antagonists of the invention, include cancers (such as follicular lymphomas, carcinomas with p53 mutations, and hormone-dependent tumors, including, but not limited to colon cancer, cardiac tumors, pancreatic cancer, melanoma, retinoblastoma, glioblastoma, lung cancer, intestinal cancer, testicular cancer, stomach cancer, neuroblastoma, myxoma, myoma, lymphoma, endothelioma, osteoblastoma, osteoclastoma, osteosarcoma, chondrosarcoma, adenoma, breast cancer, prostate cancer, Kaposi's sarcoma and ovarian cancer); autoimmune disorders such as, multiple sclerosis, Sjogren's syndrome, Hashimoto's thyroiditis, biliary cirrhosis, Behcet's disease, Crohn's disease, polymyositis, systemic lupus erythematosus and immune-related glomerulonephritis and rheumatoid arthritis) and viral infections (such as herpes viruses, pox viruses and adenoviruses), inflammation, graft v. host disease, acute graft rejection, and chronic graft rejection.

[665] In preferred embodiments, polynucleotides, polypeptides, and/or agonists or antagonists of the invention are used to inhibit growth, progression, and/or metastasis of cancers, in particular those listed above.

[666] Additional diseases or conditions associated with increased cell survival that could be diagnosed, prognosed, prevented, and/or treated by polynucleotides, polypeptides, and/or agonists or antagonists of the invention, include, but are not limited to, progression, and/or metastases of malignancies and related disorders such as leukemia (including acute leukemias (e.g., acute lymphocytic leukemia, acute myelocytic leukemia (including

myeloblastic, promyelocytic, myelomonocytic, monocytic, and erythroleukemia)) and chronic leukemias (e.g., chronic myelocytic (granulocytic) leukemia and chronic lymphocytic leukemia)), polycythemia vera, lymphomas (e.g., Hodgkin's disease and non-Hodgkin's disease), multiple myeloma, Waldenstrom's macroglobulinemia, heavy chain disease, and solid tumors including, but not limited to, sarcomas and carcinomas such as fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, osteogenic sarcoma, chordoma, angiosarcoma, endotheliosarcoma, lymphangiosarcoma, lymphangioendotheliosarcoma, synovioma, mesothelioma, Ewing's tumor, leiomyosarcoma, rhabdomyosarcoma, colon carcinoma, pancreatic cancer, breast cancer, ovarian cancer, prostate cancer, squamous cell carcinoma, basal cell carcinoma, adenocarcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma, papillary adenocarcinomas, cystadenocarcinoma, medullary carcinoma, bronchogenic carcinoma, renal cell carcinoma, hepatoma, bile duct carcinoma, choriocarcinoma, seminoma, embryonal carcinoma, Wilm's tumor, cervical cancer, testicular tumor, lung carcinoma, small cell lung carcinoma, bladder carcinoma, epithelial carcinoma, glioma, astrocytoma, medulloblastoma, craniopharyngioma, ependymoma, pinealoma, emangioblastoma, acoustic neuroma, oligodendroglioma, menangioma, melanoma, neuroblastoma, and retinoblastoma.

[667] Diseases associated with increased apoptosis that could be diagnosed, prognosed, prevented, and/or treated by polynucleotides, polypeptides, and/or agonists or antagonists of the invention, include AIDS; neurodegenerative disorders (such as Alzheimer's disease, Parkinson's disease, amyotrophic lateral sclerosis, retinitis pigmentosa, cerebellar degeneration and brain tumor or prior associated disease); autoimmune disorders (such as, multiple sclerosis, Sjogren's syndrome, Hashimoto's thyroiditis, biliary cirrhosis, Behcet's disease, Crohn's disease, polymyositis, systemic lupus erythematosus and immune-related glomerulonephritis and rheumatoid arthritis) myelodysplastic syndromes (such as aplastic anemia), graft v. host disease, ischemic injury (such as that caused by myocardial infarction, stroke and reperfusion injury), liver injury (e.g., hepatitis related liver injury, ischemia/reperfusion injury, cholestasis (bile duct injury) and liver cancer); toxin-induced liver disease (such as that caused by alcohol), septic shock, cachexia and anorexia.

[668] Hyperproliferative diseases and/or disorders that could be diagnosed, prognosed, prevented, and/or treated by polynucleotides, polypeptides, and/or agonists or antagonists of the invention, include, but are not limited to, neoplasms located in the liver, abdomen, bone,

breast, digestive system, pancreas, peritoneum, endocrine glands (adrenal, parathyroid, pituitary, testicles, ovary, thymus, thyroid), eye, head and neck, nervous system (central and peripheral), lymphatic system, pelvis, skin, soft tissue, spleen, thorax, and urogenital tract.

[669] Similarly, other hyperproliferative disorders can also be diagnosed, prognosed, prevented, and/or treated by polynucleotides, polypeptides, and/or agonists or antagonists of the invention. Examples of such hyperproliferative disorders include, but are not limited to: hypergammaglobulinemia, lymphoproliferative disorders, paraproteinemias, purpura, sarcoidosis, Sezary Syndrome, Waldenstrom's macroglobulinemia, Gaucher's Disease, histiocytosis, and any other hyperproliferative disease, besides neoplasia, located in an organ system listed above.

[670] Another preferred embodiment utilizes polynucleotides of the present invention to inhibit aberrant cellular division, by gene therapy using the present invention, and/or protein fusions or fragments thereof.

[671] Thus, the present invention provides a method for treating cell proliferative disorders by inserting into an abnormally proliferating cell a polynucleotide of the present invention, wherein said polynucleotide represses said expression.

[672] Another embodiment of the present invention provides a method of treating cell-proliferative disorders in individuals comprising administration of one or more active gene copies of the present invention to an abnormally proliferating cell or cells. In a preferred embodiment, polynucleotides of the present invention is a DNA construct comprising a recombinant expression vector effective in expressing a DNA sequence encoding said polynucleotides. In another preferred embodiment of the present invention, the DNA construct encoding the polynucleotides of the present invention is inserted into cells to be treated utilizing a retrovirus, or more preferably an adenoviral vector (See G J. Nabel, et. al., PNAS 1999 96: 324-326, which is hereby incorporated by reference). In a most preferred embodiment, the viral vector is defective and will not transform non-proliferating cells, only proliferating cells. Moreover, in a preferred embodiment, the polynucleotides of the present invention inserted into proliferating cells either alone, or in combination with or fused to other polynucleotides, can then be modulated via an external stimulus (i.e. magnetic, specific small molecule, chemical, or drug administration, etc.), which acts upon the promoter upstream of said polynucleotides to induce expression of the encoded protein product. As such the beneficial therapeutic affect of the present invention may be expressly

modulated (i.e. to increase, decrease, or inhibit expression of the present invention) based upon said external stimulus.

[673] Polynucleotides of the present invention may be useful in repressing expression of oncogenic genes or antigens. By "repressing expression of the oncogenic genes " is intended the suppression of the transcription of the gene, the degradation of the gene transcript (pre-message RNA), the inhibition of splicing, the destruction of the messenger RNA, the prevention of the post-translational modifications of the protein, the destruction of the protein, or the inhibition of the normal function of the protein.

[674] For local administration to abnormally proliferating cells, polynucleotides of the present invention may be administered by any method known to those of skill in the art including, but not limited to transfection, electroporation, microinjection of cells, or in vehicles such as liposomes, lipofectin, or as naked polynucleotides, or any other method described throughout the specification. The polynucleotide of the present invention may be delivered by known gene delivery systems such as, but not limited to, retroviral vectors (Gilboa, J. Virology 44:845 (1982); Hocke, Nature 320:275 (1986); Wilson, et al., Proc. Natl. Acad. Sci. U.S.A. 85:3014), vaccinia virus system (Chakrabarty et al., Mol. Cell Biol. 5:3403 (1985) or other efficient DNA delivery systems (Yates et al., Nature 313:812 (1985)) known to those skilled in the art. These references are exemplary only and are hereby incorporated by reference. In order to specifically deliver or transfect cells which are abnormally proliferating and spare non-dividing cells, it is preferable to utilize a retrovirus, or adenoviral (as described in the art and elsewhere herein) delivery system known to those of skill in the art. Since host DNA replication is required for retroviral DNA to integrate and the retrovirus will be unable to self replicate due to the lack of the retrovirus genes needed for its life cycle. Utilizing such a retroviral delivery system for polynucleotides of the present invention will target said gene and constructs to abnormally proliferating cells and will spare the non-dividing normal cells.

[675] The polynucleotides of the present invention may be delivered directly to cell proliferative disorder/disease sites in internal organs, body cavities and the like by use of imaging devices used to guide an injecting needle directly to the disease site. The polynucleotides of the present invention may also be administered to disease sites at the time of surgical intervention.

[676] By "cell proliferative disease" is meant any human or animal disease or disorder, affecting any one or any combination of organs, cavities, or body parts, which is

[682] It is preferred to use high affinity and/or potent *in vivo* inhibiting and/or neutralizing antibodies against polypeptides or polynucleotides of the present invention, fragments or regions thereof, for both immunoassays directed to and therapy of disorders related to polynucleotides or polypeptides, including fragments thereof, of the present invention. Such antibodies, fragments, or regions, will preferably have an affinity for polynucleotides or polypeptides, including fragments thereof. Preferred binding affinities include those with a dissociation constant or K_d less than $5 \times 10^{-6}M$, $10^{-6}M$, $5 \times 10^{-7}M$, $10^{-7}M$, $5 \times 10^{-8}M$, $10^{-8}M$, $5 \times 10^{-9}M$, $10^{-9}M$, $5 \times 10^{-10}M$, $10^{-10}M$, $5 \times 10^{-11}M$, $10^{-11}M$, $5 \times 10^{-12}M$, $10^{-12}M$, $5 \times 10^{-13}M$, $10^{-13}M$, $5 \times 10^{-14}M$, $10^{-14}M$, $5 \times 10^{-15}M$, and $10^{-15}M$.

[683] Moreover, polypeptides of the present invention are useful in inhibiting the angiogenesis of proliferative cells or tissues, either alone, as a protein fusion, or in combination with other polypeptides directly or indirectly, as described elsewhere herein. In a most preferred embodiment, said anti-angiogenesis effect may be achieved indirectly, for example, through the inhibition of hematopoietic, tumor-specific cells, such as tumor-associated macrophages (See Joseph IB, et al. J Natl Cancer Inst, 90(21):1648-53 (1998), which is hereby incorporated by reference). Antibodies directed to polypeptides or polynucleotides of the present invention may also result in inhibition of angiogenesis directly, or indirectly (See Witte L, et al., Cancer Metastasis Rev. 17(2):155-61 (1998), which is hereby incorporated by reference)).

[684] Polypeptides, including protein fusions, of the present invention, or fragments thereof may be useful in inhibiting proliferative cells or tissues through the induction of apoptosis. Said polypeptides may act either directly, or indirectly to induce apoptosis of proliferative cells and tissues, for example in the activation of a death-domain receptor, such as tumor necrosis factor (TNF) receptor-1, CD95 (Fas/APO-1), TNF-receptor-related apoptosis-mediated protein (TRAMP) and TNF-related apoptosis-inducing ligand (TRAIL) receptor-1 and -2 (See Schulze-Osthoff K, et.al., Eur J Biochem 254(3):439-59 (1998), which is hereby incorporated by reference). Moreover, in another preferred embodiment of the present invention, said polypeptides may induce apoptosis through other mechanisms, such as in the activation of other proteins which will activate apoptosis, or through stimulating the expression of said proteins, either alone or in combination with small molecule drugs or adjuvants, such as apoptonin, galectins, thioredoxins, anti-inflammatory proteins (See for example, Mutat Res 400(1-2):447-55 (1998), Med Hypotheses.50(5):423-33 (1998), Chem Biol Interact. Apr 24;111-112:23-34 (1998), J Mol Med.76(6):402-12

(1998), Int J Tissue React;20(1):3-15 (1998), which are all hereby incorporated by reference).

[685] Polypeptides, including protein fusions to, or fragments thereof, of the present invention are useful in inhibiting the metastasis of proliferative cells or tissues. Inhibition may occur as a direct result of administering polypeptides, or antibodies directed to said polypeptides as described elsewhere herein, or indirectly, such as activating the expression of proteins known to inhibit metastasis, for example alpha 4 integrins, (See, e.g., Curr Top Microbiol Immunol 1998;231:125-41, which is hereby incorporated by reference). Such therapeutic affects of the present invention may be achieved either alone, or in combination with small molecule drugs or adjuvants.

[686] In another embodiment, the invention provides a method of delivering compositions containing the polypeptides of the invention (e.g., compositions containing polypeptides or polypeptide antibodies associated with heterologous polypeptides, heterologous nucleic acids, toxins, or prodrugs) to targeted cells expressing the polypeptide of the present invention. Polypeptides or polypeptide antibodies of the invention may be associated with with heterologous polypeptides, heterologous nucleic acids, toxins, or prodrugs via hydrophobic, hydrophilic, ionic and/or covalent interactions.

[687] Polypeptides, protein fusions to, or fragments thereof, of the present invention are useful in enhancing the immunogenicity and/or antigenicity of proliferating cells or tissues, either directly, such as would occur if the polypeptides of the present invention 'vaccinated' the immune response to respond to proliferative antigens and immunogens, or indirectly, such as in activating the expression of proteins known to enhance the immune response (e.g. chemokines), to said antigens and immunogens.

Renal Disorders

[688] Polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention, may be used to treat, prevent, diagnose, and/or prognose disorders of the renal system. Renal disorders which can be diagnosed, prognosed, prevented, and/or treated with compositions of the invention include, but are not limited to, kidney failure, nephritis, blood vessel disorders of kidney, metabolic and congenital kidney disorders, urinary disorders of the kidney, autoimmune disorders, sclerosis and necrosis, electrolyte imbalance, and kidney cancers.

[689] Kidney diseases which can be diagnosed, prognosed, prevented, and/or treated with compositions of the invention include, but are not limited to, acute kidney failure, chronic kidney failure, atheroembolic renal failure, end-stage renal disease, inflammatory diseases of the kidney (e.g., acute glomerulonephritis, postinfectious glomerulonephritis, rapidly progressive glomerulonephritis, nephrotic syndrome, membranous glomerulonephritis, familial nephrotic syndrome, membranoproliferative glomerulonephritis I and II, mesangial proliferative glomerulonephritis, chronic glomerulonephritis, acute tubulointerstitial nephritis, chronic tubulointerstitial nephritis, acute post-streptococcal glomerulonephritis (PSGN), pyelonephritis, lupus nephritis, chronic nephritis, interstitial nephritis, and post-streptococcal glomerulonephritis), blood vessel disorders of the kidneys (e.g., kidney infarction, atheroembolic kidney disease, cortical necrosis, malignant nephrosclerosis, renal vein thrombosis, renal underperfusion, renal retinopathy, renal ischemia-reperfusion, renal artery embolism, and renal artery stenosis), and kidney disorders resulting from urinary tract disease (e.g., pyelonephritis, hydronephrosis, urolithiasis (renal lithiasis, nephrolithiasis), reflux nephropathy, urinary tract infections, urinary retention, and acute or chronic unilateral obstructive uropathy.)

[690] In addition, compositions of the invention can be used to diagnose, prognose, prevent, and/or treat metabolic and congenital disorders of the kidney (e.g., uremia, renal amyloidosis, renal osteodystrophy, renal tubular acidosis, renal glycosuria, nephrogenic diabetes insipidus, cystinuria, Fanconi's syndrome, renal fibrocystic osteosis (renal rickets), Hartnup disease, Bartter's syndrome, Liddle's syndrome, polycystic kidney disease, medullary cystic disease, medullary sponge kidney, Alport's syndrome, nail-patella syndrome, congenital nephrotic syndrome, CRUSH syndrome, horseshoe kidney, diabetic nephropathy, nephrogenic diabetes insipidus, analgesic nephropathy, kidney stones, and membranous nephropathy), and autoimmune disorders of the kidney (e.g., systemic lupus erythematosus (SLE), Goodpasture syndrome, IgA nephropathy, and IgM mesangial proliferative glomerulonephritis).

[691] Compositions of the invention can also be used to diagnose, prognose, prevent, and/or treat sclerotic or necrotic disorders of the kidney (e.g., glomerulosclerosis, diabetic nephropathy, focal segmental glomerulosclerosis (FSGS), necrotizing glomerulonephritis, and renal papillary necrosis), cancers of the kidney (e.g., nephroma, hypernephroma, nephroblastoma, renal cell cancer, transitional cell cancer, renal adenocarcinoma, squamous cell cancer, and Wilm's tumor), and electrolyte imbalances (e.g., nephrocalcinosis, pyuria,

edema, hydronephritis, proteinuria, hyponatremia, hypernatremia, hypokalemia, hyperkalemia, hypocalcemia, hypercalcemia, hypophosphatemia, and hyperphosphatemia).

[692] Polypeptides may be administered using any method known in the art, including, but not limited to, direct needle injection at the delivery site, intravenous injection, topical administration, catheter infusion, biolistic injectors, particle accelerators, gelfoam sponge depots, other commercially available depot materials, osmotic pumps, oral or suppositorial solid pharmaceutical formulations, decanting or topical applications during surgery, aerosol delivery. Such methods are known in the art. Polypeptides may be administered as part of a Therapeutic, described in more detail below. Methods of delivering polynucleotides are described in more detail herein.

Cardiovascular Disorders

[693] Polynucleotides or polypeptides, or agonists or antagonists of the present invention, may be used to treat, prevent, diagnose, and/or prognose cardiovascular disorders, including, but not limited to, peripheral artery disease, such as limb ischemia.

[694] Cardiovascular disorders include, but are not limited to, cardiovascular abnormalities, such as arterio-arterial fistula, arteriovenous fistula, cerebral arteriovenous malformations, congenital heart defects, pulmonary atresia, and Scimitar Syndrome. Congenital heart defects include, but are not limited to, aortic coarctation, cor triatriatum, coronary vessel anomalies, crisscross heart, dextrocardia, patent ductus arteriosus, Ebstein's anomaly, Eisenmenger complex, hypoplastic left heart syndrome, levocardia, tetralogy of fallot, transposition of great vessels, double outlet right ventricle, tricuspid atresia, persistent truncus arteriosus, and heart septal defects, such as aortopulmonary septal defect, endocardial cushion defects, Lutembacher's Syndrome, trilogly of Fallot, ventricular heart septal defects.

[695] Cardiovascular disorders also include, but are not limited to, heart disease, such as arrhythmias, carcinoid heart disease, high cardiac output, low cardiac output, cardiac tamponade, endocarditis (including bacterial), heart aneurysm, cardiac arrest, congestive heart failure, congestive cardiomyopathy, paroxysmal dyspnea, cardiac edema, heart hypertrophy, congestive cardiomyopathy, left ventricular hypertrophy, right ventricular hypertrophy, post-infarction heart rupture, ventricular septal rupture, heart valve diseases, myocardial diseases, myocardial ischemia, pericardial effusion, pericarditis (including

phlebitis, pulmonary veno-occlusive disease, Raynaud's disease, CREST syndrome, retinal vein occlusion, Scimitar syndrome, superior vena cava syndrome, telangiectasia, atacia telangiectasia, hereditary hemorrhagic telangiectasia, varicocele, varicose veins, varicose ulcer, vasculitis, and venous insufficiency.

[701] Aneurysms include, but are not limited to, dissecting aneurysms, false aneurysms, infected aneurysms, ruptured aneurysms, aortic aneurysms, cerebral aneurysms, coronary aneurysms, heart aneurysms, and iliac aneurysms.

[702] Arterial occlusive diseases include, but are not limited to, arteriosclerosis, intermittent claudication, carotid stenosis, fibromuscular dysplasias, mesenteric vascular occlusion, Moyamoya disease, renal artery obstruction, retinal artery occlusion, and thromboangiitis obliterans.

[703] Cerebrovascular disorders include, but are not limited to, carotid artery diseases, cerebral amyloid angiopathy, cerebral aneurysm, cerebral anoxia, cerebral arteriosclerosis, cerebral arteriovenous malformation, cerebral artery diseases, cerebral embolism and thrombosis, carotid artery thrombosis, sinus thrombosis, Wallenberg's syndrome, cerebral hemorrhage, epidural hematoma, subdural hematoma, subarachnoid hemorrhage, cerebral infarction, cerebral ischemia (including transient), subclavian steal syndrome, periventricular leukomalacia, vascular headache, cluster headache, migraine, and vertebrobasilar insufficiency.

[704] Embolisms include, but are not limited to, air embolisms, amniotic fluid embolisms, cholesterol embolisms, blue toe syndrome, fat embolisms, pulmonary embolisms, and thromboembolisms. Thrombosis include, but are not limited to, coronary thrombosis, hepatic vein thrombosis, retinal vein occlusion, carotid artery thrombosis, sinus thrombosis, Wallenberg's syndrome, and thrombophlebitis.

[705] Ischemic disorders include, but are not limited to, cerebral ischemia, ischemic colitis, compartment syndromes, anterior compartment syndrome, myocardial ischemia, reperfusion injuries, and peripheral limb ischemia. Vasculitis includes, but is not limited to, aortitis, arteritis, Behcet's Syndrome, Churg-Strauss Syndrome, mucocutaneous lymph node syndrome, thromboangiitis obliterans, hypersensitivity vasculitis, Schoenlein-Henoch purpura, allergic cutaneous vasculitis, and Wegener's granulomatosis.

[706] Polypeptides may be administered using any method known in the art, including, but not limited to, direct needle injection at the delivery site, intravenous injection, topical administration, catheter infusion, biolistic injectors, particle accelerators, gelfoam sponge

depots, other commercially available depot materials, osmotic pumps, oral or suppositorial solid pharmaceutical formulations, decanting or topical applications during surgery, aerosol delivery. Such methods are known in the art. Polypeptides may be administered as part of a Therapeutic, described in more detail below. Methods of delivering polynucleotides are described in more detail herein.

Respiratory Disorders

[707] Polynucleotides or polypeptides, or agonists or antagonists of the present invention may be used to treat, prevent, diagnose, and/or prognose diseases and/or disorders of the respiratory system.

[708] Diseases and disorders of the respiratory system include, but are not limited to, nasal vestibulitis, nonallergic rhinitis (e.g., acute rhinitis, chronic rhinitis, atrophic rhinitis, vasomotor rhinitis), nasal polyps, and sinusitis, juvenile angiofibromas, cancer of the nose and juvenile papillomas, vocal cord polyps, nodules (singer's nodules), contact ulcers, vocal cord paralysis, laryngoceles, pharyngitis (e.g., viral and bacterial), tonsillitis, tonsillar cellulitis, parapharyngeal abscess, laryngitis, laryngoceles, and throat cancers (e.g., cancer of the nasopharynx, tonsil cancer, larynx cancer), lung cancer (e.g., squamous cell carcinoma, small cell (oat cell) carcinoma, large cell carcinoma, and adenocarcinoma), allergic disorders (eosinophilic pneumonia, hypersensitivity pneumonitis (e.g., extrinsic allergic alveolitis, allergic interstitial pneumonitis, organic dust pneumoconiosis, allergic bronchopulmonary aspergillosis, asthma, Wegener's granulomatosis (granulomatous vasculitis), Goodpasture's syndrome)), pneumonia (e.g., bacterial pneumonia (e.g., *Streptococcus pneumoniae* (pneumococcal pneumonia), *Staphylococcus aureus* (staphylococcal pneumonia), Gram-negative bacterial pneumonia (caused by, e.g., *Klebsiella* and *Pseudomonas spp.*), *Mycoplasma pneumoniae* pneumonia, *Hemophilus influenzae* pneumonia, *Legionella pneumophila* (Legionnaires' disease), and *Chlamydia psittaci* (Psittacosis)), and viral pneumonia (e.g., influenza, chickenpox (varicella).

[709] Additional diseases and disorders of the respiratory system include, but are not limited to bronchiolitis, polio (poliomyelitis), croup, respiratory syncytial viral infection, mumps, erythema infectiosum (fifth disease), roseola infantum, progressive rubella panencephalitis, german measles, and subacute sclerosing panencephalitis), fungal pneumonia (e.g., Histoplasmosis, Coccidioidomycosis, Blastomycosis, fungal infections in

people with severely suppressed immune systems (e.g., cryptococcosis, caused by *Cryptococcus neoformans*; aspergillosis, caused by *Aspergillus spp.*; candidiasis, caused by *Candida*; and mucormycosis)), *Pneumocystis carinii* (pneumocystis pneumonia), atypical pneumonias (e.g., *Mycoplasma* and *Chlamydia spp.*), opportunistic infection pneumonia, nosocomial pneumonia, chemical pneumonitis, and aspiration pneumonia, pleural disorders (e.g., pleurisy, pleural effusion, and pneumothorax (e.g., simple spontaneous pneumothorax, complicated spontaneous pneumothorax, tension pneumothorax)), obstructive airway diseases (e.g., asthma, chronic obstructive pulmonary disease (COPD), emphysema, chronic or acute bronchitis), occupational lung diseases (e.g., silicosis, black lung (coal workers' pneumoconiosis), asbestosis, berylliosis, occupational asthma, byssinosis, and benign pneumoconioses), Infiltrative Lung Disease (e.g., pulmonary fibrosis (e.g., fibrosing alveolitis, usual interstitial pneumonia), idiopathic pulmonary fibrosis, desquamative interstitial pneumonia, lymphoid interstitial pneumonia, histiocytosis X (e.g., Letterer-Siwe disease, Hand-Schüller-Christian disease, eosinophilic granuloma), idiopathic pulmonary hemosiderosis, sarcoidosis and pulmonary alveolar proteinosis), Acute respiratory distress syndrome (also called, e.g., adult respiratory distress syndrome), edema, pulmonary embolism, bronchitis (e.g., viral, bacterial), bronchiectasis, atelectasis, lung abscess (caused by, e.g., *Staphylococcus aureus* or *Legionella pneumophila*), and cystic fibrosis.

Anti-Angiogenesis Activity

[710] The naturally occurring balance between endogenous stimulators and inhibitors of angiogenesis is one in which inhibitory influences predominate. Rastinejad *et al.*, *Cell* 56:345-355 (1989). In those rare instances in which neovascularization occurs under normal physiological conditions, such as wound healing, organ regeneration, embryonic development, and female reproductive processes, angiogenesis is stringently regulated and spatially and temporally delimited. Under conditions of pathological angiogenesis such as that characterizing solid tumor growth, these regulatory controls fail. Unregulated angiogenesis becomes pathologic and sustains progression of many neoplastic and non-neoplastic diseases. A number of serious diseases are dominated by abnormal neovascularization including solid tumor growth and metastases, arthritis, some types of eye disorders, and psoriasis. See, e.g., reviews by Moses *et al.*, *Biotech.* 9:630-634 (1991); Folkman *et al.*, *N. Engl. J. Med.*, 333:1757-1763 (1995); Auerbach *et al.*, *J. Microvasc.*

09950083-091201

Res. 29:401-411 (1985); Folkman, *Advances in Cancer Research*, eds. Klein and Weinhouse, Academic Press, New York, pp. 175-203 (1985); Patz, *Am. J. Ophthalmol.* 94:715-743 (1982); and Folkman *et al.*, *Science* 221:719-725 (1983). In a number of pathological conditions, the process of angiogenesis contributes to the disease state. For example, significant data have accumulated which suggest that the growth of solid tumors is dependent on angiogenesis. Folkman and Klagsbrun, *Science* 235:442-447 (1987).

[711] The present invention provides for treatment of diseases or disorders associated with neovascularization by administration of the polynucleotides and/or polypeptides of the invention, as well as agonists or antagonists of the present invention. Malignant and metastatic conditions which can be treated with the polynucleotides and polypeptides, or agonists or antagonists of the invention include, but are not limited to, malignancies, solid tumors, and cancers described herein and otherwise known in the art (for a review of such disorders, see Fishman *et al.*, *Medicine*, 2d Ed., J. B. Lippincott Co., Philadelphia (1985)). Thus, the present invention provides a method of treating an angiogenesis-related disease and/or disorder, comprising administering to an individual in need thereof a therapeutically effective amount of a polynucleotide, polypeptide, antagonist and/or agonist of the invention. For example, polynucleotides, polypeptides, antagonists and/or agonists may be utilized in a variety of additional methods in order to therapeutically treat a cancer or tumor. Cancers which may be treated with polynucleotides, polypeptides, antagonists and/or agonists include, but are not limited to solid tumors, including prostate, lung, breast, ovarian, stomach, pancreas, larynx, esophagus, testes, liver, parotid, biliary tract, colon, rectum, cervix, uterus, endometrium, kidney, bladder, thyroid cancer; primary tumors and metastases; melanomas; glioblastoma; Kaposi's sarcoma; leiomyosarcoma; non-small cell lung cancer; colorectal cancer; advanced malignancies; and blood born tumors such as leukemias. For example, polynucleotides, polypeptides, antagonists and/or agonists may be delivered topically, in order to treat cancers such as skin cancer, head and neck tumors, breast tumors, and Kaposi's sarcoma.

[712] Within yet other aspects, polynucleotides, polypeptides, antagonists and/or agonists may be utilized to treat superficial forms of bladder cancer by, for example, intravesical administration. Polynucleotides, polypeptides, antagonists and/or agonists may be delivered directly into the tumor, or near the tumor site, via injection or a catheter. Of course, as the artisan of ordinary skill will appreciate, the appropriate mode of administration will vary according to the cancer to be treated. Other modes of delivery are

discussed herein.

[713] Polynucleotides, polypeptides, antagonists and/or agonists may be useful in treating other disorders, besides cancers, which involve angiogenesis. These disorders include, but are not limited to: benign tumors, for example hemangiomas, acoustic neuromas, neurofibromas, trachomas, and pyogenic granulomas; arteriosclerotic plaques; ocular angiogenic diseases, for example, diabetic retinopathy, retinopathy of prematurity, macular degeneration, corneal graft rejection, neovascular glaucoma, retrolental fibroplasia, rubeosis, retinoblastoma, uveitis and Pterygia (abnormal blood vessel growth) of the eye; rheumatoid arthritis; psoriasis; delayed wound healing; endometriosis; vasculogenesis; granulations; hypertrophic scars (keloids); nonunion fractures; scleroderma; trachoma; vascular adhesions; myocardial angiogenesis; coronary collaterals; cerebral collaterals; arteriovenous malformations; ischemic limb angiogenesis; Osler-Webber Syndrome; plaque neovascularization; telangiectasia; hemophilic joints; angiofibroma; fibromuscular dysplasia; wound granulation; Crohn's disease; and atherosclerosis.

[714] For example, within one aspect of the present invention methods are provided for treating hypertrophic scars and keloids, comprising the step of administering a polynucleotide, polypeptide, antagonist and/or agonist of the invention to a hypertrophic scar or keloid.

[715] Within one embodiment of the present invention polynucleotides, polypeptides, antagonists and/or agonists of the invention are directly injected into a hypertrophic scar or keloid, in order to prevent the progression of these lesions. This therapy is of particular value in the prophylactic treatment of conditions which are known to result in the development of hypertrophic scars and keloids (e.g., burns), and is preferably initiated after the proliferative phase has had time to progress (approximately 14 days after the initial injury), but before hypertrophic scar or keloid development. As noted above, the present invention also provides methods for treating neovascular diseases of the eye, including for example, corneal neovascularization, neovascular glaucoma, proliferative diabetic retinopathy, retrolental fibroplasia and macular degeneration.

[716] Moreover, Ocular disorders associated with neovascularization which can be treated with the polynucleotides and polypeptides of the present invention (including agonists and/or antagonists) include, but are not limited to: neovascular glaucoma, diabetic retinopathy, retinoblastoma, retrolental fibroplasia, uveitis, retinopathy of prematurity macular degeneration, corneal graft neovascularization, as well as other eye inflammatory

diseases, ocular tumors and diseases associated with choroidal or iris neovascularization. See, e.g., reviews by Waltman *et al.*, *Am. J. Ophthalm.* 85:704-710 (1978) and Gartner *et al.*, *Surv. Ophthalm.* 22:291-312 (1978).

[717] Thus, within one aspect of the present invention methods are provided for treating neovascular diseases of the eye such as corneal neovascularization (including corneal graft neovascularization), comprising the step of administering to a patient a therapeutically effective amount of a compound (as described above) to the cornea, such that the formation of blood vessels is inhibited. Briefly, the cornea is a tissue which normally lacks blood vessels. In certain pathological conditions however, capillaries may extend into the cornea from the pericorneal vascular plexus of the limbus. When the cornea becomes vascularized, it also becomes clouded, resulting in a decline in the patient's visual acuity. Visual loss may become complete if the cornea completely opacitates. A wide variety of disorders can result in corneal neovascularization, including for example, corneal infections (e.g., trachoma, herpes simplex keratitis, leishmaniasis and onchocerciasis), immunological processes (e.g., graft rejection and Stevens-Johnson's syndrome), alkali burns, trauma, inflammation (of any cause), toxic and nutritional deficiency states, and as a complication of wearing contact lenses.

[718] Within particularly preferred embodiments of the invention, may be prepared for topical administration in saline (combined with any of the preservatives and antimicrobial agents commonly used in ocular preparations), and administered in eyedrop form. The solution or suspension may be prepared in its pure form and administered several times daily. Alternatively, anti-angiogenic compositions, prepared as described above, may also be administered directly to the cornea. Within preferred embodiments, the anti-angiogenic composition is prepared with a muco-adhesive polymer which binds to cornea. Within further embodiments, the anti-angiogenic factors or anti-angiogenic compositions may be utilized as an adjunct to conventional steroid therapy. Topical therapy may also be useful prophylactically in corneal lesions which are known to have a high probability of inducing an angiogenic response (such as chemical burns). In these instances the treatment, likely in combination with steroids, may be instituted immediately to help prevent subsequent complications.

[719] Within other embodiments, the compounds described above may be injected directly into the corneal stroma by an ophthalmologist under microscopic guidance. The preferred site of injection may vary with the morphology of the individual lesion, but the

goal of the administration would be to place the composition at the advancing front of the vasculature (i.e., interspersed between the blood vessels and the normal cornea). In most cases this would involve perilimbic corneal injection to "protect" the cornea from the advancing blood vessels. This method may also be utilized shortly after a corneal insult in order to prophylactically prevent corneal neovascularization. In this situation the material could be injected in the perilimbic cornea interspersed between the corneal lesion and its undesired potential limbic blood supply. Such methods may also be utilized in a similar fashion to prevent capillary invasion of transplanted corneas. In a sustained-release form injections might only be required 2-3 times per year. A steroid could also be added to the injection solution to reduce inflammation resulting from the injection itself.

[720] Within another aspect of the present invention, methods are provided for treating neovascular glaucoma, comprising the step of administering to a patient a therapeutically effective amount of a polynucleotide, polypeptide, antagonist and/or agonist to the eye, such that the formation of blood vessels is inhibited. In one embodiment, the compound may be administered topically to the eye in order to treat early forms of neovascular glaucoma. Within other embodiments, the compound may be implanted by injection into the region of the anterior chamber angle. Within other embodiments, the compound may also be placed in any location such that the compound is continuously released into the aqueous humor. Within another aspect of the present invention, methods are provided for treating proliferative diabetic retinopathy, comprising the step of administering to a patient a therapeutically effective amount of a polynucleotide, polypeptide, antagonist and/or agonist to the eyes, such that the formation of blood vessels is inhibited.

[721] Within particularly preferred embodiments of the invention, proliferative diabetic retinopathy may be treated by injection into the aqueous humor or the vitreous, in order to increase the local concentration of the polynucleotide, polypeptide, antagonist and/or agonist in the retina. Preferably, this treatment should be initiated prior to the acquisition of severe disease requiring photocoagulation.

[722] Within another aspect of the present invention, methods are provided for treating retrolental fibroplasia, comprising the step of administering to a patient a therapeutically effective amount of a polynucleotide, polypeptide, antagonist and/or agonist to the eye, such that the formation of blood vessels is inhibited. The compound may be administered topically, via intravitreal injection and/or via intraocular implants.

[723] Additionally, disorders which can be treated with the polynucleotides,

polypeptides, agonists and/or agonists include, but are not limited to, hemangioma, arthritis, psoriasis, angiofibroma, atherosclerotic plaques, delayed wound healing, granulations, hemophilic joints, hypertrophic scars, nonunion fractures, Osler-Weber syndrome, pyogenic granuloma, scleroderma, trachoma, and vascular adhesions.

[724] Moreover, disorders and/or states, which can be treated, prevented, diagnosed, and/or prognosed with the the polynucleotides, polypeptides, agonists and/or agonists of the invention include, but are not limited to, solid tumors, blood born tumors such as leukemias, tumor metastasis, Kaposi's sarcoma, benign tumors, for example hemangiomas, acoustic neuromas, neurofibromas, trachomas, and pyogenic granulomas, rheumatoid arthritis, psoriasis, ocular angiogenic diseases, for example, diabetic retinopathy, retinopathy of prematurity, macular degeneration, corneal graft rejection, neovascular glaucoma, retrolental fibroplasia, rubeosis, retinoblastoma, and uvietis, delayed wound healing, endometriosis, vasculogenesis, granulations, hypertrophic scars (keloids), nonunion fractures, scleroderma, trachoma, vascular adhesions, myocardial angiogenesis, coronary collaterals, cerebral collaterals, arteriovenous malformations, ischemic limb angiogenesis, Osler-Webber Syndrome, plaque neovascularization, telangiectasia, hemophiliac joints, angiofibroma fibromuscular dysplasia, wound granulation, Crohn's disease, atherosclerosis, birth control agent by preventing vascularization required for embryo implantation controlling menstruation, diseases that have angiogenesis as a pathologic consequence such as cat scratch disease (*Rochela minalia quintosa*), ulcers (*Helicobacter pylori*), Bartonellosis and bacillary angiomatosis.

[725] In one aspect of the birth control method, an amount of the compound sufficient to block embryo implantation is administered before or after intercourse and fertilization have occurred, thus providing an effective method of birth control, possibly a "morning after" method. Polynucleotides, polypeptides, agonists and/or agonists may also be used in controlling menstruation or administered as either a peritoneal lavage fluid or for peritoneal implantation in the treatment of endometriosis.

[726] Polynucleotides, polypeptides, agonists and/or agonists of the present invention may be incorporated into surgical sutures in order to prevent stitch granulomas.

[727] Polynucleotides, polypeptides, agonists and/or agonists may be utilized in a wide variety of surgical procedures. For example, within one aspect of the present invention a compositions (in the form of, for example, a spray or film) may be utilized to coat or spray an area prior to removal of a tumor, in order to isolate normal surrounding tissues from

09950083-091201

malignant tissue, and/or to prevent the spread of disease to surrounding tissues. Within other aspects of the present invention, compositions (e.g., in the form of a spray) may be delivered via endoscopic procedures in order to coat tumors, or inhibit angiogenesis in a desired locale. Within yet other aspects of the present invention, surgical meshes which have been coated with anti-angiogenic compositions of the present invention may be utilized in any procedure wherein a surgical mesh might be utilized. For example, within one embodiment of the invention a surgical mesh laden with an anti-angiogenic composition may be utilized during abdominal cancer resection surgery (e.g., subsequent to colon resection) in order to provide support to the structure, and to release an amount of the anti-angiogenic factor.

[728] Within further aspects of the present invention, methods are provided for treating tumor excision sites, comprising administering a polynucleotide, polypeptide, agonist and/or agonist to the resection margins of a tumor subsequent to excision, such that the local recurrence of cancer and the formation of new blood vessels at the site is inhibited. Within one embodiment of the invention, the anti-angiogenic compound is administered directly to the tumor excision site (e.g., applied by swabbing, brushing or otherwise coating the resection margins of the tumor with the anti-angiogenic compound). Alternatively, the anti-angiogenic compounds may be incorporated into known surgical pastes prior to administration. Within particularly preferred embodiments of the invention, the anti-angiogenic compounds are applied after hepatic resections for malignancy, and after neurosurgical operations.

[729] Within one aspect of the present invention, polynucleotides, polypeptides, agonists and/or agonists may be administered to the resection margin of a wide variety of tumors, including for example, breast, colon, brain and hepatic tumors. For example, within one embodiment of the invention, anti-angiogenic compounds may be administered to the site of a neurological tumor subsequent to excision, such that the formation of new blood vessels at the site are inhibited.

[730] The polynucleotides, polypeptides, agonists and/or agonists of the present invention may also be administered along with other anti-angiogenic factors. Representative examples of other anti-angiogenic factors include: Anti-Invasive Factor, retinoic acid and derivatives thereof, paclitaxel, Suramin, Tissue Inhibitor of Metalloproteinase-1, Tissue Inhibitor of Metalloproteinase-2, Plasminogen Activator

Inhibitor-1, Plasminogen Activator Inhibitor-2, and various forms of the lighter "d group" transition metals.

[731] Lighter "d group" transition metals include, for example, vanadium, molybdenum, tungsten, titanium, niobium, and tantalum species. Such transition metal species may form transition metal complexes. Suitable complexes of the above-mentioned transition metal species include oxo transition metal complexes.

[732] Representative examples of vanadium complexes include oxo vanadium complexes such as vanadate and vanadyl complexes. Suitable vanadate complexes include metavanadate and orthovanadate complexes such as, for example, ammonium metavanadate, sodium metavanadate, and sodium orthovanadate. Suitable vanadyl complexes include, for example, vanadyl acetylacetonate and vanadyl sulfate including vanadyl sulfate hydrates such as vanadyl sulfate mono- and trihydrates.

[733] Representative examples of tungsten and molybdenum complexes also include oxo complexes. Suitable oxo tungsten complexes include tungstate and tungsten oxide complexes. Suitable tungstate complexes include ammonium tungstate, calcium tungstate, sodium tungstate dihydrate, and tungstic acid. Suitable tungsten oxides include tungsten (IV) oxide and tungsten (VI) oxide. Suitable oxo molybdenum complexes include molybdate, molybdenum oxide, and molybdenyl complexes. Suitable molybdate complexes include ammonium molybdate and its hydrates, sodium molybdate and its hydrates, and potassium molybdate and its hydrates. Suitable molybdenum oxides include molybdenum (VI) oxide, molybdenum (VI) oxide, and molybdic acid. Suitable molybdenyl complexes include, for example, molybdenyl acetylacetonate. Other suitable tungsten and molybdenum complexes include hydroxo derivatives derived from, for example, glycerol, tartaric acid, and sugars.

[734] A wide variety of other anti-angiogenic factors may also be utilized within the context of the present invention. Representative examples include platelet factor 4; protamine sulphate; sulphated chitin derivatives (prepared from queen crab shells), (Murata et al., Cancer Res. 51:22-26, 1991); Sulphated Polysaccharide Peptidoglycan Complex (SP-PG) (the function of this compound may be enhanced by the presence of steroids such as estrogen, and tamoxifen citrate); Staurosporine; modulators of matrix metabolism, including for example, proline analogs, cishydroxyproline, d,L-3,4-dehydroproline, Thiaproline, alpha,alpha-dipyridyl, aminopropionitrile fumarate; 4-propyl-5-(4-pyridinyl)-2(3H)-oxazolone; Methotrexate; Mitoxantrone; Heparin; Interferons; 2 Macroglobulin-

serum; ChIMP-3 (Pavloff et al., J. Bio. Chem. 267:17321-17326, 1992); Chymostatin (Tomkinson et al., Biochem J. 286:475-480, 1992); Cyclodextrin Tetradecasulfate; Eponemycin; Camptothecin; Fumagillin (Ingber et al., Nature 348:555-557, 1990); Gold Sodium Thiomalate ("GST"; Matsubara and Ziff, J. Clin. Invest. 79:1440-1446, 1987); anticollagenase-serum; alpha2-antiplasmin (Holmes et al., J. Biol. Chem. 262(4):1659-1664, 1987); Bisantrone (National Cancer Institute); Lobenzarit disodium (N-(2)-carboxyphenyl-4- chloroanthronilic acid disodium or "CCA"; Takeuchi et al., Agents Actions 36:312-316, 1992); Thalidomide; Angostatic steroid; AGM-1470; carboxynaminolmidazole; and metalloproteinase inhibitors such as BB94.

Diseases at the Cellular Level

[735] Diseases associated with increased cell survival or the inhibition of apoptosis that could be treated, prevented, diagnosed, and/or prognosed using polynucleotides or polypeptides, as well as antagonists or agonists of the present invention, include cancers (such as follicular lymphomas, carcinomas with p53 mutations, and hormone-dependent tumors, including, but not limited to colon cancer, cardiac tumors, pancreatic cancer, melanoma, retinoblastoma, glioblastoma, lung cancer, intestinal cancer, testicular cancer, stomach cancer, neuroblastoma, myxoma, myoma, lymphoma, endothelioma, osteoblastoma, osteoclastoma, osteosarcoma, chondrosarcoma, adenoma, breast cancer, prostate cancer, Kaposi's sarcoma and ovarian cancer); autoimmune disorders (such as, multiple sclerosis, Sjogren's syndrome, Hashimoto's thyroiditis, biliary cirrhosis, Behcet's disease, Crohn's disease, polymyositis, systemic lupus erythematosus and immune-related glomerulonephritis and rheumatoid arthritis) and viral infections (such as herpes viruses, pox viruses and adenoviruses), inflammation, graft v. host disease, acute graft rejection, and chronic graft rejection.

[736] In preferred embodiments, polynucleotides, polypeptides, and/or antagonists of the invention are used to inhibit growth, progression, and/or metasis of cancers, in particular those listed above.

[737] Additional diseases or conditions associated with increased cell survival that could be treated or detected by polynucleotides or polypeptides, or agonists or antagonists of the present invention include, but are not limited to, progression, and/or metastases of malignancies and related disorders such as leukemia (including acute leukemias (e.g., acute lymphocytic leukemia, acute myelocytic leukemia (including myeloblastic, promyelocytic,

myelomonocytic, monocytic, and erythroleukemia)) and chronic leukemias (e.g., chronic myelocytic (granulocytic) leukemia and chronic lymphocytic leukemia)), polycythemia vera, lymphomas (e.g., Hodgkin's disease and non-Hodgkin's disease), multiple myeloma, Waldenstrom's macroglobulinemia, heavy chain disease, and solid tumors including, but not limited to, sarcomas and carcinomas such as fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, osteogenic sarcoma, chordoma, angiosarcoma, endotheliosarcoma, lymphangiosarcoma, lymphangioendotheliosarcoma, synovioma, mesothelioma, Ewing's tumor, leiomyosarcoma, rhabdomyosarcoma, colon carcinoma, pancreatic cancer, breast cancer, ovarian cancer, prostate cancer, squamous cell carcinoma, basal cell carcinoma, adenocarcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma, papillary adenocarcinomas, cystadenocarcinoma, medullary carcinoma, bronchogenic carcinoma, renal cell carcinoma, hepatoma, bile duct carcinoma, choriocarcinoma, seminoma, embryonal carcinoma, Wilm's tumor, cervical cancer, testicular tumor, lung carcinoma, small cell lung carcinoma, bladder carcinoma, epithelial carcinoma, glioma, astrocytoma, medulloblastoma, craniopharyngioma, ependymoma, pinealoma, hemangioblastoma, acoustic neuroma, oligodendroglioma, menangioma, melanoma, neuroblastoma, and retinoblastoma.

[738] Diseases associated with increased apoptosis that could be treated, prevented, diagnosed, and/or prognosed using polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, include, but are not limited to, AIDS; neurodegenerative disorders (such as Alzheimer's disease, Parkinson's disease, Amyotrophic lateral sclerosis, Retinitis pigmentosa, Cerebellar degeneration and brain tumor or prior associated disease); autoimmune disorders (such as, multiple sclerosis, Sjogren's syndrome, Hashimoto's thyroiditis, biliary cirrhosis, Behcet's disease, Crohn's disease, polymyositis, systemic lupus erythematosus and immune-related glomerulonephritis and rheumatoid arthritis) myelodysplastic syndromes (such as aplastic anemia), graft v. host disease, ischemic injury (such as that caused by myocardial infarction, stroke and reperfusion injury), liver injury (e.g., hepatitis related liver injury, ischemia/reperfusion injury, cholestasis (bile duct injury) and liver cancer); toxin-induced liver disease (such as that caused by alcohol), septic shock, cachexia and anorexia.

Wound Healing and Epithelial Cell Proliferation

[739] In accordance with yet a further aspect of the present invention, there is provided a process for utilizing polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, for therapeutic purposes, for example, to stimulate epithelial cell proliferation and basal keratinocytes for the purpose of wound healing, and to stimulate hair follicle production and healing of dermal wounds. Polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, may be clinically useful in stimulating wound healing including surgical wounds, excisional wounds, deep wounds involving damage of the dermis and epidermis, eye tissue wounds, dental tissue wounds, oral cavity wounds, diabetic ulcers, dermal ulcers, cubitus ulcers, arterial ulcers, venous stasis ulcers, burns resulting from heat exposure or chemicals, and other abnormal wound healing conditions such as uremia, malnutrition, vitamin deficiencies and complications associated with systemic treatment with steroids, radiation therapy and antineoplastic drugs and antimetabolites. Polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, could be used to promote dermal reestablishment subsequent to dermal loss

[740] Polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, could be used to increase the adherence of skin grafts to a wound bed and to stimulate re-epithelialization from the wound bed. The following are types of grafts that polynucleotides or polypeptides, agonists or antagonists of the present invention, could be used to increase adherence to a wound bed: autografts, artificial skin, allografts, autodermic graft, autoepdermic grafts, avacular grafts, Blair-Brown grafts, bone graft, brephoplastic grafts, cutis graft, delayed graft, dermic graft, epidermic graft, fascia graft, full thickness graft, heterologous graft, xenograft, homologous graft, hyperplastic graft, lamellar graft, mesh graft, mucosal graft, Ollier-Thiersch graft, omenpal graft, patch graft, pedicle graft, penetrating graft, split skin graft, thick split graft. Polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, can be used to promote skin strength and to improve the appearance of aged skin.

[741] It is believed that polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, will also produce changes in hepatocyte proliferation, and epithelial cell proliferation in the lung, breast, pancreas, stomach, small intestine, and large intestine. Polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, could promote proliferation of epithelial cells such as sebocytes, hair

follicles, hepatocytes, type II pneumocytes, mucin-producing goblet cells, and other epithelial cells and their progenitors contained within the skin, lung, liver, and gastrointestinal tract. Polynucleotides or polypeptides, agonists or antagonists of the present invention, may promote proliferation of endothelial cells, keratinocytes, and basal keratinocytes.

[742] Polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, could also be used to reduce the side effects of gut toxicity that result from radiation, chemotherapy treatments or viral infections. Polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, may have a cytoprotective effect on the small intestine mucosa. Polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, may also stimulate healing of mucositis (mouth ulcers) that result from chemotherapy and viral infections.

[743] Polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, could further be used in full regeneration of skin in full and partial thickness skin defects, including burns, (i.e., repopulation of hair follicles, sweat glands, and sebaceous glands), treatment of other skin defects such as psoriasis. Polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, could be used to treat epidermolysis bullosa, a defect in adherence of the epidermis to the underlying dermis which results in frequent, open and painful blisters by accelerating reepithelialization of these lesions. Polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, could also be used to treat gastric and duodenal ulcers and help heal by scar formation of the mucosal lining and regeneration of glandular mucosa and duodenal mucosal lining more rapidly. Inflammatory bowel diseases, such as Crohn's disease and ulcerative colitis, are diseases which result in destruction of the mucosal surface of the small or large intestine, respectively. Thus, polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, could be used to promote the resurfacing of the mucosal surface to aid more rapid healing and to prevent progression of inflammatory bowel disease. Treatment with polynucleotides or polypeptides, agonists or antagonists of the present invention, is expected to have a significant effect on the production of mucus throughout the gastrointestinal tract and could be used to protect the intestinal mucosa from injurious substances that are ingested or following surgery. Polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, could be used to treat diseases associate with the under expression.

[744] Moreover, polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, could be used to prevent and heal damage to the lungs due to various pathological states. Polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, which could stimulate proliferation and differentiation and promote the repair of alveoli and bronchiolar epithelium to prevent or treat acute or chronic lung damage. For example, emphysema, which results in the progressive loss of aveoli, and inhalation injuries, i.e., resulting from smoke inhalation and burns, that cause necrosis of the bronchiolar epithelium and alveoli could be effectively treated using polynucleotides or polypeptides, agonists or antagonists of the present invention. Also, polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, could be used to stimulate the proliferation of and differentiation of type II pneumocytes, which may help treat or prevent disease such as hyaline membrane diseases, such as infant respiratory distress syndrome and bronchopulmonary dysplasia, in premature infants.

[745] Polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, could stimulate the proliferation and differentiation of hepatocytes and, thus, could be used to alleviate or treat liver diseases and pathologies such as fulminant liver failure caused by cirrhosis, liver damage caused by viral hepatitis and toxic substances (i.e., acetaminophen, carbon tetrachloride and other hepatotoxins known in the art).

[746] In addition, polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, could be used treat or prevent the onset of diabetes mellitus. In patients with newly diagnosed Types I and II diabetes, where some islet cell function remains, polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, could be used to maintain the islet function so as to alleviate, delay or prevent permanent manifestation of the disease. Also, polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, could be used as an auxiliary in islet cell transplantation to improve or promote islet cell function.

Neural Activity and Neurological Diseases

[747] The polynucleotides, polypeptides and agonists or antagonists of the invention may be used for the diagnosis and/or treatment of diseases, disorders, damage or injury of the brain and/or nervous system. Nervous system disorders that can be treated with the compositions of the invention (e.g., polypeptides, polynucleotides, and/or agonists or antagonists), include, but are not limited to, nervous system injuries, and diseases or

disorders which result in either a disconnection of axons, a diminution or degeneration of neurons, or demyelination. Nervous system lesions which may be treated in a patient (including human and non-human mammalian patients) according to the methods of the invention, include but are not limited to, the following lesions of either the central (including spinal cord, brain) or peripheral nervous systems: (1) ischemic lesions, in which a lack of oxygen in a portion of the nervous system results in neuronal injury or death, including cerebral infarction or ischemia, or spinal cord infarction or ischemia; (2) traumatic lesions, including lesions caused by physical injury or associated with surgery, for example, lesions which sever a portion of the nervous system, or compression injuries; (3) malignant lesions, in which a portion of the nervous system is destroyed or injured by malignant tissue which is either a nervous system associated malignancy or a malignancy derived from non-nervous system tissue; (4) infectious lesions, in which a portion of the nervous system is destroyed or injured as a result of infection, for example, by an abscess or associated with infection by human immunodeficiency virus, herpes zoster, or herpes simplex virus or with Lyme disease, tuberculosis, or syphilis; (5) degenerative lesions, in which a portion of the nervous system is destroyed or injured as a result of a degenerative process including but not limited to, degeneration associated with Parkinson's disease, Alzheimer's disease, Huntington's chorea, or amyotrophic lateral sclerosis (ALS); (6) lesions associated with nutritional diseases or disorders, in which a portion of the nervous system is destroyed or injured by a nutritional disorder or disorder of metabolism including, but not limited to, vitamin B12 deficiency, folic acid deficiency, Wernicke disease, tobacco-alcohol amblyopia, Marchiafava-Bignami disease (primary degeneration of the corpus callosum), and alcoholic cerebellar degeneration; (7) neurological lesions associated with systemic diseases including, but not limited to, diabetes (diabetic neuropathy, Bell's palsy), systemic lupus erythematosus, carcinoma, or sarcoidosis; (8) lesions caused by toxic substances including alcohol, lead, or particular neurotoxins; and (9) demyelinated lesions in which a portion of the nervous system is destroyed or injured by a demyelinating disease including, but not limited to, multiple sclerosis, human immunodeficiency virus-associated myelopathy, transverse myelopathy or various etiologies, progressive multifocal leukoencephalopathy, and central pontine myelinolysis.

[748] In one embodiment, the polypeptides, polynucleotides, or agonists or antagonists of the invention are used to protect neural cells from the damaging effects of hypoxia. In a further preferred embodiment, the polypeptides, polynucleotides, or agonists or antagonists

of the invention are used to protect neural cells from the damaging effects of cerebral hypoxia. According to this embodiment, the compositions of the invention are used to treat or prevent neural cell injury associated with cerebral hypoxia. In one non-exclusive aspect of this embodiment, the polypeptides, polynucleotides, or agonists or antagonists of the invention, are used to treat or prevent neural cell injury associated with cerebral ischemia. In another non-exclusive aspect of this embodiment, the polypeptides, polynucleotides, or agonists or antagonists of the invention are used to treat or prevent neural cell injury associated with cerebral infarction.

[749] In another preferred embodiment, the polypeptides, polynucleotides, or agonists or antagonists of the invention are used to treat or prevent neural cell injury associated with a stroke. In a specific embodiment, the polypeptides, polynucleotides, or agonists or antagonists of the invention are used to treat or prevent cerebral neural cell injury associated with a stroke.

[750] In another preferred embodiment, the polypeptides, polynucleotides, or agonists or antagonists of the invention are used to treat or prevent neural cell injury associated with a heart attack. In a specific embodiment, the polypeptides, polynucleotides, or agonists or antagonists of the invention are used to treat or prevent cerebral neural cell injury associated with a heart attack.

[751] The compositions of the invention which are useful for treating or preventing a nervous system disorder may be selected by testing for biological activity in promoting the survival or differentiation of neurons. For example, and not by way of limitation, compositions of the invention which elicit any of the following effects may be useful according to the invention: (1) increased survival time of neurons in culture either in the presence or absence of hypoxia or hypoxic conditions; (2) increased sprouting of neurons in culture or *in vivo*; (3) increased production of a neuron-associated molecule in culture or *in vivo*, e.g., choline acetyltransferase or acetylcholinesterase with respect to motor neurons; or (4) decreased symptoms of neuron dysfunction *in vivo*. Such effects may be measured by any method known in the art. In preferred, non-limiting embodiments, increased survival of neurons may routinely be measured using a method set forth herein or otherwise known in the art, such as, for example, in Zhang *et al.*, *Proc Natl Acad Sci USA* 97:3637-42 (2000) or in Arakawa *et al.*, *J. Neurosci.*, 10:3507-15 (1990); increased sprouting of neurons may be detected by methods known in the art, such as, for example, the methods set forth in Pestronk *et al.*, *Exp. Neurol.*, 70:65-82 (1980), or Brown *et al.*, *Ann. Rev.*

Neurosci., 4:17-42 (1981); increased production of neuron-associated molecules may be measured by bioassay, enzymatic assay, antibody binding, Northern blot assay, etc., using techniques known in the art and depending on the molecule to be measured; and motor neuron dysfunction may be measured by assessing the physical manifestation of motor neuron disorder, e.g., weakness, motor neuron conduction velocity, or functional disability.

[752] In specific embodiments, motor neuron disorders that may be treated according to the invention include, but are not limited to, disorders such as infarction, infection, exposure to toxin, trauma, surgical damage, degenerative disease or malignancy that may affect motor neurons as well as other components of the nervous system, as well as disorders that selectively affect neurons such as amyotrophic lateral sclerosis, and including, but not limited to, progressive spinal muscular atrophy, progressive bulbar palsy, primary lateral sclerosis, infantile and juvenile muscular atrophy, progressive bulbar paralysis of childhood (Fazio-Londe syndrome), poliomyelitis and the post polio syndrome, and Hereditary Motorsensory Neuropathy (Charcot-Marie-Tooth Disease).

[753] Further, polypeptides or polynucleotides of the invention may play a role in neuronal survival; synapse formation; conductance; neural differentiation, etc. Thus, compositions of the invention (including polynucleotides, polypeptides, and agonists or antagonists) may be used to diagnose and/or treat or prevent diseases or disorders associated with these roles, including, but not limited to, learning and/or cognition disorders. The compositions of the invention may also be useful in the treatment or prevention of neurodegenerative disease states and/or behavioural disorders. Such neurodegenerative disease states and/or behavioral disorders include, but are not limited to, Alzheimer's Disease, Parkinson's Disease, Huntington's Disease, Tourette Syndrome, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder, panic disorder, learning disabilities, ALS, psychoses, autism, and altered behaviors, including disorders in feeding, sleep patterns, balance, and perception. In addition, compositions of the invention may also play a role in the treatment, prevention and/or detection of developmental disorders associated with the developing embryo, or sexually-linked disorders.

[754] Additionally, polypeptides, polynucleotides and/or agonists or antagonists of the invention, may be useful in protecting neural cells from diseases, damage, disorders, or injury, associated with cerebrovascular disorders including, but not limited to, carotid artery diseases (e.g., carotid artery thrombosis, carotid stenosis, or Moyamoya Disease), cerebral amyloid angiopathy, cerebral aneurysm, cerebral anoxia, cerebral arteriosclerosis, cerebral

arteriovenous malformations, cerebral artery diseases, cerebral embolism and thrombosis (e.g., carotid artery thrombosis, sinus thrombosis, or Wallenberg's Syndrome), cerebral hemorrhage (e.g., epidural or subdural hematoma, or subarachnoid hemorrhage), cerebral infarction, cerebral ischemia (e.g., transient cerebral ischemia, Subclavian Steal Syndrome, or vertebrobasilar insufficiency), vascular dementia (e.g., multi-infarct), leukomalacia, periventricular, and vascular headache (e.g., cluster headache or migraines).

[755] In accordance with yet a further aspect of the present invention, there is provided a process for utilizing polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, for therapeutic purposes, for example, to stimulate neurological cell proliferation and/or differentiation. Therefore, polynucleotides, polypeptides, agonists and/or antagonists of the invention may be used to treat and/or detect neurologic diseases. Moreover, polynucleotides or polypeptides, or agonists or antagonists of the invention, can be used as a marker or detector of a particular nervous system disease or disorder.

[756] Examples of neurologic diseases which can be treated or detected with polynucleotides, polypeptides, agonists, and/or antagonists of the present invention include brain diseases, such as metabolic brain diseases which includes phenylketonuria such as maternal phenylketonuria, pyruvate carboxylase deficiency, pyruvate dehydrogenase complex deficiency, Wernicke's Encephalopathy, brain edema, brain neoplasms such as cerebellar neoplasms which include infratentorial neoplasms, cerebral ventricle neoplasms such as choroid plexus neoplasms, hypothalamic neoplasms, supratentorial neoplasms, canavan disease, cerebellar diseases such as cerebellar ataxia which include spinocerebellar degeneration such as ataxia telangiectasia, cerebellar dyssynergia, Friederich's Ataxia, Machado-Joseph Disease, olivopontocerebellar atrophy, cerebellar neoplasms such as infratentorial neoplasms, diffuse cerebral sclerosis such as encephalitis periaxialis, globoid cell leukodystrophy, metachromatic leukodystrophy and subacute sclerosing panencephalitis.

[757] Additional neurologic diseases which can be treated or detected with polynucleotides, polypeptides, agonists, and/or antagonists of the present invention include cerebrovascular disorders (such as carotid artery diseases which include carotid artery thrombosis, carotid stenosis and Moyamoya Disease), cerebral amyloid angiopathy, cerebral aneurysm, cerebral anoxia, cerebral arteriosclerosis, cerebral arteriovenous malformations, cerebral artery diseases, cerebral embolism and thrombosis such as carotid artery thrombosis, sinus thrombosis and Wallenberg's Syndrome, cerebral hemorrhage such

as epidural hematoma, subdural hematoma and subarachnoid hemorrhage, cerebral infarction, cerebral ischemia such as transient cerebral ischemia, Subclavian Steal Syndrome and vertebrobasilar insufficiency, vascular dementia such as multi-infarct dementia, periventricular leukomalacia, vascular headache such as cluster headache and migraine.

[758] Additional neurologic diseases which can be treated or detected with polynucleotides, polypeptides, agonists, and/or antagonists of the present invention include dementia such as AIDS Dementia Complex, presenile dementia such as Alzheimer's Disease and Creutzfeldt-Jakob Syndrome, senile dementia such as Alzheimer's Disease and progressive supranuclear palsy, vascular dementia such as multi-infarct dementia, encephalitis which include encephalitis periaxialis, viral encephalitis such as epidemic encephalitis, Japanese Encephalitis, St. Louis Encephalitis, tick-borne encephalitis and West Nile Fever, acute disseminated encephalomyelitis, meningoencephalitis such as uveomeningoencephalitic syndrome, Postencephalitic Parkinson Disease and subacute sclerosing panencephalitis, encephalomalacia such as periventricular leukomalacia, epilepsy such as generalized epilepsy which includes infantile spasms, absence epilepsy, myoclonic epilepsy which includes MERRF Syndrome, tonic-clonic epilepsy, partial epilepsy such as complex partial epilepsy, frontal lobe epilepsy and temporal lobe epilepsy, post-traumatic epilepsy, status epilepticus such as Epilepsia Partialis Continua, and Hallervorden-Spatz Syndrome.

[759] Additional neurologic diseases which can be treated or detected with polynucleotides, polypeptides, agonists, and/or antagonists of the present invention include hydrocephalus such as Dandy-Walker Syndrome and normal pressure hydrocephalus, hypothalamic diseases such as hypothalamic neoplasms, cerebral malaria, narcolepsy which includes cataplexy, bulbar poliomyelitis, cerebri pseudotumor, Rett Syndrome, Reye's Syndrome, thalamic diseases, cerebral toxoplasmosis, intracranial tuberculoma and Zellweger Syndrome, central nervous system infections such as AIDS Dementia Complex, Brain Abscess, subdural empyema, encephalomyelitis such as Equine Encephalomyelitis, Venezuelan Equine Encephalomyelitis, Necrotizing Hemorrhagic Encephalomyelitis, Visna, and cerebral malaria.

[760] Additional neurologic diseases which can be treated or detected with polynucleotides, polypeptides, agonists, and/or antagonists of the present invention include meningitis such as arachnoiditis, aseptic meningitis such as viral meningitis which

includes lymphocytic choriomeningitis, Bacterial meningitis which includes Haemophilus Meningitis, Listeria Meningitis, Meningococcal Meningitis such as Waterhouse-Friderichsen Syndrome, Pneumococcal Meningitis and meningeal tuberculosis, fungal meningitis such as Cryptococcal Meningitis, subdural effusion, meningoencephalitis such as uve-meningoencephalitic syndrome, myelitis such as transverse myelitis, neurosyphilis such as tabes dorsalis, poliomyelitis which includes bulbar poliomyelitis and postpoliomyelitis syndrome, prion diseases (such as Creutzfeldt-Jakob Syndrome, Bovine Spongiform Encephalopathy, Gerstmann-Straussler Syndrome, Kuru, Scrapie), and cerebral toxoplasmosis.

[761] Additional neurologic diseases which can be treated or detected with polynucleotides, polypeptides, agonists, and/or antagonists of the present invention include central nervous system neoplasms such as brain neoplasms that include cerebellar neoplasms such as infratentorial neoplasms, cerebral ventricle neoplasms such as choroid plexus neoplasms, hypothalamic neoplasms and supratentorial neoplasms, meningeal neoplasms, spinal cord neoplasms which include epidural neoplasms, demyelinating diseases such as Canavan Diseases, diffuse cerebral sclerolysis which includes adrenoleukodystrophy, encephalitis periaxialis, globoid cell leukodystrophy, diffuse cerebral sclerosis such as metachromatic leukodystrophy, allergic encephalomyelitis, necrotizing hemorrhagic encephalomyelitis, progressive multifocal leukoencephalopathy, multiple sclerosis, central pontine myelinolysis, transverse myelitis, neuromyelitis optica, Scrapie, Swayback, Chronic Fatigue Syndrome, Visna, High Pressure Nervous Syndrome, Meningism, spinal cord diseases such as amyotonia congenita, amyotrophic lateral sclerosis, spinal muscular atrophy such as Werdnig-Hoffmann Disease, spinal cord compression, spinal cord neoplasms such as epidural neoplasms, syringomyelia, Tabes Dorsalis, Stiff-Man Syndrome, mental retardation such as Angelman Syndrome, Cri-du-Chat Syndrome, De Lange's Syndrome, Down Syndrome, Gangliosidoses such as gangliosidoses G(M1), Sandhoff Disease, Tay-Sachs Disease, Hartnup Disease, homocystinuria, Laurence-Moon-Biedl Syndrome, Lesch-Nyhan Syndrome, Maple Syrup Urine Disease, mucopolysaccharidosis such as fucosidosis, neuronal ceroid-lipofuscinosis, oculocerebrorenal syndrome, phenylketonuria such as maternal phenylketonuria, Prader-Willi Syndrome, Rett Syndrome, Rubinstein-Taybi Syndrome, Tuberous Sclerosis, WAGR Syndrome, nervous system abnormalities such as holoprosencephaly, neural tube defects such as anencephaly which includes hydranencephaly, Arnold-Chiari Deformity,

encephalocele, meningocele, meningomyelocele, spinal dysraphism such as spina bifida cystica and spina bifida occulta.

[762] Additional neurologic diseases which can be treated or detected with polynucleotides, polypeptides, agonists, and/or antagonists of the present invention include hereditary motor and sensory neuropathies which include Charcot-Marie Disease, Hereditary optic atrophy, Refsum's Disease, hereditary spastic paraplegia, Werdnig-Hoffmann Disease, Hereditary Sensory and Autonomic Neuropathies such as Congenital Analgesia and Familial Dysautonomia, Neurologic manifestations (such as agnosia that include Gerstmann's Syndrome, Amnesia such as retrograde amnesia, apraxia, neurogenic bladder, cataplexy, communicative disorders such as hearing disorders that includes deafness, partial hearing loss, loudness recruitment and tinnitus, language disorders such as aphasia which include agraphia, anomia, broca aphasia, and Wernicke Aphasia, Dyslexia such as Acquired Dyslexia, language development disorders, speech disorders such as aphasia which includes anomia, broca aphasia and Wernicke Aphasia, articulation disorders, communicative disorders such as speech disorders which include dysarthria, echolalia, mutism and stuttering, voice disorders such as aphonia and hoarseness, decerebrate state, delirium, fasciculation, hallucinations, meningism, movement disorders such as angelman syndrome, ataxia, athetosis, chorea, dystonia, hypokinesia, muscle hypotonia, myoclonus, tic, torticollis and tremor, muscle hypertonia such as muscle rigidity such as stiff-man syndrome, muscle spasticity, paralysis such as facial paralysis which includes Herpes Zoster Oticus, Gastroparesis, Hemiplegia, ophthalmoplegia such as diplopia, Duane's Syndrome, Horner's Syndrome, Chronic progressive external ophthalmoplegia such as Kearns Syndrome, Bulbar Paralysis, Tropical Spastic Paraparesis, Paraplegia such as Brown-Sequard Syndrome, quadriplegia, respiratory paralysis and vocal cord paralysis, paresis, phantom limb, taste disorders such as ageusia and dysgeusia, vision disorders such as amblyopia, blindness, color vision defects, diplopia, hemianopsia, scotoma and subnormal vision, sleep disorders such as hypersomnia which includes Kleine-Levin Syndrome, insomnia, and somnambulism, spasm such as trismus, unconsciousness such as coma, persistent vegetative state and syncope and vertigo, neuromuscular diseases such as amyotonia congenita, amyotrophic lateral sclerosis, Lambert-Eaton Myasthenic Syndrome, motor neuron disease, muscular atrophy such as spinal muscular atrophy, Charcot-Marie Disease and Werdnig-Hoffmann Disease, Postpoliomyelitis Syndrome, Muscular Dystrophy, Myasthenia Gravis, Myotonia Atrophica, Myotonia Confenita,

Nemaline Myopathy, Familial Periodic Paralysis, Multiplex Paramyoclonus, Tropical Spastic Paraparesis and Stiff-Man Syndrome, peripheral nervous system diseases such as acrodynia, amyloid neuropathies, autonomic nervous system diseases such as Adie's Syndrome, Barre-Lieou Syndrome, Familial Dysautonomia, Horner's Syndrome, Reflex Sympathetic Dystrophy and Shy-Drager Syndrome, Cranial Nerve Diseases such as Acoustic Nerve Diseases such as Acoustic Neuroma which includes Neurofibromatosis 2, Facial Nerve Diseases such as Facial Neuralgia, Melkersson-Rosenthal Syndrome, ocular motility disorders which includes amblyopia, nystagmus, oculomotor nerve paralysis, ophthalmoplegia such as Duane's Syndrome, Horner's Syndrome, Chronic Progressive External Ophthalmoplegia which includes Kearns Syndrome, Strabismus such as Esotropia and Exotropia, Oculomotor Nerve Paralysis, Optic Nerve Diseases such as Optic Atrophy which includes Hereditary Optic Atrophy, Optic Disk Drusen, Optic Neuritis such as Neuromyelitis Optica, Papilledema, Trigeminal Neuralgia, Vocal Cord Paralysis, Demyelinating Diseases such as Neuromyelitis Optica and Swayback, and Diabetic neuropathies such as diabetic foot.

[763] Additional neurologic diseases which can be treated or detected with polynucleotides, polypeptides, agonists, and/or antagonists of the present invention include nerve compression syndromes such as carpal tunnel syndrome, tarsal tunnel syndrome, thoracic outlet syndrome such as cervical rib syndrome, ulnar nerve compression syndrome, neuralgia such as causalgia, cervico-brachial neuralgia, facial neuralgia and trigeminal neuralgia, neuritis such as experimental allergic neuritis, optic neuritis, polyneuritis, polyradiculoneuritis and radiculitis such as polyradiculitis, hereditary motor and sensory neuropathies such as Charcot-Marie Disease, Hereditary Optic Atrophy, Refsum's Disease, Hereditary Spastic Paraplegia and Werdnig-Hoffmann Disease, Hereditary Sensory and Autonomic Neuropathies which include Congenital Analgesia and Familial Dysautonomia, POEMS Syndrome, Sciatica, Gustatory Sweating and Tetany).

Endocrine Disorders

[764] Polynucleotides or polypeptides, or agonists or antagonists of the present invention, may be used to treat, prevent, diagnose, and/or prognose disorders and/or diseases related to hormone imbalance, and/or disorders or diseases of the endocrine system.

[765] Hormones secreted by the glands of the endocrine system control physical growth, sexual function, metabolism, and other functions. Disorders may be classified in two ways: disturbances in the production of hormones, and the inability of tissues to respond to hormones. The etiology of these hormone imbalance or endocrine system diseases, disorders or conditions may be genetic, somatic, such as cancer and some autoimmune diseases, acquired (e.g., by chemotherapy, injury or toxins), or infectious. Moreover, polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention can be used as a marker or detector of a particular disease or disorder related to the endocrine system and/or hormone imbalance.

[766] Endocrine system and/or hormone imbalance and/or diseases encompass disorders of uterine motility including, but not limited to: complications with pregnancy and labor (e.g., pre-term labor, post-term pregnancy, spontaneous abortion, and slow or stopped labor); and disorders and/or diseases of the menstrual cycle (e.g., dysmenorrhea and endometriosis).

[767] Endocrine system and/or hormone imbalance disorders and/or diseases include disorders and/or diseases of the pancreas, such as, for example, diabetes mellitus, diabetes insipidus, congenital pancreatic agenesis, pheochromocytoma--islet cell tumor syndrome; disorders and/or diseases of the adrenal glands such as, for example, Addison's Disease, corticosteroid deficiency, virilizing disease, hirsutism, Cushing's Syndrome, hyperaldosteronism, pheochromocytoma; disorders and/or diseases of the pituitary gland, such as, for example, hyperpituitarism, hypopituitarism, pituitary dwarfism, pituitary adenoma, panhypopituitarism, acromegaly, gigantism; disorders and/or diseases of the thyroid, including but not limited to, hyperthyroidism, hypothyroidism, Plummer's disease, Graves' disease (toxic diffuse goiter), toxic nodular goiter, thyroiditis (Hashimoto's thyroiditis, subacute granulomatous thyroiditis, and silent lymphocytic thyroiditis), Pendred's syndrome, myxedema, cretinism, thyrotoxicosis, thyroid hormone coupling defect, thymic aplasia, Hurthle cell tumours of the thyroid, thyroid cancer, thyroid carcinoma, Medullary thyroid carcinoma; disorders and/or diseases of the parathyroid, such as, for example, hyperparathyroidism, hypoparathyroidism; disorders and/or diseases of the hypothalamus.

[768] In addition, endocrine system and/or hormone imbalance disorders and/or diseases may also include disorders and/or diseases of the testes or ovaries, including cancer. Other disorders and/or diseases of the testes or ovaries further include, for example,

ovarian cancer, polycystic ovary syndrome, Klinefelter's syndrome, vanishing testes syndrome (bilateral anorchia), congenital absence of Leydig's cells, cryptorchidism, Noonan's syndrome, myotonic dystrophy, capillary haemangioma of the testis (benign), neoplasias of the testis and neo-testis.

[769] Moreover, endocrine system and/or hormone imbalance disorders and/or diseases may also include disorders and/or diseases such as, for example, polyglandular deficiency syndromes, pheochromocytoma, neuroblastoma, multiple Endocrine neoplasia, and disorders and/or cancers of endocrine tissues.

[770] In another embodiment, a polypeptide of the invention, or polynucleotides, antibodies, agonists, or antagonists corresponding to that polypeptide, may be used to diagnose, prognose, prevent, and/or treat endocrine diseases and/or disorders associated with the tissue(s) in which the polypeptide of the invention is expressed, including one, two, three, four, five, or more tissues disclosed in Table 1B, column 8 (Tissue Distribution Library Code).

Reproductive System Disorders

[771] The polynucleotides or polypeptides, or agonists or antagonists of the invention may be used for the diagnosis, treatment, or prevention of diseases and/or disorders of the reproductive system. Reproductive system disorders that can be treated by the compositions of the invention, include, but are not limited to, reproductive system injuries, infections, neoplastic disorders, congenital defects, and diseases or disorders which result in infertility, complications with pregnancy, labor, or parturition, and postpartum difficulties.

[772] Reproductive system disorders and/or diseases include diseases and/or disorders of the testes, including testicular atrophy, testicular feminization, cryptorchism (unilateral and bilateral), anorchia, ectopic testis, epididymitis and orchitis (typically resulting from infections such as, for example, gonorrhea, mumps, tuberculosis, and syphilis), testicular torsion, vasitis nodosa, germ cell tumors (e.g., seminomas, embryonal cell carcinomas, teratocarcinomas, choriocarcinomas, yolk sac tumors, and teratomas), stromal tumors (e.g., Leydig cell tumors), hydrocele, hematocele, varicocele, spermatocele, inguinal hernia, and disorders of sperm production (e.g., immotile cilia syndrome, aspermia, asthenozoospermia, azoospermia, oligospermia, and teratozoospermia).

[773] Reproductive system disorders also include disorders of the prostate gland, such as acute non-bacterial prostatitis, chronic non-bacterial prostatitis, acute bacterial prostatitis, chronic bacterial prostatitis, prostatodystonia, prostatosis, granulomatous prostatitis, malacoplakia, benign prostatic hypertrophy or hyperplasia, and prostate neoplastic disorders, including adenocarcinomas, transitional cell carcinomas, ductal carcinomas, and squamous cell carcinomas.

[774] Additionally, the compositions of the invention may be useful in the diagnosis, treatment, and/or prevention of disorders or diseases of the penis and urethra, including inflammatory disorders, such as balanoposthitis, balanitis xerotica obliterans, phimosis, paraphimosis, syphilis, herpes simplex virus, gonorrhea, non-gonococcal urethritis, chlamydia, mycoplasma, trichomonas, HIV, AIDS, Reiter's syndrome, condyloma acuminatum, condyloma latum, and pearly penile papules; urethral abnormalities, such as hypospadias, epispadias, and phimosis; premalignant lesions, including Erythroplasia of Queyrat, Bowen's disease, Bowenoid paplosis, giant condyloma of Buscke-Lowenstein, and verrucous carcinoma; penile cancers, including squamous cell carcinomas, carcinoma in situ, verrucous carcinoma, and disseminated penile carcinoma; urethral neoplastic disorders, including penile urethral carcinoma, bulbomembranous urethral carcinoma, and prostatic urethral carcinoma; and erectile disorders, such as priapism, Peyronie's disease, erectile dysfunction, and impotence.

[775] Moreover, diseases and/or disorders of the vas deferens include vasculitis and CBAVD (congenital bilateral absence of the vas deferens); additionally, the polynucleotides, polypeptides, and agonists or antagonists of the present invention may be used in the diagnosis, treatment, and/or prevention of diseases and/or disorders of the seminal vesicles, including hydatid disease, congenital chloride diarrhea, and polycystic kidney disease.

[776] Other disorders and/or diseases of the male reproductive system include, for example, Klinefelter's syndrome, Young's syndrome, premature ejaculation, diabetes mellitus, cystic fibrosis, Kartagener's syndrome, high fever, multiple sclerosis, and gynecomastia.

[777] Further, the polynucleotides, polypeptides, and agonists or antagonists of the present invention may be used in the diagnosis, treatment, and/or prevention of diseases and/or disorders of the vagina and vulva, including bacterial vaginosis, candida vaginitis, herpes simplex virus, chancroid, granuloma inguinale, lymphogranuloma venereum,

scabies, human papillomavirus, vaginal trauma, vulvar trauma, adenositis, chlamydia vaginitis, gonorrhea, trichomonas vaginitis, condyloma acuminatum, syphilis, molluscum contagiosum, atrophic vaginitis, Paget's disease, lichen sclerosus, lichen planus, vulvodynia, toxic shock syndrome, vaginismus, vulvovaginitis, vulvar vestibulitis, and neoplastic disorders, such as squamous cell hyperplasia, clear cell carcinoma, basal cell carcinoma, melanomas, cancer of Bartholin's gland, and vulvar intraepithelial neoplasia.

[778] Disorders and/or diseases of the uterus include dysmenorrhea, retroverted uterus, endometriosis, fibroids, adenomyosis, anovulatory bleeding, amenorrhea, Cushing's syndrome, hydatidiform moles, Asherman's syndrome, premature menopause, precocious puberty, uterine polyps, dysfunctional uterine bleeding (e.g., due to aberrant hormonal signals), and neoplastic disorders, such as adenocarcinomas, leiomyosarcomas, and sarcomas. Additionally, the polypeptides, polynucleotides, or agonists or antagonists of the invention may be useful as a marker or detector of, as well as in the diagnosis, treatment, and/or prevention of congenital uterine abnormalities, such as bicornuate uterus, septate uterus, simple unicornuate uterus, unicornuate uterus with a noncavitary rudimentary horn, unicornuate uterus with a non-communicating cavitary rudimentary horn, unicornuate uterus with a communicating cavitary horn, arcuate uterus, uterine didelphys, and T-shaped uterus.

[779] Ovarian diseases and/or disorders include anovulation, polycystic ovary syndrome (Stein-Leventhal syndrome), ovarian cysts, ovarian hypofunction, ovarian insensitivity to gonadotropins, ovarian overproduction of androgens, right ovarian vein syndrome, amenorrhea, hirsutism, and ovarian cancer (including, but not limited to, primary and secondary cancerous growth, Sertoli-Leydig tumors, endometrioid carcinoma of the ovary, ovarian papillary serous adenocarcinoma, ovarian mucinous adenocarcinoma, and Ovarian Krukenberg tumors).

[780] Cervical diseases and/or disorders include cervicitis, chronic cervicitis, mucopurulent cervicitis, cervical dysplasia, cervical polyps, Nabothian cysts, cervical erosion, cervical incompetence, and cervical neoplasms (including, for example, cervical carcinoma, squamous metaplasia, squamous cell carcinoma, adenosquamous cell neoplasia, and columnar cell neoplasia).

[781] Additionally, diseases and/or disorders of the reproductive system include disorders and/or diseases of pregnancy, including miscarriage and stillbirth, such as early abortion, late abortion, spontaneous abortion, induced abortion, therapeutic abortion,

threatened abortion, missed abortion, incomplete abortion, complete abortion, habitual abortion, missed abortion, and septic abortion; ectopic pregnancy, anemia, Rh incompatibility, vaginal bleeding during pregnancy, gestational diabetes, intrauterine growth retardation, polyhydramnios, HELLP syndrome, abruptio placentae, placenta previa, hyperemesis, preeclampsia, eclampsia, herpes gestationis, and urticaria of pregnancy. Additionally, the polynucleotides, polypeptides, and agonists or antagonists of the present invention may be used in the diagnosis, treatment, and/or prevention of diseases that can complicate pregnancy, including heart disease, heart failure, rheumatic heart disease, congenital heart disease, mitral valve prolapse, high blood pressure, anemia, kidney disease, infectious disease (e.g., rubella, cytomegalovirus, toxoplasmosis, infectious hepatitis, chlamydia, HIV, AIDS, and genital herpes), diabetes mellitus, Graves' disease, thyroiditis, hypothyroidism, Hashimoto's thyroiditis, chronic active hepatitis, cirrhosis of the liver, primary biliary cirrhosis, asthma, systemic lupus eryematosus, rheumatoid arthritis, myasthenia gravis, idiopathic thrombocytopenic purpura, appendicitis, ovarian cysts, gallbladder disorders, and obstruction of the intestine.

[782] Complications associated with labor and parturition include premature rupture of the membranes, pre-term labor, post-term pregnancy, postmaturity, labor that progresses too slowly, fetal distress (e.g., abnormal heart rate (fetal or maternal), breathing problems, and abnormal fetal position), shoulder dystocia, prolapsed umbilical cord, amniotic fluid embolism, and aberrant uterine bleeding.

[783] Further, diseases and/or disorders of the postdelivery period, including endometritis, myometritis, parametritis, peritonitis, pelvic thrombophlebitis, pulmonary embolism, endotoxemia, pyelonephritis, saphenous thrombophlebitis, mastitis, cystitis, postpartum hemorrhage, and inverted uterus.

[784] Other disorders and/or diseases of the female reproductive system that may be diagnosed, treated, and/or prevented by the polynucleotides, polypeptides, and agonists or antagonists of the present invention include, for example, Turner's syndrome, pseudohermaphroditism, premenstrual syndrome, pelvic inflammatory disease, pelvic congestion (vascular engorgement), frigidity, anorgasmia, dyspareunia, ruptured fallopian tube, and Mittelschmerz.

Infectious Disease

[785] Polynucleotides or polypeptides, as well as agonists or antagonists of the present invention can be used to treat or detect infectious agents. For example, by increasing the immune response, particularly increasing the proliferation and differentiation of B and/or T cells, infectious diseases may be treated. The immune response may be increased by either enhancing an existing immune response, or by initiating a new immune response. Alternatively, polynucleotides or polypeptides, as well as agonists or antagonists of the present invention may also directly inhibit the infectious agent, without necessarily eliciting an immune response.

[786] Viruses are one example of an infectious agent that can cause disease or symptoms that can be treated or detected by a polynucleotide or polypeptide and/or agonist or antagonist of the present invention. Examples of viruses, include, but are not limited to Examples of viruses, include, but are not limited to the following DNA and RNA viruses and viral families: Arbovirus, Adenoviridae, Arenaviridae, Arterivirus, Birnaviridae, Bunyaviridae, Caliciviridae, Circoviridae, Coronaviridae, Dengue, EBV, HIV, Flaviviridae, Hepadnaviridae (Hepatitis), Herpesviridae (such as, Cytomegalovirus, Herpes Simplex, Herpes Zoster), Mononegavirus (e.g., Paramyxoviridae, Morbillivirus, Rhabdoviridae), Orthomyxoviridae (e.g., Influenza A, Influenza B, and parainfluenza), Papiloma virus, Papovaviridae, Parvoviridae, Picornaviridae, Poxviridae (such as Smallpox or Vaccinia), Reoviridae (e.g., Rotavirus), Retroviridae (HTLV-I, HTLV-II, Lentivirus), and Togaviridae (e.g., Rubivirus). Viruses falling within these families can cause a variety of diseases or symptoms, including, but not limited to: arthritis, bronchiollitis, respiratory syncytial virus, encephalitis, eye infections (e.g., conjunctivitis, keratitis), chronic fatigue syndrome, hepatitis (A, B, C, E, Chronic Active, Delta), Japanese B encephalitis, Junin, Chikungunya, Rift Valley fever, yellow fever, meningitis, opportunistic infections (e.g., AIDS), pneumonia, Burkitt's Lymphoma, chickenpox, hemorrhagic fever, Measles, Mumps, Parainfluenza, Rabies, the common cold, Polio, leukemia, Rubella, sexually transmitted diseases, skin diseases (e.g., Kaposi's, warts), and viremia. polynucleotides or polypeptides, or agonists or antagonists of the invention, can be used to treat or detect any of these symptoms or diseases. In specific embodiments, polynucleotides, polypeptides, or agonists or antagonists of the invention are used to treat: meningitis, Dengue, EBV, and/or hepatitis (e.g., hepatitis B). In an additional specific embodiment polynucleotides, polypeptides, or agonists or antagonists of the invention are used to treat patients nonresponsive to one or more other commercially available hepatitis vaccines. In a further specific embodiment

polynucleotides, polypeptides, or agonists or antagonists of the invention are used to treat AIDS.

[787] Similarly, bacterial and fungal agents that can cause disease or symptoms and that can be treated or detected by a polynucleotide or polypeptide and/or agonist or antagonist of the present invention include, but not limited to, the following Gram-Negative and Gram-positive bacteria, bacterial families, and fungi: Actinomyces (e.g., Norcardia), Acinetobacter, *Cryptococcus neoformans*, Aspergillus, Bacillaceae (e.g., *Bacillus anthraxis*), Bacteroides (e.g., *Bacteroides fragilis*), Blastomycosis, Bordetella, Borrelia (e.g., *Borrelia burgdorferi*), Brucella, Candidia, Campylobacter, Chlamydia, Clostridium (e.g., *Clostridium botulinum*, *Clostridium difficile*, *Clostridium perfringens*, *Clostridium tetani*), Coccidioides, Corynebacterium (e.g., *Corynebacterium diphtheriae*), Cryptococcus, Dermatocycoses, *E. coli* (e.g., Enterotoxigenic *E. coli* and Enterohemorrhagic *E. coli*), Enterobacter (e.g. *Enterobacter aerogenes*), Enterobacteriaceae (Klebsiella, Salmonella (e.g., *Salmonella typhi*, *Salmonella enteritidis*, *Salmonella typhi*), Serratia, Yersinia, Shigella), Erysipelothrix, Haemophilus (e.g., *Haemophilus influenza* type B), Helicobacter, Legionella (e.g., *Legionella pneumophila*), Leptospira, Listeria (e.g., *Listeria monocytogenes*), Mycoplasma, Mycobacterium (e.g., *Mycobacterium leprae* and *Mycobacterium tuberculosis*), Vibrio (e.g., *Vibrio cholerae*), Neisseriaceae (e.g., *Neisseria gonorrhea*, *Neisseria meningitidis*), Pasteurellacea, Proteus, Pseudomonas (e.g., *Pseudomonas aeruginosa*), Rickettsiaceae, Spirochetes (e.g., Treponema spp., Leptospira spp., Borrelia spp.), Shigella spp., Staphylococcus (e.g., *Staphylococcus aureus*), Meningiococcus, Pneumococcus and Streptococcus (e.g., *Streptococcus pneumoniae* and Groups A, B, and C Streptococci), and Ureaplasmas. These bacterial, parasitic, and fungal families can cause diseases or symptoms, including, but not limited to: antibiotic-resistant infections, bacteremia, endocarditis, septicemia, eye infections (e.g., conjunctivitis), uveitis, tuberculosis, gingivitis, bacterial diarrhea, opportunistic infections (e.g., AIDS related infections), paronychia, prosthesis-related infections, dental caries, Reiter's Disease, respiratory tract infections, such as Whooping Cough or Empyema, sepsis, Lyme Disease, Cat-Scratch Disease, dysentery, paratyphoid fever, food poisoning, Legionella disease, chronic and acute inflammation, erythema, yeast infections, typhoid, pneumonia, gonorrhea, meningitis (e.g., meningitis types A and B), chlamydia, syphilis, diphtheria, leprosy, brucellosis, peptic ulcers, anthrax, spontaneous abortions, birth defects, pneumonia, lung infections, ear infections, deafness, blindness, lethargy, malaise, vomiting, chronic

diarrhea, Crohn's disease, colitis, vaginosis, sterility, pelvic inflammatory diseases, candidiasis, paratuberculosis, tuberculosis, lupus, botulism, gangrene, tetanus, impetigo, Rheumatic Fever, Scarlet Fever, sexually transmitted diseases, skin diseases (e.g., cellulitis, dermatocycoses), toxemia, urinary tract infections, wound infections, noscomial infections. Polynucleotides or polypeptides, agonists or antagonists of the invention, can be used to treat or detect any of these symptoms or diseases. In specific embodiments, polynucleotides, polypeptides, agonists or antagonists of the invention are used to treat: tetanus, diphtheria, botulism, and/or meningitis type B.

[788] Moreover, parasitic agents causing disease or symptoms that can be treated, prevented, and/or diagnosed by a polynucleotide or polypeptide and/or agonist or antagonist of the present invention include, but not limited to, the following families or class: Amebiasis, Babesiosis, Coccidiosis, Cryptosporidiosis, Dientamoebiasis, Dourine, Ectoparasitic, Giardiasis, Helminthiasis, Leishmaniasis, Schistosoma, Theileriasis, Toxoplasmosis, Trypanosomiasis, and Trichomonas and Sporozoans (e.g., *Plasmodium virax*, *Plasmodium falciparum*, *Plasmodium malariae* and *Plasmodium ovale*). These parasites can cause a variety of diseases or symptoms, including, but not limited to: Scabies, Trombiculiasis, eye infections, intestinal disease (e.g., dysentery, giardiasis), liver disease, lung disease, opportunistic infections (e.g., AIDS related), malaria, pregnancy complications, and toxoplasmosis. polynucleotides or polypeptides, or agonists or antagonists of the invention, can be used to treat, prevent, and/or diagnose any of these symptoms or diseases. In specific embodiments, polynucleotides, polypeptides, or agonists or antagonists of the invention are used to treat, prevent, and/or diagnose malaria.

[789] Polynucleotides or polypeptides, as well as agonists or antagonists of the present invention of the present invention could either be by administering an effective amount of a polypeptide to the patient, or by removing cells from the patient, supplying the cells with a polynucleotide of the present invention, and returning the engineered cells to the patient (ex vivo therapy). Moreover, the polypeptide or polynucleotide of the present invention can be used as an antigen in a vaccine to raise an immune response against infectious disease.

Regeneration

[790] Polynucleotides or polypeptides, as well as agonists or antagonists of the present invention can be used to differentiate, proliferate, and attract cells, leading to the regeneration of tissues. (See, Science 276:59-87 (1997)). The regeneration of tissues could

be used to repair, replace, or protect tissue damaged by congenital defects, trauma (wounds, burns, incisions, or ulcers), age, disease (e.g. osteoporosis, osteoarthritis, periodontal disease, liver failure), surgery, including cosmetic plastic surgery, fibrosis, reperfusion injury, or systemic cytokine damage.

[791] Tissues that could be regenerated using the present invention include organs (e.g., pancreas, liver, intestine, kidney, skin, endothelium), muscle (smooth, skeletal or cardiac), vasculature (including vascular and lymphatics), nervous, hematopoietic, and skeletal (bone, cartilage, tendon, and ligament) tissue. Preferably, regeneration occurs without or decreased scarring. Regeneration also may include angiogenesis.

[792] Moreover, polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, may increase regeneration of tissues difficult to heal. For example, increased tendon/ligament regeneration would quicken recovery time after damage. Polynucleotides or polypeptides, as well as agonists or antagonists of the present invention could also be used prophylactically in an effort to avoid damage. Specific diseases that could be treated include of tendinitis, carpal tunnel syndrome, and other tendon or ligament defects. A further example of tissue regeneration of non-healing wounds includes pressure ulcers, ulcers associated with vascular insufficiency, surgical, and traumatic wounds.

[793] Similarly, nerve and brain tissue could also be regenerated by using polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, to proliferate and differentiate nerve cells. Diseases that could be treated using this method include central and peripheral nervous system diseases, neuropathies, or mechanical and traumatic disorders (e.g., spinal cord disorders, head trauma, cerebrovascular disease, and stroke). Specifically, diseases associated with peripheral nerve injuries, peripheral neuropathy (e.g., resulting from chemotherapy or other medical therapies), localized neuropathies, and central nervous system diseases (e.g., Alzheimer's disease, Parkinson's disease, Huntington's disease, amyotrophic lateral sclerosis, and Shy-Drager syndrome), could all be treated using the polynucleotides or polypeptides, as well as agonists or antagonists of the present invention.

Gastrointestinal Disorders

[794] Polynucleotides or polypeptides, or agonists or antagonists of the present invention, may be used to treat, prevent, diagnose, and/or prognose gastrointestinal

disorders, including inflammatory diseases and/or conditions, infections, cancers (e.g., intestinal neoplasms (carcinoid tumor of the small intestine, non-Hodgkin's lymphoma of the small intestine, small bowel lymphoma)), and ulcers, such as peptic ulcers.

[795] Gastrointestinal disorders include dysphagia, odynophagia, inflammation of the esophagus, peptic esophagitis, gastric reflux, submucosal fibrosis and stricturing, Mallory-Weiss lesions, leiomyomas, lipomas, epidermal cancers, adeoncarcinomas, gastric retention disorders, gastroenteritis, gastric atrophy, gastric/stomach cancers, polyps of the stomach, autoimmune disorders such as pernicious anemia, pyloric stenosis, gastritis (bacterial, viral, eosinophilic, stress-induced, chronic erosive, atrophic, plasma cell, and Ménétrier's), and peritoneal diseases (e.g., chyloperitoneum, hemoperitoneum, mesenteric cyst, mesenteric lymphadenitis, mesenteric vascular occlusion, panniculitis, neoplasms, peritonitis, pneumoperitoneum, bubphrenic abscess,).

[796] Gastrointestinal disorders also include disorders associated with the small intestine, such as malabsorption syndromes, distension, irritable bowel syndrome, sugar intolerance, celiac disease, duodenal ulcers, duodenitis, tropical sprue, Whipple's disease, intestinal lymphangiectasia, Crohn's disease, appendicitis, obstructions of the ileum, Meckel's diverticulum, multiple diverticula, failure of complete rotation of the small and large intestine, lymphoma, and bacterial and parasitic diseases (such as Traveler's diarrhea, typhoid and paratyphoid, cholera, infection by Roundworms (*Ascariasis lumbricoides*), Hookworms (*Ancylostoma duodenale*), Threadworms (*Enterobius vermicularis*), Tapeworms (*Taenia saginata*, *Echinococcus granulosus*, *Diphyllobothrium spp.*, and *T. solium*).

[797] Liver diseases and/or disorders include intrahepatic cholestasis (alagille syndrome, biliary liver cirrhosis), fatty liver (alcoholic fatty liver, reye syndrome), hepatic vein thrombosis, hepatolentricular degeneration, hepatomegaly, hepatopulmonary syndrome, hepatorenal syndrome, portal hypertension (esophageal and gastric varices), liver abscess (amebic liver abscess), liver cirrhosis (alcoholic, biliary and experimental), alcoholic liver diseases (fatty liver, hepatitis, cirrhosis), parasitic (hepatic echinococcosis, fascioliasis, amebic liver abscess), jaundice (hemolytic, hepatocellular, and cholestatic), cholestasis, portal hypertension, liver enlargement, ascites, hepatitis (alcoholic hepatitis, animal hepatitis, chronic hepatitis (autoimmune, hepatitis B, hepatitis C, hepatitis D, drug induced), toxic hepatitis, viral human hepatitis (hepatitis A, hepatitis B, hepatitis C, hepatitis D, hepatitis E), Wilson's disease, granulomatous hepatitis, secondary biliary

cirrhosis, hepatic encephalopathy, portal hypertension, varices, hepatic encephalopathy, primary biliary cirrhosis, primary sclerosing cholangitis, hepatocellular adenoma, hemangiomas, bile stones, liver failure (hepatic encephalopathy, acute liver failure), and liver neoplasms (angiomyolipoma, calcified liver metastases, cystic liver metastases, epithelial tumors, fibrolamellar hepatocarcinoma, focal nodular hyperplasia, hepatic adenoma, hepatobiliary cystadenoma, hepatoblastoma, hepatocellular carcinoma, hepatoma, liver cancer, liver hemangioendothelioma, mesenchymal hamartoma, mesenchymal tumors of liver, nodular regenerative hyperplasia, benign liver tumors (Hepatic cysts [Simple cysts, Polycystic liver disease, Hepatobiliary cystadenoma, Choledochal cyst], Mesenchymal tumors [Mesenchymal hamartoma, Infantile hemangioendothelioma, Hemangioma, Peliosis hepatis, Lipomas, Inflammatory pseudotumor, Miscellaneous], Epithelial tumors [Bile duct epithelium (Bile duct hamartoma, Bile duct adenoma), Hepatocyte (Adenoma, Focal nodular hyperplasia, Nodular regenerative hyperplasia)]), malignant liver tumors [hepatocellular, hepatoblastoma, hepatocellular carcinoma, cholangiocellular, cholangiocarcinoma, cystadenocarcinoma, tumors of blood vessels, angiosarcoma, Kaposi's sarcoma, hemangioendothelioma, other tumors, embryonal sarcoma, fibrosarcoma, leiomyosarcoma, rhabdomyosarcoma, carcinosarcoma, teratoma, carcinoid, squamous carcinoma, primary lymphoma]), peliosis hepatis, erythrohepatic porphyria, hepatic porphyria (acute intermittent porphyria, porphyria cutanea tarda), Zellweger syndrome).

[798] Pancreatic diseases and/or disorders include acute pancreatitis, chronic pancreatitis (acute necrotizing pancreatitis, alcoholic pancreatitis), neoplasms (adenocarcinoma of the pancreas, cystadenocarcinoma, insulinoma, gastrinoma, and glucagonoma, cystic neoplasms, islet-cell tumors, pancreoblastoma), and other pancreatic diseases (e.g., cystic fibrosis, cyst (pancreatic pseudocyst, pancreatic fistula, insufficiency)).

[799] Gallbladder diseases include gallstones (cholelithiasis and choledocholithiasis), postcholecystectomy syndrome, diverticulosis of the gallbladder, acute cholecystitis, chronic cholecystitis, bile duct tumors, and mucocele.

[800] Diseases and/or disorders of the large intestine include antibiotic-associated colitis, diverticulitis, ulcerative colitis, acquired megacolon, abscesses, fungal and bacterial infections, anorectal disorders (e.g., fissures, hemorrhoids), colonic diseases (colitis, colonic neoplasms [colon cancer, adenomatous colon polyps (e.g., villous adenoma), colon carcinoma, colorectal cancer], colonic diverticulitis, colonic diverticulosis, megacolon

[Hirschsprung disease, toxic megacolon]; sigmoid diseases [proctocolitis, sigmoid neoplasms]), constipation, Crohn's disease, diarrhea (infantile diarrhea, dysentery), duodenal diseases (duodenal neoplasms, duodenal obstruction, duodenal ulcer, duodenitis), enteritis (enterocolitis), HIV enteropathy, ileal diseases (ileal neoplasms, ileitis), immunoproliferative small intestinal disease, inflammatory bowel disease (ulcerative colitis, Crohn's disease), intestinal atresia, parasitic diseases (anisakiasis, balantidiasis, blastocystis infections, cryptosporidiosis, dientamoebiasis, amebic dysentery, giardiasis), intestinal fistula (rectal fistula), intestinal neoplasms (cecal neoplasms, colonic neoplasms, duodenal neoplasms, ileal neoplasms, intestinal polyps, jejunal neoplasms, rectal neoplasms), intestinal obstruction (afferent loop syndrome, duodenal obstruction, impacted feces, intestinal pseudo-obstruction [cecal volvulus], intussusception), intestinal perforation, intestinal polyps (colonic polyps, gardner syndrome, peutz-jeghers syndrome), jejunal diseases (jejunal neoplasms), malabsorption syndromes (blind loop syndrome, celiac disease, lactose intolerance, short bowel syndrome, tropical sprue, whipple's disease), mesenteric vascular occlusion, pneumatosis cystoides intestinalis, protein-losing enteropathies (intestinal lymphagiectasis), rectal diseases (anus diseases, fecal incontinence, hemorrhoids, proctitis, rectal fistula, rectal prolapse, rectocele), peptic ulcer (duodenal ulcer, peptic esophagitis, hemorrhage, perforation, stomach ulcer, Zollinger-Ellison syndrome), postgastrectomy syndromes (dumping syndrome), stomach diseases (e.g., achlorhydria, duodenogastric reflux (bile reflux), gastric antral vascular ectasia, gastric fistula, gastric outlet obstruction, gastritis (atrophic or hypertrophic), gastroparesis, stomach dilatation, stomach diverticulum, stomach neoplasms (gastric cancer, gastric polyps, gastric adenocarcinoma, hyperplastic gastric polyp), stomach rupture, stomach ulcer, stomach volvulus), tuberculosis, visceroptosis, vomiting (e.g., hematemesis, hyperemesis gravidarum, postoperative nausea and vomiting) and hemorrhagic colitis.

[801] Further diseases and/or disorders of the gastrointestinal system include biliary tract diseases, such as, gastroschisis, fistula (e.g., biliary fistula, esophageal fistula, gastric fistula, intestinal fistula, pancreatic fistula), neoplasms (e.g., biliary tract neoplasms, esophageal neoplasms, such as adenocarcinoma of the esophagus, esophageal squamous cell carcinoma, gastrointestinal neoplasms, pancreatic neoplasms, such as adenocarcinoma of the pancreas, mucinous cystic neoplasm of the pancreas, pancreatic cystic neoplasms, pancreatoblastoma, and peritoneal neoplasms), esophageal disease (e.g., bullous diseases, candidiasis, glycogenic acanthosis, ulceration, barrett esophagus varices, atresia, cyst,

diverticulum (e.g., Zenker's diverticulum), fistula (e.g., tracheoesophageal fistula), motility disorders (e.g., CREST syndrome, deglutition disorders, achalasia, spasm, gastroesophageal reflux), neoplasms, perforation (e.g., Boerhaave syndrome, Mallory-Weiss syndrome), stenosis, esophagitis, diaphragmatic hernia (e.g., hiatal hernia); gastrointestinal diseases, such as, gastroenteritis (e.g., cholera morbus, norwalk virus infection), hemorrhage (e.g., hematemesis, melena, peptic ulcer hemorrhage), stomach neoplasms (gastric cancer, gastric polyps, gastric adenocarcinoma, stomach cancer)), hernia (e.g., congenital diaphragmatic hernia, femoral hernia, inguinal hernia, obturator hernia, umbilical hernia, ventral hernia), and intestinal diseases (e.g., cecal diseases (appendicitis, cecal neoplasms)).

Chemotaxis

[802] Polynucleotides or polypeptides, as well as agonists or antagonists of the present invention may have chemotaxis activity. A chemotactic molecule attracts or mobilizes cells (e.g., monocytes, fibroblasts, neutrophils, T-cells, mast cells, eosinophils, epithelial and/or endothelial cells) to a particular site in the body, such as inflammation, infection, or site of hyperproliferation. The mobilized cells can then fight off and/or heal the particular trauma or abnormality.

[803] Polynucleotides or polypeptides, as well as agonists or antagonists of the present invention may increase chemotactic activity of particular cells. These chemotactic molecules can then be used to treat inflammation, infection, hyperproliferative disorders, or any immune system disorder by increasing the number of cells targeted to a particular location in the body. For example, chemotactic molecules can be used to treat wounds and other trauma to tissues by attracting immune cells to the injured location. Chemotactic molecules of the present invention can also attract fibroblasts, which can be used to treat wounds.

[804] It is also contemplated that polynucleotides or polypeptides, as well as agonists or antagonists of the present invention may inhibit chemotactic activity. These molecules could also be used to treat disorders. Thus, polynucleotides or polypeptides, as well as agonists or antagonists of the present invention could be used as an inhibitor of chemotaxis.

Binding Activity

[805] A polypeptide of the present invention may be used to screen for molecules that bind to the polypeptide or for molecules to which the polypeptide binds. The binding of the polypeptide and the molecule may activate (agonist), increase, inhibit (antagonist), or decrease activity of the polypeptide or the molecule bound. Examples of such molecules include antibodies, oligonucleotides, proteins (e.g., receptors), or small molecules.

[806] Preferably, the molecule is closely related to the natural ligand of the polypeptide, e.g., a fragment of the ligand, or a natural substrate, a ligand, a structural or functional mimetic. (See, Coligan et al., Current Protocols in Immunology 1(2):Chapter 5 (1991)). Similarly, the molecule can be closely related to the natural receptor to which the polypeptide binds, or at least, a fragment of the receptor capable of being bound by the polypeptide (e.g., active site). In either case, the molecule can be rationally designed using known techniques.

[807] Preferably, the screening for these molecules involves producing appropriate cells which express the polypeptide. Preferred cells include cells from mammals, yeast, *Drosophila*, or *E. coli*. Cells expressing the polypeptide (or cell membrane containing the expressed polypeptide) are then preferably contacted with a test compound potentially containing the molecule to observe binding, stimulation, or inhibition of activity of either the polypeptide or the molecule.

[808] The assay may simply test binding of a candidate compound to the polypeptide, wherein binding is detected by a label, or in an assay involving competition with a labeled competitor. Further, the assay may test whether the candidate compound results in a signal generated by binding to the polypeptide.

[809] Alternatively, the assay can be carried out using cell-free preparations, polypeptide/molecule affixed to a solid support, chemical libraries, or natural product mixtures. The assay may also simply comprise the steps of mixing a candidate compound with a solution containing a polypeptide, measuring polypeptide/molecule activity or binding, and comparing the polypeptide/molecule activity or binding to a standard.

[810] Preferably, an ELISA assay can measure polypeptide level or activity in a sample (e.g., biological sample) using a monoclonal or polyclonal antibody. The antibody can measure polypeptide level or activity by either binding, directly or indirectly, to the polypeptide or by competing with the polypeptide for a substrate.

[811] Additionally, the receptor to which the polypeptide of the present invention binds can be identified by numerous methods known to those of skill in the art, for example,

ligand panning and FACS sorting (Coligan, et al., Current Protocols in Immun., 1(2), Chapter 5, (1991)). For example, expression cloning is employed wherein polyadenylated RNA is prepared from a cell responsive to the polypeptides, for example, NIH3T3 cells which are known to contain multiple receptors for the FGF family proteins, and SC-3 cells, and a cDNA library created from this RNA is divided into pools and used to transfect COS cells or other cells that are not responsive to the polypeptides. Transfected cells which are grown on glass slides are exposed to the polypeptide of the present invention, after they have been labeled. The polypeptides can be labeled by a variety of means including iodination or inclusion of a recognition site for a site-specific protein kinase.

[812] Following fixation and incubation, the slides are subjected to auto-radiographic analysis. Positive pools are identified and sub-pools are prepared and re-transfected using an iterative sub-pooling and re-screening process, eventually yielding a single clones that encodes the putative receptor.

[813] As an alternative approach for receptor identification, the labeled polypeptides can be photoaffinity linked with cell membrane or extract preparations that express the receptor molecule. Cross-linked material is resolved by PAGE analysis and exposed to X-ray film. The labeled complex containing the receptors of the polypeptides can be excised, resolved into peptide fragments, and subjected to protein microsequencing. The amino acid sequence obtained from microsequencing would be used to design a set of degenerate oligonucleotide probes to screen a cDNA library to identify the genes encoding the putative receptors.

[814] Moreover, the techniques of gene-shuffling, motif-shuffling, exon-shuffling, and/or codon-shuffling (collectively referred to as "DNA shuffling") may be employed to modulate the activities of the polypeptide of the present invention thereby effectively generating agonists and antagonists of the polypeptide of the present invention. *See generally*, U.S. Patent Nos. 5,605,793, 5,811,238, 5,830,721, 5,834,252, and 5,837,458, and Patten, P. A., *et al.*, *Curr. Opinion Biotechnol.* 8:724-33 (1997); Harayama, S. *Trends Biotechnol.* 16(2):76-82 (1998); Hansson, L. O., *et al.*, *J. Mol. Biol.* 287:265-76 (1999); and Lorenzo, M. M. and Blasco, R. *Biotechniques* 24(2):308-13 (1998); each of these patents and publications are hereby incorporated by reference). In one embodiment, alteration of polynucleotides and corresponding polypeptides may be achieved by DNA shuffling. DNA shuffling involves the assembly of two or more DNA segments into a desired molecule by homologous, or site-specific, recombination. In another embodiment,

polynucleotides and corresponding polypeptides may be altered by being subjected to random mutagenesis by error-prone PCR, random nucleotide insertion or other methods prior to recombination. In another embodiment, one or more components, motifs, sections, parts, domains, fragments, etc., of the polypeptide of the present invention may be recombined with one or more components, motifs, sections, parts, domains, fragments, etc. of one or more heterologous molecules. In preferred embodiments, the heterologous molecules are family members. In further preferred embodiments, the heterologous molecule is a growth factor such as, for example, platelet-derived growth factor (PDGF), insulin-like growth factor (IGF-I), transforming growth factor (TGF)-alpha, epidermal growth factor (EGF), fibroblast growth factor (FGF), TGF-beta, bone morphogenetic protein (BMP)-2, BMP-4, BMP-5, BMP-6, BMP-7, activins A and B, decapentaplegic(dpp), 60A, OP-2, dorsalin, growth differentiation factors (GDFs), nodal, MIS, inhibin-alpha, TGF-beta1, TGF-beta2, TGF-beta3, TGF-beta5, and glial-derived neurotrophic factor (GDNF).

[815] Other preferred fragments are biologically active fragments of the polypeptide of the present invention. Biologically active fragments are those exhibiting activity similar, but not necessarily identical, to an activity of the polypeptide of the present invention. The biological activity of the fragments may include an improved desired activity, or a decreased undesirable activity.

[816] Additionally, this invention provides a method of screening compounds to identify those which modulate the action of the polypeptide of the present invention. An example of such an assay comprises combining a mammalian fibroblast cell, a the polypeptide of the present invention, the compound to be screened and $^3\text{[H]}$ thymidine under cell culture conditions where the fibroblast cell would normally proliferate. A control assay may be performed in the absence of the compound to be screened and compared to the amount of fibroblast proliferation in the presence of the compound to determine if the compound stimulates proliferation by determining the uptake of $^3\text{[H]}$ thymidine in each case. The amount of fibroblast cell proliferation is measured by liquid scintillation chromatography which measures the incorporation of $^3\text{[H]}$ thymidine. Both agonist and antagonist compounds may be identified by this procedure.

[817] In another method, a mammalian cell or membrane preparation expressing a receptor for a polypeptide of the present invention is incubated with a labeled polypeptide

of the present invention in the presence of the compound. The ability of the compound to enhance or block this interaction could then be measured. Alternatively, the response of a known second messenger system following interaction of a compound to be screened and the receptor is measured and the ability of the compound to bind to the receptor and elicit a second messenger response is measured to determine if the compound is a potential agonist or antagonist. Such second messenger systems include but are not limited to, cAMP guanylate cyclase, ion channels or phosphoinositide hydrolysis.

[818] All of these above assays can be used as diagnostic or prognostic markers. The molecules discovered using these assays can be used to treat disease or to bring about a particular result in a patient (e.g., blood vessel growth) by activating or inhibiting the polypeptide/molecule. Moreover, the assays can discover agents which may inhibit or enhance the production of the polypeptides of the invention from suitably manipulated cells or tissues.

[819] Therefore, the invention includes a method of identifying compounds which bind to a polypeptide of the invention comprising the steps of: (a) incubating a candidate binding compound with a polypeptide of the present invention; and (b) determining if binding has occurred. Moreover, the invention includes a method of identifying agonists/antagonists comprising the steps of: (a) incubating a candidate compound with a polypeptide of the present invention, (b) assaying a biological activity, and (b) determining if a biological activity of the polypeptide has been altered.

Targeted Delivery

[820] In another embodiment, the invention provides a method of delivering compositions to targeted cells expressing a receptor for a polypeptide of the invention, or cells expressing a cell bound form of a polypeptide of the invention.

[821] As discussed herein, polypeptides or antibodies of the invention may be associated with heterologous polypeptides, heterologous nucleic acids, toxins, or prodrugs via hydrophobic, hydrophilic, ionic and/or covalent interactions. In one embodiment, the invention provides a method for the specific delivery of compositions of the invention to cells by administering polypeptides of the invention (including antibodies) that are associated with heterologous polypeptides or nucleic acids. In one example, the invention provides a method for delivering a therapeutic protein into the targeted cell. In another example, the invention provides a method for delivering a single stranded nucleic acid (e.g.,

antisense or ribozymes) or double stranded nucleic acid (e.g., DNA that can integrate into the cell's genome or replicate episomally and that can be transcribed) into the targeted cell.

[822] In another embodiment, the invention provides a method for the specific destruction of cells (e.g., the destruction of tumor cells) by administering polypeptides of the invention (e.g., polypeptides of the invention or antibodies of the invention) in association with toxins or cytotoxic prodrugs.

[823] By "toxin" is meant compounds that bind and activate endogenous cytotoxic effector systems, radioisotopes, holotoxins, modified toxins, catalytic subunits of toxins, or any molecules or enzymes not normally present in or on the surface of a cell that under defined conditions cause the cell's death. Toxins that may be used according to the methods of the invention include, but are not limited to, radioisotopes known in the art, compounds such as, for example, antibodies (or complement fixing containing portions thereof) that bind an inherent or induced endogenous cytotoxic effector system, thymidine kinase, endonuclease, RNase, alpha toxin, ricin, abrin, *Pseudomonas* exotoxin A, diphtheria toxin, saporin, momordin, gelonin, pokeweed antiviral protein, alpha-sarcin and cholera toxin. By "cytotoxic prodrug" is meant a non-toxic compound that is converted by an enzyme, normally present in the cell, into a cytotoxic compound. Cytotoxic prodrugs that may be used according to the methods of the invention include, but are not limited to, glutamyl derivatives of benzoic acid mustard alkylating agent, phosphate derivatives of etoposide or mitomycin C, cytosine arabinoside, daunorubisin, and phenoxyacetamide derivatives of doxorubicin.

Drug Screening

[824] Further contemplated is the use of the polypeptides of the present invention, or the polynucleotides encoding these polypeptides, to screen for molecules which modify the activities of the polypeptides of the present invention. Such a method would include contacting the polypeptide of the present invention with a selected compound(s) suspected of having antagonist or agonist activity, and assaying the activity of these polypeptides following binding.

[825] This invention is particularly useful for screening therapeutic compounds by using the polypeptides of the present invention, or binding fragments thereof, in any of a variety of drug screening techniques. The polypeptide or fragment employed in such a test

may be affixed to a solid support, expressed on a cell surface, free in solution, or located intracellularly. One method of drug screening utilizes eukaryotic or prokaryotic host cells which are stably transformed with recombinant nucleic acids expressing the polypeptide or fragment. Drugs are screened against such transformed cells in competitive binding assays. One may measure, for example, the formulation of complexes between the agent being tested and a polypeptide of the present invention.

[826] Thus, the present invention provides methods of screening for drugs or any other agents which affect activities mediated by the polypeptides of the present invention. These methods comprise contacting such an agent with a polypeptide of the present invention or a fragment thereof and assaying for the presence of a complex between the agent and the polypeptide or a fragment thereof, by methods well known in the art. In such a competitive binding assay, the agents to screen are typically labeled. Following incubation, free agent is separated from that present in bound form, and the amount of free or uncomplexed label is a measure of the ability of a particular agent to bind to the polypeptides of the present invention.

[827] Another technique for drug screening provides high throughput screening for compounds having suitable binding affinity to the polypeptides of the present invention, and is described in great detail in European Patent Application 84/03564, published on September 13, 1984, which is incorporated herein by reference herein. Briefly stated, large numbers of different small peptide test compounds are synthesized on a solid substrate, such as plastic pins or some other surface. The peptide test compounds are reacted with polypeptides of the present invention and washed. Bound polypeptides are then detected by methods well known in the art. Purified polypeptides are coated directly onto plates for use in the aforementioned drug screening techniques. In addition, non-neutralizing antibodies may be used to capture the peptide and immobilize it on the solid support.

[828] This invention also contemplates the use of competitive drug screening assays in which neutralizing antibodies capable of binding polypeptides of the present invention specifically compete with a test compound for binding to the polypeptides or fragments thereof. In this manner, the antibodies are used to detect the presence of any peptide which shares one or more antigenic epitopes with a polypeptide of the invention.

Antisense And Ribozyme (Antagonists)

[829] In specific embodiments, antagonists according to the present invention are nucleic acids corresponding to the sequences contained in SEQ ID NO:X, or the complementary strand thereof, and/or to cDNA sequences contained in cDNA Clone ID NO:Z identified for example, in Table 1A and/or 1B. In one embodiment, antisense sequence is generated internally, by the organism, in another embodiment, the antisense sequence is separately administered (see, for example, O'Connor, J., *Neurochem.* 56:560 (1991). *Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression*, CRC Press, Boca Raton, FL (1988). Antisense technology can be used to control gene expression through antisense DNA or RNA, or through triple-helix formation. Antisense techniques are discussed for example, in Okano, J., *Neurochem.* 56:560 (1991); *Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression*, CRC Press, Boca Raton, FL (1988). Triple helix formation is discussed in, for instance, Lee et al., *Nucleic Acids Research* 6:3073 (1979); Cooney et al., *Science* 241:456 (1988); and Dervan et al., *Science* 251:1300 (1991). The methods are based on binding of a polynucleotide to a complementary DNA or RNA.

[830] For example, the use of c-myc and c-myb antisense RNA constructs to inhibit the growth of the non-lymphocytic leukemia cell line HL-60 and other cell lines was previously described. (Wickstrom et al. (1988); Anfossi et al. (1989)). These experiments were performed *in vitro* by incubating cells with the oligoribonucleotide. A similar procedure for *in vivo* use is described in WO 91/15580. Briefly, a pair of oligonucleotides for a given antisense RNA is produced as follows: A sequence complimentary to the first 15 bases of the open reading frame is flanked by an EcoR1 site on the 5' end and a HindIII site on the 3' end. Next, the pair of oligonucleotides is heated at 90°C for one minute and then annealed in 2X ligation buffer (20mM TRIS HCl pH 7.5, 10mM MgCl₂, 10mM dithiothreitol (DTT) and 0.2 mM ATP) and then ligated to the EcoR1/Hind III site of the retroviral vector PMV7 (WO 91/15580).

[831] For example, the 5' coding portion of a polynucleotide that encodes the polypeptide of the present invention may be used to design an antisense RNA oligonucleotide of from about 10 to 40 base pairs in length. A DNA oligonucleotide is designed to be complementary to a region of the gene involved in transcription thereby preventing transcription and the production of the receptor. The antisense RNA oligonucleotide hybridizes to the mRNA *in vivo* and blocks translation of the mRNA molecule into receptor polypeptide.

[832] In one embodiment, the antisense nucleic acid of the invention is produced intracellularly by transcription from an exogenous sequence. For example, a vector or a portion thereof, is transcribed, producing an antisense nucleic acid (RNA) of the invention. Such a vector would contain a sequence encoding the antisense nucleic acid. Such a vector can remain episomal or become chromosomally integrated, as long as it can be transcribed to produce the desired antisense RNA. Such vectors can be constructed by recombinant DNA technology methods standard in the art. Vectors can be plasmid, viral, or others known in the art, used for replication and expression in vertebrate cells. Expression of the sequence encoding the polypeptide of the present invention or fragments thereof, can be by any promoter known in the art to act in vertebrate, preferably human cells. Such promoters can be inducible or constitutive. Such promoters include, but are not limited to, the SV40 early promoter region (Bernoist and Chambon, *Nature* 29:304-310 (1981), the promoter contained in the 3' long terminal repeat of Rous sarcoma virus (Yamamoto et al., *Cell* 22:787-797 (1980), the herpes thymidine promoter (Wagner et al., *Proc. Natl. Acad. Sci. U.S.A.* 78:1441-1445 (1981), the regulatory sequences of the metallothionein gene (Brinster, et al., *Nature* 296:39-42 (1982)), etc.

[833] The antisense nucleic acids of the invention comprise a sequence complementary to at least a portion of an RNA transcript of a gene of the present invention. However, absolute complementarity, although preferred, is not required. A sequence "complementary to at least a portion of an RNA," referred to herein, means a sequence having sufficient complementarity to be able to hybridize with the RNA, forming a stable duplex; in the case of double stranded antisense nucleic acids, a single strand of the duplex DNA may thus be tested, or triplex formation may be assayed. The ability to hybridize will depend on both the degree of complementarity and the length of the antisense nucleic acid. Generally, the larger the hybridizing nucleic acid, the more base mismatches with a RNA it may contain and still form a stable duplex (or triplex as the case may be). One skilled in the art can ascertain a tolerable degree of mismatch by use of standard procedures to determine the melting point of the hybridized complex.

[834] Oligonucleotides that are complementary to the 5' end of the message, e.g., the 5' untranslated sequence up to and including the AUG initiation codon, should work most efficiently at inhibiting translation. However, sequences complementary to the 3' untranslated sequences of mRNAs have been shown to be effective at inhibiting translation of mRNAs as well. See generally, Wagner, R., 1994, *Nature* 372:333-335. Thus,

oligonucleotides complementary to either the 5'- or 3'- non- translated, non-coding regions of polynucleotide sequences described herein could be used in an antisense approach to inhibit translation of endogenous mRNA. Oligonucleotides complementary to the 5' untranslated region of the mRNA should include the complement of the AUG start codon. Antisense oligonucleotides complementary to mRNA coding regions are less efficient inhibitors of translation but could be used in accordance with the invention. Whether designed to hybridize to the 5'-, 3'- or coding region of mRNA of the present invention, antisense nucleic acids should be at least six nucleotides in length, and are preferably oligonucleotides ranging from 6 to about 50 nucleotides in length. In specific aspects the oligonucleotide is at least 10 nucleotides, at least 17 nucleotides, at least 25 nucleotides or at least 50 nucleotides.

[835] The polynucleotides of the invention can be DNA or RNA or chimeric mixtures or derivatives or modified versions thereof, single-stranded or double-stranded. The oligonucleotide can be modified at the base moiety, sugar moiety, or phosphate backbone, for example, to improve stability of the molecule, hybridization, etc. The oligonucleotide may include other appended groups such as peptides (e.g., for targeting host cell receptors *in vivo*), or agents facilitating transport across the cell membrane (see, e.g., Letsinger et al., 1989, Proc. Natl. Acad. Sci. U.S.A. 86:6553-6556; Lemaitre et al., 1987, Proc. Natl. Acad. Sci. 84:648-652; PCT Publication No. WO88/09810, published December 15, 1988) or the blood-brain barrier (see, e.g., PCT Publication No. WO89/10134, published April 25, 1988), hybridization-triggered cleavage agents. (See, e.g., Krol et al., 1988, BioTechniques 6:958-976) or intercalating agents. (See, e.g., Zon, 1988, Pharm. Res. 5:539-549). To this end, the oligonucleotide may be conjugated to another molecule, e.g., a peptide, hybridization triggered cross-linking agent, transport agent, hybridization-triggered cleavage agent, etc.

[836] The antisense oligonucleotide may comprise at least one modified base moiety which is selected from the group including, but not limited to, 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xantine, 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-

D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid (v), 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, and 2,6-diaminopurine.

[837] The antisense oligonucleotide may also comprise at least one modified sugar moiety selected from the group including, but not limited to, arabinose, 2-fluoroarabinose, xylulose, and hexose.

[838] In yet another embodiment, the antisense oligonucleotide comprises at least one modified phosphate backbone selected from the group including, but not limited to, a phosphorothioate, a phosphorodithioate, a phosphoramidothioate, a phosphoramidate, a phosphordiamidate, a methylphosphonate, an alkyl phosphotriester, and a formacetal or analog thereof.

[839] In yet another embodiment, the antisense oligonucleotide is an a-anomeric oligonucleotide. An a-anomeric oligonucleotide forms specific double-stranded hybrids with complementary RNA in which, contrary to the usual b-units, the strands run parallel to each other (Gautier et al., 1987, Nucl. Acids Res. 15:6625-6641). The oligonucleotide is a 2'-O-methylribonucleotide (Inoue et al., 1987, Nucl. Acids Res. 15:6131-6148), or a chimeric RNA-DNA analogue (Inoue et al., 1987, FEBS Lett. 215:327-330).

[840] Polynucleotides of the invention may be synthesized by standard methods known in the art, e.g. by use of an automated DNA synthesizer (such as are commercially available from Biosearch, Applied Biosystems, etc.). As examples, phosphorothioate oligonucleotides may be synthesized by the method of Stein et al. (1988, Nucl. Acids Res. 16:3209), methylphosphonate oligonucleotides can be prepared by use of controlled pore glass polymer supports (Sarin et al., 1988, Proc. Natl. Acad. Sci. U.S.A. 85:7448-7451), etc.

[841] While antisense nucleotides complementary to the coding region sequence could be used, those complementary to the transcribed untranslated region are most preferred.

[842] Potential antagonists according to the invention also include catalytic RNA, or a ribozyme (See, e.g., PCT International Publication WO 90/11364, published October 4, 1990; Sarver et al, Science 247:1222-1225 (1990). While ribozymes that cleave mRNA at site specific recognition sequences can be used to destroy mRNAs, the use of hammerhead ribozymes is preferred. Hammerhead ribozymes cleave mRNAs at locations dictated by flanking regions that form complementary base pairs with the target mRNA. The sole

requirement is that the target mRNA have the following sequence of two bases: 5'-UG-3'. The construction and production of hammerhead ribozymes is well known in the art and is described more fully in Haseloff and Gerlach, Nature 334:585-591 (1988). There are numerous potential hammerhead ribozyme cleavage sites within the nucleotide sequence of SEQ ID NO:X. Preferably, the ribozyme is engineered so that the cleavage recognition site is located near the 5' end of the mRNA; i.e., to increase efficiency and minimize the intracellular accumulation of non-functional mRNA transcripts.

[843] As in the antisense approach, the ribozymes of the invention can be composed of modified oligonucleotides (e.g., for improved stability, targeting, etc.) and should be delivered to cells which express *in vivo*. DNA constructs encoding the ribozyme may be introduced into the cell in the same manner as described above for the introduction of antisense encoding DNA. A preferred method of delivery involves using a DNA construct "encoding" the ribozyme under the control of a strong constitutive promoter, such as, for example, pol III or pol II promoter, so that transfected cells will produce sufficient quantities of the ribozyme to destroy endogenous messages and inhibit translation. Since ribozymes unlike antisense molecules, are catalytic, a lower intracellular concentration is required for efficiency.

[844] Antagonist/agonist compounds may be employed to inhibit the cell growth and proliferation effects of the polypeptides of the present invention on neoplastic cells and tissues, i.e. stimulation of angiogenesis of tumors, and, therefore, retard or prevent abnormal cellular growth and proliferation, for example, in tumor formation or growth.

[845] The antagonist/agonist may also be employed to prevent hyper-vascular diseases, and prevent the proliferation of epithelial lens cells after extracapsular cataract surgery. Prevention of the mitogenic activity of the polypeptides of the present invention may also be desirous in cases such as restenosis after balloon angioplasty.

[846] The antagonist/agonist may also be employed to prevent the growth of scar tissue during wound healing.

[847] The antagonist/agonist may also be employed to treat the diseases described herein.

[848] Thus, the invention provides a method of treating disorders or diseases, including but not limited to the disorders or diseases listed throughout this application, associated with overexpression of a polynucleotide of the present invention by administering to a

patient (a) an antisense molecule directed to the polynucleotide of the present invention, and/or (b) a ribozyme directed to the polynucleotide of the present invention.

Binding Peptides and Other Molecules

[849] The invention also encompasses screening methods for identifying polypeptides and nonpolypeptides that bind polypeptides of the invention, and the binding molecules identified thereby. These binding molecules are useful, for example, as agonists and antagonists of the polypeptides of the invention. Such agonists and antagonists can be used, in accordance with the invention, in the therapeutic embodiments described in detail, below.

This method comprises the steps of:

- a. contacting polypeptides of the invention with a plurality of molecules;
and
- b. identifying a molecule that binds the polypeptides of the invention.

[850] The step of contacting the polypeptides of the invention with the plurality of molecules may be effected in a number of ways. For example, one may contemplate immobilizing the polypeptides on a solid support and bringing a solution of the plurality of molecules in contact with the immobilized polypeptides. Such a procedure would be akin to an affinity chromatographic process, with the affinity matrix being comprised of the immobilized polypeptides of the invention. The molecules having a selective affinity for the polypeptides can then be purified by affinity selection. The nature of the solid support, process for attachment of the polypeptides to the solid support, solvent, and conditions of the affinity isolation or selection are largely conventional and well known to those of ordinary skill in the art.

[851] Alternatively, one may also separate a plurality of polypeptides into substantially separate fractions comprising a subset of or individual polypeptides. For instance, one can separate the plurality of polypeptides by gel electrophoresis, column chromatography, or like method known to those of ordinary skill for the separation of polypeptides. The individual polypeptides can also be produced by a transformed host cell in such a way as to be expressed on or about its outer surface (e.g., a recombinant phage). Individual isolates can then be "probed" by the polypeptides of the invention, optionally in the presence of an inducer should one be required for expression, to determine if any selective affinity interaction takes place between the polypeptides and the individual clone. Prior to contacting the polypeptides with each fraction comprising individual polypeptides, the

polypeptides could first be transferred to a solid support for additional convenience. Such a solid support may simply be a piece of filter membrane, such as one made of nitrocellulose or nylon. In this manner, positive clones could be identified from a collection of transformed host cells of an expression library, which harbor a DNA construct encoding a polypeptide having a selective affinity for polypeptides of the invention. Furthermore, the amino acid sequence of the polypeptide having a selective affinity for the polypeptides of the invention can be determined directly by conventional means or the coding sequence of the DNA encoding the polypeptide can frequently be determined more conveniently. The primary sequence can then be deduced from the corresponding DNA sequence. If the amino acid sequence is to be determined from the polypeptide itself, one may use microsequencing techniques. The sequencing technique may include mass spectroscopy.

[852] In certain situations, it may be desirable to wash away any unbound polypeptides from a mixture of the polypeptides of the invention and the plurality of polypeptides prior to attempting to determine or to detect the presence of a selective affinity interaction. Such a wash step may be particularly desirable when the polypeptides of the invention or the plurality of polypeptides are bound to a solid support.

[853] The plurality of molecules provided according to this method may be provided by way of diversity libraries, such as random or combinatorial peptide or nonpeptide libraries which can be screened for molecules that specifically bind polypeptides of the invention. Many libraries are known in the art that can be used, e.g., chemically synthesized libraries, recombinant (e.g., phage display libraries), and in vitro translation-based libraries. Examples of chemically synthesized libraries are described in Fodor et al., 1991, *Science* 251:767-773; Houghten et al., 1991, *Nature* 354:84-86; Lam et al., 1991, *Nature* 354:82-84; Medynski, 1994, *Bio/Technology* 12:709-710; Gallop et al., 1994, *J. Medicinal Chemistry* 37(9):1233-1251; Ohlmeyer et al., 1993, *Proc. Natl. Acad. Sci. USA* 90:10922-10926; Erb et al., 1994, *Proc. Natl. Acad. Sci. USA* 91:11422-11426; Houghten et al., 1992, *Biotechniques* 13:412; Jayawickreme et al., 1994, *Proc. Natl. Acad. Sci. USA* 91:1614-1618; Salmon et al., 1993, *Proc. Natl. Acad. Sci. USA* 90:11708-11712; PCT Publication No. WO 93/20242; and Brenner and Lerner, 1992, *Proc. Natl. Acad. Sci. USA* 89:5381-5383.

[854] Examples of phage display libraries are described in Scott and Smith, 1990, *Science* 249:386-390; Devlin et al., 1990, *Science*, 249:404-406; Christian, R. B., et al., 1992, *J. Mol. Biol.* 227:711-718; Lenstra, 1992, *J. Immunol. Meth.* 152:149-157; Kay et

al., 1993, Gene 128:59-65; and PCT Publication No. WO 94/18318 dated Aug. 18, 1994.

[855] In vitro translation-based libraries include but are not limited to those described in PCT Publication No. WO 91/05058 dated Apr. 18, 1991; and Mattheakis et al., 1994, Proc. Natl. Acad. Sci. USA 91:9022-9026.

[856] By way of examples of nonpeptide libraries, a benzodiazepine library (see e.g., Bunin et al., 1994, Proc. Natl. Acad. Sci. USA 91:4708-4712) can be adapted for use. Peptoid libraries (Simon et al., 1992, Proc. Natl. Acad. Sci. USA 89:9367-9371) can also be used. Another example of a library that can be used, in which the amide functionalities in peptides have been permethylated to generate a chemically transformed combinatorial library, is described by Ostresh et al. (1994, Proc. Natl. Acad. Sci. USA 91:11138-11142).

[857] The variety of non-peptide libraries that are useful in the present invention is great. For example, Ecker and Crooke, 1995, Bio/Technology 13:351-360 list benzodiazepines, hydantoins, piperazinediones, biphenyls, sugar analogs, beta-mercaptoketones, arylacetic acids, acylpiperidines, benzopyrans, cubanes, xanthines, aminimides, and oxazolones as among the chemical species that form the basis of various libraries.

[858] Non-peptide libraries can be classified broadly into two types: decorated monomers and oligomers. Decorated monomer libraries employ a relatively simple scaffold structure upon which a variety functional groups is added. Often the scaffold will be a molecule with a known useful pharmacological activity. For example, the scaffold might be the benzodiazepine structure.

[859] Non-peptide oligomer libraries utilize a large number of monomers that are assembled together in ways that create new shapes that depend on the order of the monomers. Among the monomer units that have been used are carbamates, pyrrolinones, and morpholinos. Peptoids, peptide-like oligomers in which the side chain is attached to the alpha amino group rather than the alpha carbon, form the basis of another version of non-peptide oligomer libraries. The first non-peptide oligomer libraries utilized a single type of monomer and thus contained a repeating backbone. Recent libraries have utilized more than one monomer, giving the libraries added flexibility.

[860] Screening the libraries can be accomplished by any of a variety of commonly known methods. See, e.g., the following references, which disclose screening of peptide libraries: Parmley and Smith, 1989, Adv. Exp. Med. Biol. 251:215-218; Scott and Smith, 1990, Science 249:386-390; Fowlkes et al., 1992; BioTechniques 13:422-427; Oldenburg et

al., 1992, Proc. Natl. Acad. Sci. USA 89:5393-5397; Yu et al., 1994, Cell 76:933-945; Staudt et al., 1988, Science 241:577-580; Bock et al., 1992, Nature 355:564-566; Tuerk et al., 1992, Proc. Natl. Acad. Sci. USA 89:6988-6992; Ellington et al., 1992, Nature 355:850-852; U.S. Pat. No. 5,096,815, U.S. Pat. No. 5,223,409, and U.S. Pat. No. 5,198,346, all to Ladner et al.; Rebar and Pabo, 1993, Science 263:671-673; and CT Publication No. WO 94/18318.

[861] In a specific embodiment, screening to identify a molecule that binds polypeptides of the invention can be carried out by contacting the library members with polypeptides of the invention immobilized on a solid phase and harvesting those library members that bind to the polypeptides of the invention. Examples of such screening methods, termed "panning" techniques are described by way of example in Parmley and Smith, 1988, Gene 73:305-318; Fowlkes et al., 1992, BioTechniques 13:422-427; PCT Publication No. WO 94/18318; and in references cited herein.

[862] In another embodiment, the two-hybrid system for selecting interacting proteins in yeast (Fields and Song, 1989, Nature 340:245-246; Chien et al., 1991, Proc. Natl. Acad. Sci. USA 88:9578-9582) can be used to identify molecules that specifically bind to polypeptides of the invention.

[863] Where the binding molecule is a polypeptide, the polypeptide can be conveniently selected from any peptide library, including random peptide libraries, combinatorial peptide libraries, or biased peptide libraries. The term "biased" is used herein to mean that the method of generating the library is manipulated so as to restrict one or more parameters that govern the diversity of the resulting collection of molecules, in this case peptides.

[864] Thus, a truly random peptide library would generate a collection of peptides in which the probability of finding a particular amino acid at a given position of the peptide is the same for all 20 amino acids. A bias can be introduced into the library, however, by specifying, for example, that a lysine occur every fifth amino acid or that positions 4, 8, and 9 of a decapeptide library be fixed to include only arginine. Clearly, many types of biases can be contemplated, and the present invention is not restricted to any particular bias. Furthermore, the present invention contemplates specific types of peptide libraries, such as phage displayed peptide libraries and those that utilize a DNA construct comprising a lambda phage vector with a DNA insert.

[865] As mentioned above, in the case of a binding molecule that is a polypeptide, the

09550083-091201

polypeptide may have about 6 to less than about 60 amino acid residues, preferably about 6 to about 10 amino acid residues, and most preferably, about 6 to about 22 amino acids. In another embodiment, a binding polypeptide has in the range of 15-100 amino acids, or 20-50 amino acids.

[866] The selected binding polypeptide can be obtained by chemical synthesis or recombinant expression.

Other Activities

[867] A polypeptide, polynucleotide, agonist, or antagonist of the present invention, as a result of the ability to stimulate vascular endothelial cell growth, may be employed in treatment for stimulating re-vascularization of ischemic tissues due to various disease conditions such as thrombosis, arteriosclerosis, and other cardiovascular conditions. The polypeptide, polynucleotide, agonist, or antagonist of the present invention may also be employed to stimulate angiogenesis and limb regeneration, as discussed above.

[868] A polypeptide, polynucleotide, agonist, or antagonist of the present invention may also be employed for treating wounds due to injuries, burns, post-operative tissue repair, and ulcers since they are mitogenic to various cells of different origins, such as fibroblast cells and skeletal muscle cells, and therefore, facilitate the repair or replacement of damaged or diseased tissue.

[869] A polypeptide, polynucleotide, agonist, or antagonist of the present invention may also be employed stimulate neuronal growth and to treat and prevent neuronal damage which occurs in certain neuronal disorders or neuro-degenerative conditions such as Alzheimer's disease, Parkinson's disease, and AIDS-related complex. A polypeptide, polynucleotide, agonist, or antagonist of the present invention may have the ability to stimulate chondrocyte growth, therefore, they may be employed to enhance bone and periodontal regeneration and aid in tissue transplants or bone grafts.

[870] A polypeptide, polynucleotide, agonist, or antagonist of the present invention may be also be employed to prevent skin aging due to sunburn by stimulating keratinocyte growth.

[871] A polypeptide, polynucleotide, agonist, or antagonist of the present invention may also be employed for preventing hair loss, since FGF family members activate hair-forming cells and promotes melanocyte growth. Along the same lines, a polypeptide, polynucleotide, agonist, or antagonist of the present invention may be employed to

05550083-091201

stimulate growth and differentiation of hematopoietic cells and bone marrow cells when used in combination with other cytokines.

[872] A polypeptide, polynucleotide, agonist, or antagonist of the present invention may also be employed to maintain organs before transplantation or for supporting cell culture of primary tissues. A polypeptide, polynucleotide, agonist, or antagonist of the present invention may also be employed for inducing tissue of mesodermal origin to differentiate in early embryos.

[873] A polypeptide, polynucleotide, agonist, or antagonist of the present invention may also increase or decrease the differentiation or proliferation of embryonic stem cells, besides, as discussed above, hematopoietic lineage.

[874] A polypeptide, polynucleotide, agonist, or antagonist of the present invention may also be used to modulate mammalian characteristics, such as body height, weight, hair color, eye color, skin, percentage of adipose tissue, pigmentation, size, and shape (e.g., cosmetic surgery). Similarly, a polypeptide, polynucleotide, agonist, or antagonist of the present invention may be used to modulate mammalian metabolism affecting catabolism, anabolism, processing, utilization, and storage of energy.

[875] A polypeptide, polynucleotide, agonist, or antagonist of the present invention may be used to change a mammal's mental state or physical state by influencing biorhythms, cardiac rhythms, depression (including depressive disorders), tendency for violence, tolerance for pain, reproductive capabilities (preferably by Activin or Inhibin-like activity), hormonal or endocrine levels, appetite, libido, memory, stress, or other cognitive qualities.

[876] A polypeptide, polynucleotide, agonist, or antagonist of the present invention may also be used as a food additive or preservative, such as to increase or decrease storage capabilities, fat content, lipid, protein, carbohydrate, vitamins, minerals, cofactors or other nutritional components.

[877] The above-recited applications have uses in a wide variety of hosts. Such hosts include, but are not limited to, human, murine, rabbit, goat, guinea pig, camel, horse, mouse, rat, hamster, pig, micro-pig, chicken, goat, cow, sheep, dog, cat, non-human primate, and human. In specific embodiments, the host is a mouse, rabbit, goat, guinea pig, chicken, rat, hamster, pig, sheep, dog or cat. In preferred embodiments, the host is a mammal. In most preferred embodiments, the host is a human.

Other Preferred Embodiments

[878] Other preferred embodiments of the claimed invention include an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to a sequence of at least about 50 contiguous nucleotides in the nucleotide sequence of SEQ ID NO:X or the complementary strand thereto, the nucleotide sequence as defined in column 5 of Table 1B or columns 8 and 9 of Table 2 or the complementary strand thereto, and/or cDNA contained in Clone ID NO:Z.

[879] Also preferred is a nucleic acid molecule wherein said sequence of contiguous nucleotides is included in the nucleotide sequence of the portion of SEQ ID NO:X as defined in column 5, "ORF (From-To)", in Table 1B.

[880] Also preferred is a nucleic acid molecule wherein said sequence of contiguous nucleotides is included in the nucleotide sequence of the portion of SEQ ID NO:X as defined in columns 8 and 9, "NT From" and "NT To" respectively, in Table 2.

[881] Also preferred is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to a sequence of at least about 150 contiguous nucleotides in the nucleotide sequence of SEQ ID NO:X or the complementary strand thereto, the nucleotide sequence as defined in column 5 of Table 1B or columns 8 and 9 of Table 2 or the complementary strand thereto, and/or cDNA contained in Clone ID NO:Z.

[882] Further preferred is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to a sequence of at least about 500 contiguous nucleotides in the nucleotide sequence of SEQ ID NO:X or the complementary strand thereto, the nucleotide sequence as defined in column 5 of Table 1B or columns 8 and 9 of Table 2 or the complementary strand thereto, and/or cDNA contained in Clone ID NO:Z.

[883] A further preferred embodiment is a nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to the nucleotide sequence of the portion of SEQ ID NO:X defined in column 5, "ORF (From-To)", in Table 1B.

[884] A further preferred embodiment is a nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to the nucleotide sequence of the portion of SEQ ID NO:X defined in columns 8 and 9, "NT From" and "NT To", respectively, in Table 2.

[885] A further preferred embodiment is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to the complete nucleotide sequence of SEQ ID NO:X or the complementary strand thereto, the nucleotide sequence as defined in

09950003-091201

column 5 of Table 1B or columns 8 and 9 of Table 2 or the complementary strand thereto, and/or cDNA contained in Clone ID NO:Z.

[886] Also preferred is an isolated nucleic acid molecule which hybridizes under stringent hybridization conditions to a nucleic acid molecule comprising a nucleotide sequence of SEQ ID NO:X or the complementary strand thereto, the nucleotide sequence as defined in column 5 of Table 1B or columns 8 and 9 of Table 2 or the complementary strand thereto, and/or cDNA contained in Clone ID NO:Z, wherein said nucleic acid molecule which hybridizes does not hybridize under stringent hybridization conditions to a nucleic acid molecule having a nucleotide sequence consisting of only A residues or of only T residues.

[887] Also preferred is a composition of matter comprising a DNA molecule which comprises the cDNA contained in Clone ID NO:Z.

[888] Also preferred is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to a sequence of at least 50 contiguous nucleotides of the cDNA sequence contained in Clone ID NO:Z.

[889] Also preferred is an isolated nucleic acid molecule, wherein said sequence of at least 50 contiguous nucleotides is included in the nucleotide sequence of an open reading frame sequence encoded by cDNA contained in Clone ID NO:Z.

[890] Also preferred is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to sequence of at least 150 contiguous nucleotides in the nucleotide sequence encoded by cDNA contained in Clone ID NO:Z.

[891] A further preferred embodiment is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to sequence of at least 500 contiguous nucleotides in the nucleotide sequence encoded by cDNA contained in Clone ID NO:Z.

[892] A further preferred embodiment is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to the complete nucleotide sequence encoded by cDNA contained in Clone ID NO:Z.

[893] A further preferred embodiment is a method for detecting in a biological sample a nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from the group consisting of: a nucleotide sequence of SEQ ID NO:X or the complementary strand thereto; the nucleotide sequence as defined in column 5 of Table 1B or columns 8 and 9 of Table 2 or the complementary strand thereto; and a nucleotide sequence encoded by cDNA

contained in Clone ID NO:Z; which method comprises a step of comparing a nucleotide sequence of at least one nucleic acid molecule in said sample with a sequence selected from said group and determining whether the sequence of said nucleic acid molecule in said sample is at least 95% identical to said selected sequence.

[894] Also preferred is the above method wherein said step of comparing sequences comprises determining the extent of nucleic acid hybridization between nucleic acid molecules in said sample and a nucleic acid molecule comprising said sequence selected from said group. Similarly, also preferred is the above method wherein said step of comparing sequences is performed by comparing the nucleotide sequence determined from a nucleic acid molecule in said sample with said sequence selected from said group. The nucleic acid molecules can comprise DNA molecules or RNA molecules.

[895] A further preferred embodiment is a method for identifying the species, tissue or cell type of a biological sample which method comprises a step of detecting nucleic acid molecules in said sample, if any, comprising a nucleotide sequence that is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from the group consisting of: a nucleotide sequence of SEQ ID NO:X or the complementary strand thereto; the nucleotide sequence as defined in column 5 of Table 1B or columns 8 and 9 of Table 2 or the complementary strand thereto; and a nucleotide sequence of the cDNA contained in Clone ID NO:Z.

[896] The method for identifying the species, tissue or cell type of a biological sample can comprise a step of detecting nucleic acid molecules comprising a nucleotide sequence in a panel of at least two nucleotide sequences, wherein at least one sequence in said panel is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from said group.

[897] Also preferred is a method for diagnosing in a subject a pathological condition associated with abnormal structure or expression of a nucleotide sequence of SEQ ID NO:X or the complementary strand thereto; the nucleotide sequence as defined in column 5 of Table 1B or columns 8 and 9 of Table 2 or the complementary strand thereto; or the cDNA contained in Clone ID NO:Z which encodes a protein, wherein the method comprises a step of detecting in a biological sample obtained from said subject nucleic acid molecules, if any, comprising a nucleotide sequence that is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from the group consisting of: a nucleotide sequence of SEQ ID NO:X or the complementary strand thereto; the nucleotide sequence as

defined in column 5 of Table 1B or columns 8 and 9 of Table 2 or the complementary strand thereto; and a nucleotide sequence of cDNA contained in Clone ID NO:Z.

[898] The method for diagnosing a pathological condition can comprise a step of detecting nucleic acid molecules comprising a nucleotide sequence in a panel of at least two nucleotide sequences, wherein at least one sequence in said panel is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from said group.

[899] Also preferred is a composition of matter comprising isolated nucleic acid molecules wherein the nucleotide sequences of said nucleic acid molecules comprise a panel of at least two nucleotide sequences, wherein at least one sequence in said panel is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from the group consisting of: a nucleotide sequence of SEQ ID NO:X or the complementary strand thereto; the nucleotide sequence as defined in column 5 of Table 1B or columns 8 and 9 of Table 2 or the complementary strand thereto; and a nucleotide sequence encoded by cDNA contained in Clone ID NO:Z. The nucleic acid molecules can comprise DNA molecules or RNA molecules.

[900] Also preferred is a composition of matter comprising isolated nucleic acid molecules wherein the nucleotide sequences of said nucleic acid molecules comprise a DNA microarray or "chip" of at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 40, 50, 100, 150, 200, 250, 300, 500, 1000, 2000, 3000, or 4000 nucleotide sequences, wherein at least one sequence in said DNA microarray or "chip" is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from the group consisting of: a nucleotide sequence of SEQ ID NO:X wherein X is any integer as defined in Table 1A and/or 1B; and a nucleotide sequence encoded by a human cDNA clone identified by a cDNA "Clone ID" in Table 1A and/or 1B.

[901] Also preferred is an isolated polypeptide comprising an amino acid sequence at least 90% identical to a sequence of at least about 10 contiguous amino acids in the polypeptide sequence of SEQ ID NO:Y; a polypeptide encoded by SEQ ID NO:X or the complementary strand thereto; the polypeptide encoded by the nucleotide sequence as defined in columns 8 and 9 of Table 2; and/or a polypeptide encoded by cDNA contained in Clone ID NO:Z.

[902] Also preferred is an isolated polypeptide comprising an amino acid sequence at least 95% identical to a sequence of at least about 30 contiguous amino acids in the amino acid sequence of SEQ ID NO:Y; a polypeptide encoded by SEQ ID NO:X or the

complementary strand thereto; the polypeptide encoded by the nucleotide sequence as defined in columns 8 and 9 of Table 2; and/or a polypeptide encoded by cDNA contained in Clone ID NO:Z.

[903] Further preferred is an isolated polypeptide comprising an amino acid sequence at least 95% identical to a sequence of at least about 100 contiguous amino acids in the amino acid sequence of SEQ ID NO:Y; a polypeptide encoded by SEQ ID NO:X or the complementary strand thereto; the polypeptide encoded by the nucleotide sequence as defined in columns 8 and 9 of Table 2; and/or a polypeptide encoded by cDNA contained in Clone ID NO:Z.

[904] Further preferred is an isolated polypeptide comprising an amino acid sequence at least 95% identical to the complete amino acid sequence of SEQ ID NO:Y; a polypeptide encoded by SEQ ID NO:X or the complementary strand thereto; the polypeptide encoded by the nucleotide sequence as defined in columns 8 and 9 of Table 2; and/or a polypeptide encoded by cDNA contained in Clone ID NO:Z.

[905] Further preferred is an isolated polypeptide comprising an amino acid sequence at least 90% identical to a sequence of at least about 10 contiguous amino acids in the complete amino acid sequence of a polypeptide encoded by contained in Clone ID NO:Z

[906] Also preferred is a polypeptide wherein said sequence of contiguous amino acids is included in the amino acid sequence of a portion of said polypeptide encoded by cDNA contained in Clone ID NO:Z; a polypeptide encoded by SEQ ID NO:X or the complementary strand thereto; the polypeptide encoded by the nucleotide sequence as defined in columns 8 and 9 of Table 2; and/or the polypeptide sequence of SEQ ID NO:Y.

[907] Also preferred is an isolated polypeptide comprising an amino acid sequence at least 95% identical to a sequence of at least about 30 contiguous amino acids in the amino acid sequence of a polypeptide encoded by the cDNA contained in Clone ID NO:Z.

[908] Also preferred is an isolated polypeptide comprising an amino acid sequence at least 95% identical to a sequence of at least about 100 contiguous amino acids in the amino acid sequence of a polypeptide encoded by cDNA contained in Clone ID NO:Z.

[909] Also preferred is an isolated polypeptide comprising an amino acid sequence at least 95% identical to the amino acid sequence of a polypeptide encoded by the cDNA contained in Clone ID NO:Z.

[910] Further preferred is an isolated antibody which binds specifically to a polypeptide comprising an amino acid sequence that is at least 90% identical to a sequence of at least 10

09950083.091201

contiguous amino acids in a sequence selected from the group consisting of: a polypeptide sequence of SEQ ID NO:Y; a polypeptide encoded by SEQ ID NO:X or the complementary strand thereto; the polypeptide encoded by the nucleotide sequence as defined in columns 8 and 9 of Table 2; and a polypeptide encoded by the cDNA contained in Clone ID NO:Z.

[911] Further preferred is a method for detecting in a biological sample a polypeptide comprising an amino acid sequence which is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: a polypeptide sequence of SEQ ID NO:Y; a polypeptide encoded by SEQ ID NO:X or the complementary strand thereto; the polypeptide encoded by the nucleotide sequence as defined in columns 8 and 9 of Table 2; and a polypeptide encoded by the cDNA contained in Clone ID NO:Z; which method comprises a step of comparing an amino acid sequence of at least one polypeptide molecule in said sample with a sequence selected from said group and determining whether the sequence of said polypeptide molecule in said sample is at least 90% identical to said sequence of at least 10 contiguous amino acids.

[912] Also preferred is the above method wherein said step of comparing an amino acid sequence of at least one polypeptide molecule in said sample with a sequence selected from said group comprises determining the extent of specific binding of polypeptides in said sample to an antibody which binds specifically to a polypeptide comprising an amino acid sequence that is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: a polypeptide sequence of SEQ ID NO:Y; a polypeptide encoded by SEQ ID NO:X or the complementary strand thereto; the polypeptide encoded by the nucleotide sequence as defined in columns 8 and 9 of Table 2; and a polypeptide encoded by the cDNA contained in Clone ID NO:Z.

[913] Also preferred is the above method wherein said step of comparing sequences is performed by comparing the amino acid sequence determined from a polypeptide molecule in said sample with said sequence selected from said group.

[914] Also preferred is a method for identifying the species, tissue or cell type of a biological sample which method comprises a step of detecting polypeptide molecules in said sample, if any, comprising an amino acid sequence that is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: polypeptide sequence of SEQ ID NO:Y; a polypeptide encoded by SEQ ID NO:X or the complementary strand thereto; the polypeptide encoded by the nucleotide

sequence as defined in columns 8 and 9 of Table 2; and a polypeptide encoded by the cDNA contained in Clone ID NO:Z.

[915] Also preferred is the above method for identifying the species, tissue or cell type of a biological sample, which method comprises a step of detecting polypeptide molecules comprising an amino acid sequence in a panel of at least two amino acid sequences, wherein at least one sequence in said panel is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the above group.

[916] Also preferred is a method for diagnosing in a subject a pathological condition associated with abnormal structure or expression of a nucleic acid sequence identified in Table 1A, 1B or Table 2 encoding a polypeptide, which method comprises a step of detecting in a biological sample obtained from said subject polypeptide molecules comprising an amino acid sequence in a panel of at least two amino acid sequences, wherein at least one sequence in said panel is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: polypeptide sequence of SEQ ID NO:Y; a polypeptide encoded by SEQ ID NO:X or the complementary strand thereto; the polypeptide encoded by the nucleotide sequence as defined in columns 8 and 9 of Table 2; and a polypeptide encoded by the cDNA contained in Clone ID NO:Z.

[917] In any of these methods, the step of detecting said polypeptide molecules includes using an antibody.

[918] Also preferred is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to a nucleotide sequence encoding a polypeptide wherein said polypeptide comprises an amino acid sequence that is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: polypeptide sequence of SEQ ID NO:Y; a polypeptide encoded by SEQ ID NO:X or the complementary strand thereto; the polypeptide encoded by the nucleotide sequence as defined in columns 8 and 9 of Table 2; and a polypeptide encoded by the cDNA contained in Clone ID NO:Z.

[919] Also preferred is an isolated nucleic acid molecule, wherein said nucleotide sequence encoding a polypeptide has been optimized for expression of said polypeptide in a prokaryotic host.

[920] Also preferred is a polypeptide molecule, wherein said polypeptide comprises an amino acid sequence selected from the group consisting of: polypeptide sequence of SEQ

agent, or antibody of the claimed invention that are associated with toxin or cytotoxic prodrugs.

[926] Having generally described the invention, the same will be more readily understood by reference to the following examples, which are provided by way of illustration and are not intended as limiting.

0950083-091201
T02T60" (8005560

Table 6

ATCC Deposits	Deposit Date	ATCC Designation Number
LP01, LP02, LP03, LP04, LP05, LP06, LP07, LP08, LP09, LP10, LP11,	May-20-97	209059, 209060, 209061, 209062, 209063, 209064, 209065, 209066, 209067, 209068, 209069
LP12	Jan-12-98	209579
LP13	Jan-12-98	209578
LP14	Jul-16-98	203067
LP15	Jul-16-98	203068
LP16	Feb-1-99	203609
LP17	Feb-1-99	203610
LP20	Nov-17-98	203485
LP21	Jun-18-99	PTA-252
LP22	Jun-18-99	PTA-253
LP23	Dec-22-99	PTA-1081

Examples

Example 1: Isolation of a Selected cDNA Clone From the Deposited Sample

[927] Each Clone ID NO:Z is contained in a plasmid vector. Table 7 identifies the vectors used to construct the cDNA library from which each clone was isolated. In many cases, the vector used to construct the library is a phage vector from which a plasmid has been excised. The following correlates the related plasmid for each phage vector used in constructing the cDNA library. For example, where a particular clone is identified in Table 7 as being isolated in the vector "Lambda Zap," the corresponding deposited clone is in "pBluescript."

<u>Vector Used to Construct Library</u>	<u>Corresponding Deposited Plasmid</u>
Lambda Zap	pBluescript (pBS)
Uni-Zap XR	pBluescript (pBS)
Zap Express	pBK
lafmid BA	plafmid BA
pSport1	pSport1
pCMVSPORT 2.0	pCMVSPORT 2.0
pCMVSPORT 3.0	pCMVSPORT 3.0
pCR [®] 2.1	pCR [®] 2.1

[928] Vectors Lambda Zap (U.S. Patent Nos. 5,128,256 and 5,286,636), Uni-Zap XR (U.S. Patent Nos. 5,128, 256 and 5,286,636), Zap Express (U.S. Patent Nos. 5,128,256 and 5,286,636), pBluescript (pBS) (Short, J. M. et al., Nucleic Acids Res. 16:7583-7600 (1988); Alting-Mees, M. A. and Short, J. M., Nucleic Acids Res. 17:9494 (1989)) and pBK (Alting-Mees, M. A. et al., Strategies 5:58-61 (1992)) are commercially available from Stratagene Cloning Systems, Inc., 11011 N. Torrey Pines Road, La Jolla, CA, 92037. pBS contains an ampicillin resistance gene and pBK contains a neomycin resistance gene. Both can be transformed into E. coli strain XL-1 Blue, also available from Stratagene. pBS comes in 4 forms SK+, SK-, KS+ and KS-. The S and K refers to the orientation of the polylinker to the T7 and T3 primer sequences which flank the polylinker region ("S" is for SacI and "K" is for KpnI which are the first sites on each respective end of the linker). "+" or "-" refer to the orientation of the fl origin of replication ("ori"), such that in one orientation, single

stranded rescue initiated from the f1 ori generates sense strand DNA and in the other, antisense.

[929] Vectors pSport1, pCMVSPORT 2.0 and pCMVSPORT 3.0, were obtained from Life Technologies, Inc., P. O. Box 6009, Gaithersburg, MD 20897. All Sport vectors contain an ampicillin resistance gene and may be transformed into *E. coli* strain DH10B, also available from Life Technologies. (See, for instance, Gruber, C. E., et al., Focus 15:59 (1993)). Vector lafmid BA (Bento Soares, Columbia University, NY) contains an ampicillin resistance gene and can be transformed into *E. coli* strain XL-1 Blue. Vector pCR[®]2.1, which is available from Invitrogen, 1600 Faraday Avenue, Carlsbad, CA 92008, contains an ampicillin resistance gene and may be transformed into *E. coli* strain DH10B, available from Life Technologies. (See, for instance, Clark, J. M., Nuc. Acids Res. 16:9677-9686 (1988) and Mead, D. et al., Bio/Technology 9: (1991)). Preferably, a polynucleotide of the present invention does not comprise the phage vector sequences identified for the particular clone in Table 7, as well as the corresponding plasmid vector sequences designated above.

[930] The deposited material in the sample assigned the ATCC Deposit Number cited by reference to Tables 1, 2, 6 and 7 for any given cDNA clone also may contain one or more additional plasmids, each comprising a cDNA clone different from that given clone. Thus, deposits sharing the same ATCC Deposit Number contain at least a plasmid for each Clone ID NO:Z.

TABLE 7

Libraries owned by Catalog	Catalog Description	Vector	ATCC Deposit
HUKA HUKB HUKC HUKD HUKF HUKG	Human Uterine Cancer	Lambda ZAP II	LP01
HCNA HCNB	Human Colon	Lambda Zap II	LP01
HFFA	Human Fetal Brain, random primed	Lambda Zap II	LP01
HTWA	Resting T-Cell	Lambda ZAP II	LP01
HBQA	Early Stage Human Brain, random primed	Lambda ZAP II	LP01
HLMB HLMF HLMG HLMH HLMI HLMJ HLMM HLMN	breast lymph node CDNA library	Lambda ZAP II	LP01
HCQA HCQB	human colon cancer	Lambda ZAP II	LP01
HMEA HMEC HMED HMEF HMEG HMEI HMEJ HMEK HMEI	Human Microvascular Endothelial Cells, fract. A	Lambda ZAP II	LP01
HUSA HUSC	Human Umbilical Vein Endothelial	Lambda ZAP II	LP01

Libraries owned by Catalog	Catalog Description	Vector	ATCC Deposit
	Cells, fract. A		
HLQA HLQB	Hepatocellular Tumor	Lambda ZAP II	LP01
HHGA HHGB HHGC HHGD	Hemangiopericytoma	Lambda ZAP II	LP01
HSDM	Human Striatum Depression, re-rescue	Lambda ZAP II	LP01
HUSH	H Umbilical Vein Endothelial Cells, frac A, re-excision	Lambda ZAP II	LP01
HSGS	Salivary gland, subtracted	Lambda ZAP II	LP01
HFXA HFXB HFXC HFXD HFXE HFXF HFXG HFXH	Brain frontal cortex	Lambda ZAP II	LP01
HPQA HPQB HPQC	PERM TF274	Lambda ZAP II	LP01
HFXJ HFXK	Brain Frontal Cortex, re-excision	Lambda ZAP II	LP01
HCWA HCWB HCWC HCWD HCWE HCWF HCWG HCWH HCWI HCWJ HCWK	CD34 positive cells (Cord Blood)	ZAP Express	LP02
HCUA HCUB HCUC	CD34 depleted Buffy Coat (Cord Blood)	ZAP Express	LP02
HRSM	A-14 cell line	ZAP Express	LP02
HRSA	A1-CELL LINE	ZAP Express	LP02
HCUD HCUE HCUF HCUG HCUH HCUI	CD34 depleted Buffy Coat (Cord Blood), re-excision	ZAP Express	LP02
HBXE HBXF HBXG	H. Whole Brain #2, re-excision	ZAP Express	LP02
HRLM	L8 cell line	ZAP Express	LP02
HBXA HBXB HBXC HBXD	Human Whole Brain #2 - Oligo dT > 1.5Kb	ZAP Express	LP02
HUDA HUDB HUDC	Testes	ZAP Express	LP02
HHTM HHTN HHTO	H. hypothalamus, frac A; re-excision	ZAP Express	LP02
HHTL	H. hypothalamus, frac A	ZAP Express	LP02
HASA HASD	Human Adult Spleen	Uni-ZAP XR	LP03
HFKC HFKD HFKE HFKF HFKG	Human Fetal Kidney	Uni-ZAP XR	LP03
HE8A HE8B HE8C HE8D HE8E HE8F HE8M HE8N	Human 8 Week Whole Embryo	Uni-ZAP XR	LP03
HGBA HGBD HGBE HGBF HGBG HGBH HGBI	Human Gall Bladder	Uni-ZAP XR	LP03
HLHA HLHB HLHC HLHD HLHE HLHF HLHG HLHH HLHQ	Human Fetal Lung III	Uni-ZAP XR	LP03
HPMA HPMB HPMC HPMD HPME HPMF HPMG HPMH	Human Placenta	Uni-ZAP XR	LP03
HPRA HPRB HPRC HPRD	Human Prostate	Uni-ZAP XR	LP03
HSIA HSIC HSID HSIE	Human Adult Small Intestine	Uni-ZAP XR	LP03
HTEA HTEB HTEC HTED HTEE	Human Testes	Uni-ZAP XR	LP03

Libraries owned by Catalog	Catalog Description	Vector	ATCC Deposit
HTEF HTEG HTEH HTEI HTEJ HTEK			
HTPA HTPB HTPC HTPD HTPE	Human Pancreas Tumor	Uni-ZAP XR	LP03
HTTA HTTB HTTC HTTD HTTE HTTF	Human Testes Tumor	Uni-ZAP XR	LP03
HAPA HAPB HAPC HAPM	Human Adult Pulmonary	Uni-ZAP XR	LP03
HETA HETB HETC HETD HETE HETF HETG HETH HETI	Human Endometrial Tumor	Uni-ZAP XR	LP03
HHFB HHFC HHFD HHFE HHFF HHFG HHFH HHFI	Human Fetal Heart	Uni-ZAP XR	LP03
HHPB HHPC HHPD HHPE HHPP HHPG HHPH	Human Hippocampus	Uni-ZAP XR	LP03
HCE1 HCE2 HCE3 HCE4 HCE5 HCEB HCEC HCED HCEE HCEF HCEG	Human Cerebellum	Uni-ZAP XR	LP03
HUVB HUV C HUVD HUVE	Human Umbilical Vein, Endo. remake	Uni-ZAP XR	LP03
HSTA HSTB HSTC HSTD	Human Skin Tumor	Uni-ZAP XR	LP03
HTAA HTAB HTAC HTAD HTAE	Human Activated T-Cells	Uni-ZAP XR	LP03
HFEA HFEB HFEC	Human Fetal Epithelium (Skin)	Uni-ZAP XR	LP03
HJPA HJPB HJPC HJPD	HUMAN JURKAT MEMBRANE BOUND POLYSOMES	Uni-ZAP XR	LP03
HESA	Human epithelioid sarcoma	Uni-Zap XR	LP03
HLTA HLTB HLTC HLTD HLTE HLTF	Human T-Cell Lymphoma	Uni-ZAP XR	LP03
HFTA HFTB HFTC HFTD	Human Fetal Dura Mater	Uni-ZAP XR	LP03
HRDA HRDB HRDC HRDD HRDE HRDF	Human Rhabdomyosarcoma	Uni-ZAP XR	LP03
HCAA HCAB HCAC	Cem cells cyclohexamide treated	Uni-ZAP XR	LP03
HRGA HRGB HRGC HRGD	Raji Cells, cyclohexamide treated	Uni-ZAP XR	LP03
HSUA HSUB HSUC HSUM	Supt Cells, cyclohexamide treated	Uni-ZAP XR	LP03
HT4A HT4C HT4D	Activated T-Cells, 12 hrs.	Uni-ZAP XR	LP03
HE9A HE9B HE9C HE9D HE9E HE9F HE9G HE9H HE9M HE9N	Nine Week Old Early Stage Human	Uni-ZAP XR	LP03
HATA HATB HATC HATD HATE	Human Adrenal Gland Tumor	Uni-ZAP XR	LP03
HT5A	Activated T-Cells, 24 hrs.	Uni-ZAP XR	LP03
HFGA HFGM	Human Fetal Brain	Uni-ZAP XR	LP03
HNEA HNEB HNEC HNE D HNEE	Human Neutrophil	Uni-ZAP XR	LP03
HBGB HBGD	Human Primary Breast Cancer	Uni-ZAP XR	LP03
HBNA HBNB	Human Normal Breast	Uni-ZAP XR	LP03
HCAS	Cem Cells, cyclohexamide treated, subtra	Uni-ZAP XR	LP03
HHPS	Human Hippocampus, subtracted	pBS	LP03

Libraries owned by Catalog	Catalog Description	Vector	ATCC Deposit
HKCS HKCU	Human Colon Cancer, subtracted	pBS	LP03
HRGS	Raji cells, cyclohexamide treated, subtracted	pBS	LP03
HSUT	Supt cells, cyclohexamide treated, differentially expressed	pBS	LP03
HT4S	Activated T-Cells, 12 hrs, subtracted	Uni-ZAP XR	LP03
HCDA HCDB HCDC HCDD HCDE	Human Chondrosarcoma	Uni-ZAP XR	LP03
HOAA HOAB HOAC	Human Osteosarcoma	Uni-ZAP XR	LP03
HTLA HTLB HTLC HTLD HTLE HTLF	Human adult testis, large inserts	Uni-ZAP XR	LP03
HLMA HLMC HLMD	Breast Lymph node cDNA library	Uni-ZAP XR	LP03
H6EA H6EB H6EC	HL-60, PMA 4H	Uni-ZAP XR	LP03
HTXA HTXB HTXC HTXD HTXE HTXF HTXG HTXH	Activated T-Cell (12hs)/Thiouridine labelledEco	Uni-ZAP XR	LP03
HNFA HNFB HNFC HNFD HNFE HNFF HNFG HNFH HNFJ	Human Neutrophil, Activated	Uni-ZAP XR	LP03
HTOB HTOC	HUMAN TONSILS, FRACTION 2	Uni-ZAP XR	LP03
HMGB	Human OB MG63 control fraction I	Uni-ZAP XR	LP03
HOPB	Human OB HOS control fraction I	Uni-ZAP XR	LP03
HORB	Human OB HOS treated (10 nM E2) fraction I	Uni-ZAP XR	LP03
HSVA HSVB HSVC	Human Chronic Synovitis	Uni-ZAP XR	LP03
HROA	HUMAN STOMACH	Uni-ZAP XR	LP03
HBJA HBJB HBJC HBJD HBJE HBJF HBJG HBJH HBJI HBJJ HBJK	HUMAN B CELL LYMPHOMA	Uni-ZAP XR	LP03
HCRA HCRB HCRC	human corpus colosum	Uni-ZAP XR	LP03
HODA HODB HODC HODD	human ovarian cancer	Uni-ZAP XR	LP03
HDSA	Dermatofibrosarcoma Protuberance	Uni-ZAP XR	LP03
HMWA HMWB HMWC HMWD HMWE HMWF HMWG HMWH HMWI HMWJ	Bone Marrow Cell Line (RS4;11)	Uni-ZAP XR	LP03
HSOA	stomach cancer (human)	Uni-ZAP XR	LP03
HERA	SKIN	Uni-ZAP XR	LP03
HMDA	Brain-medulloblastoma	Uni-ZAP XR	LP03
HGLA HGLB HGLD	Glioblastoma	Uni-ZAP XR	LP03
HEAA	H. Atrophic Endometrium	Uni-ZAP XR	LP03
HBCA HBCB	H. Lymph node breast Cancer	Uni-ZAP XR	LP03
HPWT	Human Prostate BPH, re-excision	Uni-ZAP XR	LP03
HFVG HFVH HFVI	Fetal Liver, subtraction II	pBS	LP03
HNFI	Human Neutrophils, Activated, re-	pBS	LP03

Libraries owned by Catalog	Catalog Description	Vector	ATCC Deposit
	excision		
HBMB HBMC HBMD	Human Bone Marrow, re-excision	pBS	LP03
HKML HKMM HKMN	H. Kidney Medulla, re-excision	pBS	LP03
HKIX HKIY	H. Kidney Cortex, subtracted	pBS	LP03
HADT	H. Amygdala Depression, subtracted	pBS	LP03
H6AS	HL-60, untreated, subtracted	Uni-ZAP XR	LP03
H6ES	HL-60, PMA 4H, subtracted	Uni-ZAP XR	LP03
H6BS	HL-60, RA 4h, Subtracted	Uni-ZAP XR	LP03
H6CS	HL-60, PMA 1d, subtracted	Uni-ZAP XR	LP03
HTXJ HTXK	Activated T-cell(12h)/Thiouridine-re-excision	Uni-ZAP XR	LP03
HMSA HMSB HMSC HMSE HMSF HMSG HMSH HMSI HMSJ HMSK	Monocyte activated	Uni-ZAP XR	LP03
HAGA HAGB HAGC HAGD HAGE HAGF	Human Amygdala	Uni-ZAP XR	LP03
HSRA HSRB HSRE	STROMAL -OSTEOCLASTOMA	Uni-ZAP XR	LP03
HSRD HSRF HSRG HSRH	Human Osteoclastoma Stromal Cells - unamplified	Uni-ZAP XR	LP03
HSQA HSQB HSQC HSQD HSQE HSQF HSQG	Stromal cell TF274	Uni-ZAP XR	LP03
HSKA HSKB HSKC HSKD HSKE HSKF HSKZ	Smooth muscle, serum treated	Uni-ZAP XR	LP03
HSLA HSLB HSLC HSLD HSLE HSLF HSLG	Smooth muscle,control	Uni-ZAP XR	LP03
HSDA HSDD HSDE HSDF HSDG HSDH	Spinal cord	Uni-ZAP XR	LP03
HPWS	Prostate-BPH subtracted II	pBS	LP03
HSKW HSKX HSKY	Smooth Muscle- HASTE normalized	pBS	LP03
HFPB HFPC HFPD	H. Frontal cortex,epileptic;re-excision	Uni-ZAP XR	LP03
HSDI HSDJ HSDK	Spinal Cord, re-excision	Uni-ZAP XR	LP03
HSKN HSKO	Smooth Muscle Serum Treated, Norm	pBS	LP03
HSKG HSKH HSKI	Smooth muscle, serum induced,re-exc	pBS	LP03
HFCA HFCE HFCC HFCD HFCE HFCE	Human Fetal Brain	Uni-ZAP XR	LP04
HPTA HPTB HPTD	Human Pituitary	Uni-ZAP XR	LP04
HTHB HTHC HTHD	Human Thymus	Uni-ZAP XR	LP04
HE6B HE6C HE6D HE6E HE6F HE6G HE6S	Human Whole Six Week Old Embryo	Uni-ZAP XR	LP04
HSSA HSSB HSSC HSSD HSSE HSSF	Human Synovial Sarcoma	Uni-ZAP XR	LP04

Libraries owned by Catalog	Catalog Description	Vector	ATCC Deposit
HSSG HSSH HSSI HSSJ HSSK			
HE7T	7 Week Old Early Stage Human, subtracted	Uni-ZAP XR	LP04
HEPA HEPB HEPC	Human Epididymus	Uni-ZAP XR	LP04
HSNA HSNB HSNB HSNM HSNM	Human Synovium	Uni-ZAP XR	LP04
HPFB HPFC HPFD HPFE	Human Prostate Cancer, Stage C fraction	Uni-ZAP XR	LP04
HE2A HE2D HE2E HE2H HE2I HE2M HE2N HE2O	12 Week Old Early Stage Human	Uni-ZAP XR	LP04
HE2B HE2C HE2F HE2G HE2P HE2Q	12 Week Old Early Stage Human, II	Uni-ZAP XR	LP04
HPTS HPTT HPTU	Human Pituitary, subtracted	Uni-ZAP XR	LP04
HAUA HAUB HAUC	Amniotic Cells - TNF induced	Uni-ZAP XR	LP04
HAQA HAQB HAQC HAQD	Amniotic Cells - Primary Culture	Uni-ZAP XR	LP04
HWTA HWTB HWTG	wilm's tumor	Uni-ZAP XR	LP04
HBSD	Bone Cancer, re-excision	Uni-ZAP XR	LP04
HSGB	Salivary gland, re-excision	Uni-ZAP XR	LP04
HSJA HSJB HSJC	Smooth muscle-ILb induced	Uni-ZAP XR	LP04
HSXA HSXB HSXC HSXD	Human Substantia Nigra	Uni-ZAP XR	LP04
HSJA HSJB HSJC	Smooth muscle, IL1b induced	Uni-ZAP XR	LP04
HOUA HOUB HOUC HOUD HOUE	Adipocytes	Uni-ZAP XR	LP04
HPWA HPWB HPWC HPWD HPWE	Prostate BPH	Uni-ZAP XR	LP04
HELA HELB HELC HELD HELE HELF HELG HELH	Endothelial cells-control	Uni-ZAP XR	LP04
HEMA HEMB HEMC HEMD HEME HEMF HEMG HEMH	Endothelial-induced	Uni-ZAP XR	LP04
HBIA HBIB HBIC	Human Brain, Striatum	Uni-ZAP XR	LP04
HHSA HHSB HHSC HHSD HHSE	Human Hypothalamus, Schizophrenia	Uni-ZAP XR	LP04
HNGA HNGB HNGC HNGD HNGE HNGF HNGG HNGH HNGI HNGJ	neutrophils control	Uni-ZAP XR	LP04
HNHA HNHB HNHC HNHD HNHE HNHF HNHG HNHH HNHI HNHI	Neutrophils IL-1 and LPS induced	Uni-ZAP XR	LP04
HSDB HSDC	STRIATUM DEPRESSION	Uni-ZAP XR	LP04
HHPT	Hypothalamus	Uni-ZAP XR	LP04
HSAT HSAU HSAV HSAW HSAX HSAY HSAZ	Anergic T-cell	Uni-ZAP XR	LP04
HBMS HBMT HBMU HBMV HBMW HBMX	Bone marrow	Uni-ZAP XR	LP04
HOEA HOEB HOEC HOED HOEE HOEF HOEJ	Osteoblasts	Uni-ZAP XR	LP04
HAIA HAIB HAIC HAID HAIE HAIF	Epithelial-TNFa and INF induced	Uni-ZAP XR	LP04

Libraries owned by Catalog	Catalog Description	Vector	ATCC Deposit
HTGA HTGB HTGC HTGD	Apoptotic T-cell	Uni-ZAP XR	LP04
HMCA HMCB HMCC HMCD HMCE	Macrophage-oxLDL	Uni-ZAP XR	LP04
HMAA HMAB HMAC HMAD HMAE HMAF HMAG	Macrophage (GM-CSF treated)	Uni-ZAP XR	LP04
HPHA	Normal Prostate	Uni-ZAP XR	LP04
HPIA HPIB HPIC	LNCAP prostate cell line	Uni-ZAP XR	LP04
HPJA HPJB HPJC	PC3 Prostate cell line	Uni-ZAP XR	LP04
HOSE HOSF HOSG	Human Osteoclastoma, re-excision	Uni-ZAP XR	LP04
HTGE HTGF	Apoptotic T-cell, re-excision	Uni-ZAP XR	LP04
HMAJ HMAK	H Macrophage (GM-CSF treated), re-excision	Uni-ZAP XR	LP04
HACB HACC HACD	Human Adipose Tissue, re-excision	Uni-ZAP XR	LP04
HFPA	H. Frontal Cortex, Epileptic	Uni-ZAP XR	LP04
HFAA HFAB HFAC HFAD HFAE	Alzheimer's, spongy change	Uni-ZAP XR	LP04
HFAM	Frontal Lobe, Dementia	Uni-ZAP XR	LP04
HMIA HMIB HMIC	Human Manic Depression Tissue	Uni-ZAP XR	LP04
HTSA HTSE HTSF HTSG HTSH	Human Thymus	pBS	LP05
HPBA HPBB HPBC HPBD HPBE	Human Pineal Gland	pBS	LP05
HSAA HSAB HSAC	HSA 172 Cells	pBS	LP05
HSBA HSBB HSBC HSBM	HSC172 cells	pBS	LP05
HJAA HJAB HJAC HJAD	Jurkat T-cell G1 phase	pBS	LP05
HJBA HJBB HJBC HJBD	Jurkat T-Cell, S phase	pBS	LP05
HAFA HAFB	Aorta endothelial cells + TNF-a	pBS	LP05
HAWA HAWB HAWC	Human White Adipose	pBS	LP05
HTNA HTNB	Human Thyroid	pBS	LP05
HONA	Normal Ovary, Premenopausal	pBS	LP05
HARA HARB	Human Adult Retina	pBS	LP05
HLJA HLJB	Human Lung	pCMVSPORT 1	LP06
HOFM HOFN HOFO	H. Ovarian Tumor, II, OV5232	pCMVSPORT 2.0	LP07
HOGA HOGB HOGC	OV 10-3-95	pCMVSPORT 2.0	LP07
HCGL	CD34+cells, II	pCMVSPORT 2.0	LP07
HDLA	Hodgkin's Lymphoma I	pCMVSPORT 2.0	LP07
HDTA HDTB HDTC HDTD HDTE	Hodgkin's Lymphoma II	pCMVSPORT 2.0	LP07
HKAA HKAB HKAC HKAD HKAE HKAF HKAG HKAH	Keratinocyte	pCMVSPORT2.0	LP07
HCIM	CAPFINDER, Crohn's Disease, lib 2	pCMVSPORT 2.0	LP07
HKAL	Keratinocyte, lib 2	pCMVSPORT2.0	LP07
HKAT	Keratinocyte, lib 3	pCMVSPORT2.0	LP07
HNDA	Nasal polyps	pCMVSPORT2.0	LP07
HDRA	H. Primary Dendritic Cells, lib 3	pCMVSPORT2.0	LP07

Libraries owned by Catalog	Catalog Description	Vector	ATCC Deposit
HOHA HOHB HOHC	Human Osteoblasts II	pCMVSPORT2.0	LP07
HLDA HLDB HLDC	Liver, Hepatoma	pCMVSPORT3.0	LP08
HLDN HLDO HLDP	Human Liver, normal	pCMVSPORT3.0	LP08
HMTA	pBMC stimulated w/ poly I/C	pCMVSPORT3.0	LP08
HNTA	NTERA2, control	pCMVSPORT3.0	LP08
HDP A HDPB HDP C HDPD HDPF HDPG HDPH HDPI HDPJ HDPK	Primary Dendritic Cells, lib 1	pCMVSPORT3.0	LP08
HDPM HDPN HDPO HDPP	Primary Dendritic cells, frac 2	pCMVSPORT3.0	LP08
HMUA HMUB HMUC	Myeloid Progenitor Cell Line	pCMVSPORT3.0	LP08
HHEA HHEB HHEC HHED	T Cell helper I	pCMVSPORT3.0	LP08
HHEM HHEN HHEO HHEP	T cell helper II	pCMVSPORT3.0	LP08
HEQA HEQB HEQC	Human endometrial stromal cells	pCMVSPORT3.0	LP08
HJMA HJMB	Human endometrial stromal cells- treated with progesterone	pCMVSPORT3.0	LP08
HSWA HSWB HSWC	Human endometrial stromal cells- treated with estradiol	pCMVSPORT3.0	LP08
HSYA HSYB HSYC	Human Thymus Stromal Cells	pCMVSPORT3.0	LP08
HLWA HLWB HLWC	Human Placenta	pCMVSPORT3.0	LP08
HRAA HRAB HRAC	Rejected Kidney, lib 4	pCMVSPORT3.0	LP08
HMTM	PCR, pBMC I/C treated	PCR II	LP09
HMJA	H. Meningioma, M6	pSport 1	LP10
HMKA HMKB HMKC HMKD HMKE	H. Meningioma, M1	pSport 1	LP10
HUSG HUSI	Human umbilical vein endothelial cells, IL-4 induced	pSport 1	LP10
HUSX HUSY	Human Umbilical Vein Endothelial Cells, uninduced	pSport 1	LP10
HOFA	Ovarian Tumor I, OV5232	pSport 1	LP10
HCFA HCFB HCFC HCFD	T-Cell PHA 16 hrs	pSport 1	LP10
HCFL HCFM HCFN HCFO	T-Cell PHA 24 hrs	pSport 1	LP10
HADA HADC HADD HADE HADF HADG	Human Adipose	pSport 1	LP10
HOVA HOVB HOVC	Human Ovary	pSport 1	LP10
HTWB HTWC HTWD HTWE HTWF	Resting T-Cell Library, II	pSport 1	LP10
HMMA	Spleen metastatic melanoma	pSport 1	LP10
HLYA HLYB HLYC HLYD HLYE	Spleen, Chronic lymphocytic leukemia	pSport 1	LP10
HCGA	CD34+ cell, I	pSport 1	LP10
HEOM HEON	Human Eosinophils	pSport 1	LP10
HTDA	Human Tonsil, Lib 3	pSport 1	LP10
HSPA	Salivary Gland, Lib 2	pSport 1	LP10
HCHA HCHB HCHC	Breast Cancer cell line, MDA 36	pSport 1	LP10

Libraries owned by Catalog	Catalog Description	Vector	ATCC Deposit
HCHM HCHN	Breast Cancer Cell line, angiogenic	pSport 1	LP10
HCIA	Crohn's Disease	pSport 1	LP10
HDAA HDAB HDAC	HEL cell line	pSport 1	LP10
HABA	Human Astrocyte	pSport 1	LP10
HUFA HUFB HUFC	Ulcerative Colitis	pSport 1	LP10
HNTM	NTERA2 + retinoic acid, 14 days	pSport 1	LP10
HDQA	Primary Dendritic cells,CapFinder2, frac 1	pSport 1	LP10
HDQM	Primary Dendritic Cells, CapFinder, frac 2	pSport 1	LP10
HLDX	Human Liver, normal,CapFinder	pSport 1	LP10
HULA HULB HULC	Human Dermal Endothelial Cells,untreated	pSport1	LP10
HUMA	Human Dermal Endothelial cells,treated	pSport1	LP10
HCJA	Human Stromal Endometrial fibroblasts, untreated	pSport1	LP10
HCJM	Human Stromal endometrial fibroblasts, treated w/ estradiol	pSport1	LP10
HEDA	Human Stromal endometrial fibroblasts, treated with progesterone	pSport1	LP10
HFNA	Human ovary tumor cell OV350721	pSport1	LP10
HKGA HKGB HKGC HKGD	Merkel Cells	pSport1	LP10
HISA HISB HISC	Pancreas Islet Cell Tumor	pSport1	LP10
HLSA	Skin, burned	pSport1	LP10
HBZA	Prostate,BPH, Lib 2	pSport 1	LP10
HBZS	Prostate BPH,Lib 2, subtracted	pSport 1	LP10
HFIA HFIB HFIC	Synovial Fibroblasts (control)	pSport 1	LP10
HFIH HFII HFUJ	Synovial hypoxia	pSport 1	LP10
HFIT HFIU HFIV	Synovial IL-1/TNF stimulated	pSport 1	LP10
HGCA	Messangial cell, frac 1	pSport1	LP10
HMVA HMOV HMVC	Bone Marrow Stromal Cell, untreated	pSport1	LP10
HFIX HFIY HFIZ	Synovial Fibroblasts (Il1/TNF), subt	pSport1	LP10
HFOX HFOY HFOZ	Synovial hypoxia-RSF subtracted	pSport1	LP10
HMQA HMQB HMQC HMQD	Human Activated Monocytes	Uni-ZAP XR	LP11
HLIA HLIB HLIC	Human Liver	pCMVSPORT 1	LP012
HHBA HHBB HHBC HHBD HHBE	Human Heart	pCMVSPORT 1	LP012
HBBA HBBB	Human Brain	pCMVSPORT 1	LP012
HLJA HLJB HLJC HLJD HLJE	Human Lung	pCMVSPORT 1	LP012
HOGA HOGB HOGC	Ovarian Tumor	pCMVSPORT 2.0	LP012
HTJM	Human Tonsils, Lib 2	pCMVSPORT 2.0	LP012

Libraries owned by Catalog	Catalog Description	Vector	ATCC Deposit
HAMF HAMG	KMH2	pCMVSPORT 3.0	LP012
HAJA HAJB HAJC	L428	pCMVSPORT 3.0	LP012
HWBA HWBB HWBC HWBD HWBE	Dendritic cells, pooled	pCMVSPORT 3.0	LP012
HWAA HWAB HWAC HWAD HWAE	Human Bone Marrow, treated	pCMVSPORT 3.0	LP012
HYAA HYAB HYAC	B Cell lymphoma	pCMVSPORT 3.0	LP012
HWHG HWHH HWHI	Healing groin wound, 6.5 hours post incision	pCMVSPORT 3.0	LP012
HWHP HWHQ HWHR	Healing groin wound; 7.5 hours post incision	pCMVSPORT 3.0	LP012
HARM	Healing groin wound - zero hr post-incision (control)	pCMVSPORT 3.0	LP012
HBIM	Olfactory epithelium; nasalcavity	pCMVSPORT 3.0	LP012
HWDA	Healing Abdomen wound; 70&90 min post incision	pCMVSPORT 3.0	LP012
HWEA	Healing Abdomen Wound;15 days post incision	pCMVSPORT 3.0	LP012
HWJA	Healing Abdomen Wound;21&29 days	pCMVSPORT 3.0	LP012
HNAL	Human Tongue, frac 2	pSport1	LP012
HMJA	H. Meningioma, M6	pSport1	LP012
HMKA HMKB HMKC HMKD HMKE	H. Meningioma, M1	pSport1	LP012
HOFA	Ovarian Tumor I, OV5232	pSport1	LP012
HCFA HCFB HCFC HCFD	T-Cell PHA 16 hrs	pSport1	LP012
HCFL HCFM HCFN HCFO	T-Cell PHA 24 hrs	pSport1	LP012
HMMA HMMB HMMC	Spleen metastatic melanoma	pSport1	LP012
HTDA	Human Tonsil, Lib 3	pSport1	LP012
HDBA	Human Fetal Thymus	pSport1	LP012
HDBA	Pericardium	pSport1	LP012
HBZA	Prostate,BPH, Lib 2	pSport1	LP012
HWCA	Larynx tumor	pSport1	LP012
HWKA	Normal lung	pSport1	LP012
HSMB	Bone marrow stroma,treated	pSport1	LP012
HBHM	Normal trachea	pSport1	LP012
HLFC	Human Larynx	pSport1	LP012
HLRB	Siebben Polyposis	pSport1	LP012
HNIA	Mammary Gland	pSport1	LP012
HNJB	Palate carcinoma	pSport1	LP012
HNKA	Palate normal	pSport1	LP012
HMZA	Pharynx carcinoma	pSport1	LP012
HABG	Cheek Carcinoma	pSport1	LP012
HMZM	Pharynx Carcinoma	pSport1	LP012

Libraries owned by Catalog	Catalog Description	Vector	ATCC Deposit
HDRM	Larynx Carcinoma	pSport1	LP012
HVAA	Pancreas normal PCA4 No	pSport1	LP012
HICA	Tongue carcinoma	pSport1	LP012
HUKA HUKB HUKC HUKD HUKF	Human Uterine Cancer	Lambda ZAP II	LP013
HFFA	Human Fetal Brain, random primed	Lambda ZAP II	LP013
HTUA	Activated T-cell labeled with 4-thioluri	Lambda ZAP II	LP013
HBQA	Early Stage Human Brain, random primed	Lambda ZAP II	LP013
HMEB	Human microvascular Endothelial cells, fract. B	Lambda ZAP II	LP013
HUSH	Human Umbilical Vein Endothelial cells, fract. A, re-excision	Lambda ZAP II	LP013
HLQC HLQD	Hepatocellular tumor, re-excision	Lambda ZAP II	LP013
HTWJ HTWK HTWL	Resting T-cell, re-excision	Lambda ZAP II	LP013
HF6S	Human Whole 6 week Old Embryo (II), subt	pBluescript	LP013
HHPS	Human Hippocampus, subtracted	pBluescript	LP013
HL1S	LNCAP, differential expression	pBluescript	LP013
HLHS HLHT	Early Stage Human Lung, Subtracted	pBluescript	LP013
HSUS	Supt cells, cyclohexamide treated, subtracted	pBluescript	LP013
HSUT	Supt cells, cyclohexamide treated, differentially expressed	pBluescript	LP013
HSDS	H. Striatum Depression, subtracted	pBluescript	LP013
HPTZ	Human Pituitary, Subtracted VII	pBluescript	LP013
HSDX	H. Striatum Depression, subt II	pBluescript	LP013
HSDZ	H. Striatum Depression, subt	pBluescript	LP013
HPBA HPBB HPBC HPBD HPBE	Human Pineal Gland	pBluescript SK-	LP013
HRTA	Colorectal Tumor	pBluescript SK-	LP013
HSBA HSBB HSBC HSBM	HSC172 cells	pBluescript SK-	LP013
HJAA HJAB HJAC HJAD	Jurkat T-cell G1 phase	pBluescript SK-	LP013
HJBA HJBB HJBC HJBD	Jurkat T-cell, S1 phase	pBluescript SK-	LP013
HTNA HTNB	Human Thyroid	pBluescript SK-	LP013
HAHA HAHB	Human Adult Heart	Uni-ZAP XR	LP013
HE6A	Whole 6 week Old Embryo	Uni-ZAP XR	LP013
HFCA HFCB HFCC HFCD HFCE	Human Fetal Brain	Uni-ZAP XR	LP013
HFKC HFKD HFKE HFKF HFKG	Human Fetal Kidney	Uni-ZAP XR	LP013
HGBA HGBD HGBE HGBF HGBG	Human Gall Bladder	Uni-ZAP XR	LP013
HPRA HPRB HPRC HPRD	Human Prostate	Uni-ZAP XR	LP013
HTEA HTEB HTEC HTED HTEE	Human Testes	Uni-ZAP XR	LP013

Libraries owned by Catalog	Catalog Description	Vector	ATCC Deposit
HTTA HTTB HTTC HTTD HTTE	Human Testes Tumor	Uni-ZAP XR	LP013
HYBA HYBB	Human Fetal Bone	Uni-ZAP XR	LP013
HFLA	Human Fetal Liver	Uni-ZAP XR	LP013
HHFB HHFC HHFD HHFE HHFF	Human Fetal Heart	Uni-ZAP XR	LP013
HUVB HUV C HUVD HUVE	Human Umbilical Vein, End. remake	Uni-ZAP XR	LP013
HTHB HTHC HTHD	Human Thymus	Uni-ZAP XR	LP013
HSTA HSTB HSTC HSTD	Human Skin Tumor	Uni-ZAP XR	LP013
HTAA HTAB HTAC HTAD HTAE	Human Activated T-cells	Uni-ZAP XR	LP013
HFEA HFEB HFEC	Human Fetal Epithelium (skin)	Uni-ZAP XR	LP013
HJPA HJPB HJPC HJPD	Human Jurkat Membrane Bound Polysomes	Uni-ZAP XR	LP013
HESA	Human Epithelioid Sarcoma	Uni-ZAP XR	LP013
HALS	Human Adult Liver, Subtracted	Uni-ZAP XR	LP013
HFTA HFTB HFTC HFTD	Human Fetal Dura Mater	Uni-ZAP XR	LP013
HCAA HCAB HCAC	Cem cells, cyclohexamide treated	Uni-ZAP XR	LP013
HRGA HRGB HRGC HRGD	Raji Cells, cyclohexamide treated	Uni-ZAP XR	LP013
HE9A HE9B HE9C HE9D HE9E	Nine Week Old Early Stage Human	Uni-ZAP XR	LP013
HSFA	Human Fibrosarcoma	Uni-ZAP XR	LP013
HATA HATB HATC HATD HATE	Human Adrenal Gland Tumor	Uni-ZAP XR	LP013
HTRA	Human Trachea Tumor	Uni-ZAP XR	LP013
HE2A HE2D HE2E HE2H HE2I	12 Week Old Early Stage Human	Uni-ZAP XR	LP013
HE2B HE2C HE2F HE2G HE2P	12 Week Old Early Stage Human, II	Uni-ZAP XR	LP013
HNEA HNEB HNEC HNED HNEE	Human Neutrophil	Uni-ZAP XR	LP013
HBGA	Human Primary Breast Cancer	Uni-ZAP XR	LP013
HPTS HPTT HPTU	Human Pituitary, subtracted	Uni-ZAP XR	LP013
HMQA HMQB HMQC HMQD	Human Activated Monocytes	Uni-ZAP XR	LP013
HOAA HOAB HOAC	Human Osteosarcoma	Uni-ZAP XR	LP013
HTOA HTOD HTOE HTOF HTOG	human tonsils	Uni-ZAP XR	LP013
HMGB	Human OB MG63 control fraction I	Uni-ZAP XR	LP013
HOPB	Human OB HOS control fraction I	Uni-ZAP XR	LP013
HOQB	Human OB HOS treated (1 nM E2) fraction I	Uni-ZAP XR	LP013
HAUA HAUB HAUC	Amniotic Cells - TNF induced	Uni-ZAP XR	LP013
HAQA HAQB HAQC HAQD	Amniotic Cells - Primary Culture	Uni-ZAP XR	LP013
HROA HROC	HUMAN STOMACH	Uni-ZAP XR	LP013
HBJA HBJB HBJC HBJD HBJE	HUMAN B CELL LYMPHOMA	Uni-ZAP XR	LP013
HODA HODB HODC HODD	human ovarian cancer	Uni-ZAP XR	LP013
HCPA	Corpus Callosum	Uni-ZAP XR	LP013
HSOA	stomach cancer (human)	Uni-ZAP XR	LP013
HERA	SKIN	Uni-ZAP XR	LP013

09950087 091201

Libraries owned by Catalog	Catalog Description	Vector	ATCC Deposit
HMDA	Brain-medulloblastoma	Uni-ZAP XR	LP013
HGLA HGLB HGLD	Glioblastoma	Uni-ZAP XR	LP013
HWTA HWTB HWTC	wilm's tumor	Uni-ZAP XR	LP013
HEAA	H. Atrophic Endometrium	Uni-ZAP XR	LP013
HAPN HAPO HAPQ HAPR	Human Adult Pulmonary;re-excision	Uni-ZAP XR	LP013
HLTG HLTH	Human T-cell lymphoma;re-excision	Uni-ZAP XR	LP013
HAHC HAHD HAHE	Human Adult Heart;re-excision	Uni-ZAP XR	LP013
HAGA HAGB HAGC HAGD HAGE	Human Amygdala	Uni-ZAP XR	LP013
HSJA HSJB HSJC	Smooth muscle-ILb induced	Uni-ZAP XR	LP013
HSJA HSJB HSJC	Smooth muscle, IL1b induced	Uni-ZAP XR	LP013
HPWA HPWB HPWC HPWD HPWE	Prostate BPH	Uni-ZAP XR	LP013
HPJA HPJB HPJC	PC3 Prostate cell line	Uni-ZAP XR	LP013
HPJA HPJB HPJC	PC3 Prostate cell line	Uni-ZAP XR	LP013
HBTB	Bone Marrow Stroma, TNF&LPS ind	Uni-ZAP XR	LP013
HMCB HMCB HMCB HMCB HMCB	Macrophage-oxLDL; re-excision	Uni-ZAP XR	LP013
HAGG HAGH HAGI	Human Amygdala;re-excision	Uni-ZAP XR	LP013
HACA	H. Adipose Tissue	Uni-ZAP XR	LP013
HKFB	K562 + PMA (36 hrs),re-excision	ZAP Express	LP013
HCWT HCWU HCWV	CD34 positive cells (cord blood),re-ex	ZAP Express	LP013
HBWA	Whole brain	ZAP Express	LP013
HBXA HBXB HBXC HBXD	Human Whole Brain #2 - Oligo dT > 1.5Kb	ZAP Express	LP013
HAVM	Temporal cortex-Alzheimer	pT-Adv	LP014
HAVT	Hippocampus, Alzheimer Subtracted	pT-Adv	LP014
HHAS	CHME Cell Line	Uni-ZAP XR	LP014
HAJR	Larynx normal	pSport 1	LP014
HWLE HWLF HWLG HWLH	Colon Normal	pSport 1	LP014
HCRM HCRN HCRO	Colon Carcinoma	pSport 1	LP014
HWLI HWLJ HWLK	Colon Normal	pSport 1	LP014
HWLQ HWLR HWLS HWLT	Colon Tumor	pSport 1	LP014
HBFM	Gastrocnemius Muscle	pSport 1	LP014
HBOD HBOE	Quadriceps Muscle	pSport 1	LP014
HBKD HBKE	Soleus Muscle	pSport 1	LP014
HCCM	Pancreatic Langerhans	pSport 1	LP014
HWGA	Larynx carcinoma	pSport 1	LP014
HWGM HWGN	Larynx carcinoma	pSport 1	LP014
HWLA HWLB HWLC	Normal colon	pSport 1	LP014
HWLM HWLN	Colon Tumor	pSport 1	LP014
HVAM HVAN HVAO	Pancreas Tumor	pSport 1	LP014
HWGQ	Larynx carcinoma	pSport 1	LP014

Libraries owned by Catalog	Catalog Description	Vector	ATCC Deposit
HAQM HAQN	Salivary Gland	pSport 1	LP014
HASM	Stomach; normal	pSport 1	LP014
HBCM	Uterus; normal	pSport 1	LP014
HCDM	Testis; normal	pSport 1	LP014
HDJM	Brain; normal	pSport 1	LP014
HEFM	Adrenal Gland, normal	pSport 1	LP014
HBAA	Rectum normal	pSport 1	LP014
HFDM	Rectum tumour	pSport 1	LP014
HGAM	Colon, normal	pSport 1	LP014
HHMM	Colon, tumour	pSport 1	LP014
HCLB HCLC	Human Lung Cancer	Lambda Zap II	LP015
HRLA	L1 Cell line	ZAP Express	LP015
HHAM	Hypothalamus, Alzheimer's	pCMVSPORT 3.0	LP015
HKBA	Ku 812F Basophils Line	pSport 1	LP015
HS2S	Saos2, Dexamethosone Treated	pSport 1	LP016
HA5A	Lung Carcinoma A549 TNFalpha activated	pSport 1	LP016
HTFM	TF-1 Cell Line GM-CSF Treated	pSport 1	LP016
HYAS	Thyroid Tumour	pSport 1	LP016
HUTS	Larynx Normal	pSport 1	LP016
HXOA	Larynx Tumor	pSport 1	LP016
HEAH	Ea.hy.926 cell line	pSport 1	LP016
HINA	Adenocarcinoma Human	pSport 1	LP016
HRMA	Lung Mesothelium	pSport 1	LP016
HLCL	Human Pre-Differentiated Adipocytes	Uni-Zap XR	LP017
HS2A	Saos2 Cells	pSport 1	LP020
HS2I	Saos2 Cells; Vitamin D3 Treated	pSport 1	LP020
HUCM	CHME Cell Line, untreated	pSport 1	LP020
HEPN	Aryepiglottis Normal	pSport 1	LP020
HPSN	Sinus Piniformis Tumour	pSport 1	LP020
HNSA	Stomach Normal	pSport 1	LP020
HNSM	Stomach Tumour	pSport 1	LP020
HNLA	Liver Normal Met5No	pSport 1	LP020
HUTA	Liver Tumour Met 5 Tu	pSport 1	LP020
HOCN	Colon Normal	pSport 1	LP020
HOCT	Colon Tumor	pSport 1	LP020
HTNT	Tongue Tumour	pSport 1	LP020
HLXN	Larynx Normal	pSport 1	LP020
HLXT	Larynx Tumour	pSport 1	LP020
HTYN	Thymus	pSport 1	LP020

Libraries owned by Catalog	Catalog Description	Vector	ATCC Deposit
HPLN	Placenta	pSport 1	LP020
HTNG	Tongue Normal	pSport 1	LP020
HZAA	Thyroid Normal (SDCA2 No)	pSport 1	LP020
HWES	Thyroid Thyroiditis	pSport 1	LP020
HFHD	Ficolled Human Stromal Cells, 5Fu treated	pTrip1Ex2	LP021
HFHM,HFHN	Ficolled Human Stromal Cells, Untreated	pTrip1Ex2	LP021
HPCI	Hep G2 Cells, lambda library	lambda Zap-CMV XR	LP021
HBCA,HBCB,HBCC	H. Lymph node breast Cancer	Uni-ZAP XR	LP021
HCOK	Chondrocytes	pSPORT1	LP022
HDCA, HDCB, HDCC	Dendritic Cells From CD34 Cells	pSPORT1	LP022
HDMA, HDMB	CD40 activated monocyte dendritic cells	pSPORT1	LP022
HDDM, HDDN, HDDO	LPS activated derived dendritic cells	pSPORT1	LP022
HPCR	Hep G2 Cells, PCR library	lambda Zap-CMV XR	LP022
HAAA, HAAB, HAAC	Lung, Cancer (4005313A3): Invasive Poorly Differentiated Lung Adenocarcinoma	pSPORT1	LP022
HIPA, HIPB, HIPC	Lung, Cancer (4005163 B7): Invasive, Poorly Diff. Adenocarcinoma, Metastatic	pSPORT1	LP022
HOOH, HOOI	Ovary, Cancer: (4004562 B6) Papillary Serous Cystic Neoplasm, Low Malignant Pot	pSPORT1	LP022
HIDA	Lung, Normal: (4005313 B1)	pSPORT1	LP022
HUJA,HUJB,HUJC,HUJD,HUJE	B-Cells	pCMVSPORT 3.0	LP022
HNOA,HNOB,HNOC,HNOD	Ovary, Normal: (9805C040R)	pSPORT1	LP022
HNLM	Lung, Normal: (4005313 B1)	pSPORT1	LP022
HSCL	Stromal Cells	pSPORT1	LP022
HAAX	Lung, Cancer: (4005313 A3) Invasive Poorly-differentiated Metastatic lung adenocarcinoma	pSPORT1	LP022
HUUA,HUUB,HUUC,HUUD	B-cells (unstimulated)	pTrip1Ex2	LP022
HWWA,HWWB,HWWC,HWWD,HWWE,HWWF,HWWG	B-cells (stimulated)	pSPORT1	LP022
HCCC	Colon, Cancer: (9808C064R)	pCMVSPORT 3.0	LP023
HPDO HPDP HPDQ HPDR HPD	Ovary, Cancer (9809C332): Poorly differentiated adenocarcinoma	pSport 1	LP023
HPCO HPCP HPCQ HPCT	Ovary, Cancer (15395A1F): Grade II	pSport 1	LP023

Libraries owned by Catalog	Catalog Description	Vector	ATCC Deposit
	Papillary Carcinoma		
HOCM HOCO HOCF HOCQ	Ovary, Cancer: (15799A1F) Poorly differentiated carcinoma	pSport 1	LP023
HCBM HCBN HCBO	Breast, Cancer: (4004943 A5)	pSport 1	LP023
HNBT HNBU HNBV	Breast, Normal: (4005522B2)	pSport 1	LP023
HBCP HBCQ	Breast, Cancer: (4005522 A2)	pSport 1	LP023
HBCJ	Breast, Cancer: (9806C012R)	pSport 1	LP023
HSAM HSAN	Stromal cells 3.88	pSport 1	LP023
HVCA HVCB HVCC HVCD	Ovary, Cancer: (4004332 A2)	pSport 1	LP023
HSCK HSEN HSEO	Stromal cells (HBM3.18)	pSport 1	LP023
HSCP HSCQ	stromal cell clone 2.5	pSport 1	LP023
HUXA	Breast Cancer: (4005385 A2)	pSport 1	LP023
HCOM HCON HCOO HCOP HCOQ	Ovary, Cancer (4004650 A3): Well-Differentiated Micropapillary Serous Carcinoma	pSport 1	LP023
HBNM	Breast, Cancer: (9802C020E)	pSport 1	LP023
HVVA HVVB HVVC HVVD HVVE	Human Bone Marrow, treated	pSport 1	LP023

[931] Two nonlimiting examples are provided below for isolating a particular clone from the deposited sample of plasmid cDNAs cited for that clone in Table 7. First, a plasmid is directly isolated by screening the clones using a polynucleotide probe corresponding to the nucleotide sequence of SEQ ID NO:X.

[932] Particularly, a specific polynucleotide with 30-40 nucleotides is synthesized using an Applied Biosystems DNA synthesizer according to the sequence reported. The oligonucleotide is labeled, for instance, with ^{32}P - γ -ATP using T4 polynucleotide kinase and purified according to routine methods. (E.g., Maniatis et al., Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Press, Cold Spring, NY (1982)). The plasmid mixture is transformed into a suitable host, as indicated above (such as XL-1 Blue (Stratagene)) using techniques known to those of skill in the art, such as those provided by the vector supplier or in related publications or patents cited above. The transformants are plated on 1.5% agar plates (containing the appropriate selection agent, e.g., ampicillin) to a density of about 150 transformants (colonies) per plate. These plates are screened using Nylon membranes according to routine methods for bacterial colony screening (e.g., Sambrook et al., Molecular Cloning: A Laboratory Manual, 2nd Edit., (1989), Cold Spring

Harbor Laboratory Press, pages 1.93 to 1.104), or other techniques known to those of skill in the art.

[933] Alternatively, two primers of 17-20 nucleotides derived from both ends of the nucleotide sequence of SEQ ID NO:X are synthesized and used to amplify the desired cDNA using the deposited cDNA plasmid as a template. The polymerase chain reaction is carried out under routine conditions, for instance, in 25 μ l of reaction mixture with 0.5 ug of the above cDNA template. A convenient reaction mixture is 1.5-5 mM $MgCl_2$, 0.01% (w/v) gelatin, 20 μ M each of dATP, dCTP, dGTP, dTTP, 25 pmol of each primer and 0.25 Unit of Taq polymerase. Thirty five cycles of PCR (denaturation at 94°C for 1 min; annealing at 55°C for 1 min; elongation at 72°C for 1 min) are performed with a Perkin-Elmer Cetus automated thermal cycler. The amplified product is analyzed by agarose gel electrophoresis and the DNA band with expected molecular weight is excised and purified. The PCR product is verified to be the selected sequence by subcloning and sequencing the DNA product.

[934] Several methods are available for the identification of the 5' or 3' non-coding portions of a gene which may not be present in the deposited clone. These methods include but are not limited to, filter probing, clone enrichment using specific probes, and protocols similar or identical to 5' and 3' "RACE" protocols which are well known in the art. For instance, a method similar to 5' RACE is available for generating the missing 5' end of a desired full-length transcript. (Fromont-Racine et al., *Nucleic Acids Res.* 21(7):1683-1684 (1993)).

[935] Briefly, a specific RNA oligonucleotide is ligated to the 5' ends of a population of RNA presumably containing full-length gene RNA transcripts. A primer set containing a primer specific to the ligated RNA oligonucleotide and a primer specific to a known sequence of the gene of interest is used to PCR amplify the 5' portion of the desired full-length gene. This amplified product may then be sequenced and used to generate the full length gene.

[936] This above method starts with total RNA isolated from the desired source, although poly-A+ RNA can be used. The RNA preparation can then be treated with phosphatase if necessary to eliminate 5' phosphate groups on degraded or damaged RNA which may interfere with the later RNA ligase step. The phosphatase should then be inactivated and the RNA treated with tobacco acid pyrophosphatase in order to remove the cap structure present at the 5' ends of messenger RNAs. This reaction leaves a 5' phosphate

group at the 5' end of the cap cleaved RNA which can then be ligated to an RNA oligonucleotide using T4 RNA ligase.

[937] This modified RNA preparation is used as a template for first strand cDNA synthesis using a gene specific oligonucleotide. The first strand synthesis reaction is used as a template for PCR amplification of the desired 5' end using a primer specific to the ligated RNA oligonucleotide and a primer specific to the known sequence of the gene of interest. The resultant product is then sequenced and analyzed to confirm that the 5' end sequence belongs to the desired gene.

Example 2: Isolation of Genomic Clones Corresponding to a Polynucleotide

[938] A human genomic P1 library (Genomic Systems, Inc.) is screened by PCR using primers selected for the sequence corresponding to SEQ ID NO:X according to the method described in Example 1. (See also, Sambrook.)

Example 3: Tissue specific expression analysis

[939] The Human Genome Sciences, Inc. (HGS) database is derived from sequencing tissue and/or disease specific cDNA libraries. Libraries generated from a particular tissue are selected and the specific tissue expression pattern of EST groups or assembled contigs within these libraries is determined by comparison of the expression patterns of those groups or contigs within the entire database. ESTs and assembled contigs which show tissue specific expression are selected.

[940] The original clone from which the specific EST sequence was generated, or in the case of an assembled contig, the clone from which the 5' most EST sequence was generated, is obtained from the catalogued library of clones and the insert amplified by PCR using methods known in the art. The PCR product is denatured and then transferred in 96 or 384 well format to a nylon membrane (Schleicher and Scheull) generating an array filter of tissue specific clones. Housekeeping genes, maize genes, and known tissue specific genes are included on the filters. These targets can be used in signal normalization and to validate assay sensitivity. Additional targets are included to monitor probe length and specificity of hybridization.

[941] Radioactively labeled hybridization probes are generated by first strand cDNA synthesis per the manufacturer's instructions (Life Technologies) from mRNA/RNA samples prepared from the specific tissue being analyzed (e.g., prostate, prostate cancer, ovarian, ovarian cancer, etc.). The hybridization probes are purified by gel exclusion chromatography, quantitated, and hybridized with the array filters in hybridization bottles at 65°C overnight. The filters are washed under stringent conditions and signals are captured using a Fuji phosphorimager.

[942] Data is extracted using AIS software and following background subtraction, signal normalization is performed. This includes a normalization of filter-wide expression levels between different experimental runs. Genes that are differentially expressed in the tissue of interest are identified.

Example 4: Chromosomal Mapping of the Polynucleotides

[943] An oligonucleotide primer set is designed according to the sequence at the 5' end of SEQ ID NO:X. This primer preferably spans about 100 nucleotides. This primer set is then used in a polymerase chain reaction under the following set of conditions: 30 seconds, 95°C; 1 minute, 56°C; 1 minute, 70°C. This cycle is repeated 32 times followed by one 5 minute cycle at 70°C. Human, mouse, and hamster DNA is used as template in addition to a somatic cell hybrid panel containing individual chromosomes or chromosome fragments (Bios, Inc). The reactions are analyzed on either 8% polyacrylamide gels or 3.5 % agarose gels. Chromosome mapping is determined by the presence of an approximately 100 bp PCR fragment in the particular somatic cell hybrid.

Example 5: Bacterial Expression of a Polypeptide

[944] A polynucleotide encoding a polypeptide of the present invention is amplified using PCR oligonucleotide primers corresponding to the 5' and 3' ends of the DNA sequence, as outlined in Example 1, to synthesize insertion fragments. The primers used to amplify the cDNA insert should preferably contain restriction sites, such as BamHI and XbaI, at the 5' end of the primers in order to clone the amplified product into the expression

vector. For example, BamHI and XbaI correspond to the restriction enzyme sites on the bacterial expression vector pQE-9. (Qiagen, Inc., Chatsworth, CA). This plasmid vector encodes antibiotic resistance (Amp^r), a bacterial origin of replication (ori), an IPTG-regulatable promoter/operator (P/O), a ribosome binding site (RBS), a 6-histidine tag (6-His), and restriction enzyme cloning sites.

[945] The pQE-9 vector is digested with BamHI and XbaI and the amplified fragment is ligated into the pQE-9 vector maintaining the reading frame initiated at the bacterial RBS. The ligation mixture is then used to transform the *E. coli* strain M15/rep4 (Qiagen, Inc.) which contains multiple copies of the plasmid pREP4, which expresses the lacI repressor and also confers kanamycin resistance (Kan^r). Transformants are identified by their ability to grow on LB plates and ampicillin/kanamycin resistant colonies are selected. Plasmid DNA is isolated and confirmed by restriction analysis.

[946] Clones containing the desired constructs are grown overnight (O/N) in liquid culture in LB media supplemented with both Amp (100 ug/ml) and Kan (25 ug/ml). The O/N culture is used to inoculate a large culture at a ratio of 1:100 to 1:250. The cells are grown to an optical density 600 ($O.D.^{600}$) of between 0.4 and 0.6. IPTG (Isopropyl-B-D-thiogalacto pyranoside) is then added to a final concentration of 1 mM. IPTG induces by inactivating the lacI repressor, clearing the P/O leading to increased gene expression.

[947] Cells are grown for an extra 3 to 4 hours. Cells are then harvested by centrifugation (20 mins at 6000Xg). The cell pellet is solubilized in the chaotropic agent 6 Molar Guanidine HCl by stirring for 3-4 hours at 4°C. The cell debris is removed by centrifugation, and the supernatant containing the polypeptide is loaded onto a nickel-nitrilo-tri-acetic acid ("Ni-NTA") affinity resin column (available from QIAGEN, Inc., *supra*). Proteins with a 6 x His tag bind to the Ni-NTA resin with high affinity and can be purified in a simple one-step procedure (for details see: The QIAexpressionist (1995) QIAGEN, Inc., *supra*).

[948] Briefly, the supernatant is loaded onto the column in 6 M guanidine-HCl, pH 8. The column is first washed with 10 volumes of 6 M guanidine-HCl, pH 8, then washed with 10 volumes of 6 M guanidine-HCl pH 6, and finally the polypeptide is eluted with 6 M guanidine-HCl, pH 5.

[949] The purified protein is then renatured by dialyzing it against phosphate-buffered saline (PBS) or 50 mM Na-acetate, pH 6 buffer plus 200 mM NaCl. Alternatively, the protein can be successfully refolded while immobilized on the Ni-NTA column. The

recommended conditions are as follows: renature using a linear 6M-1M urea gradient in 500 mM NaCl, 20% glycerol, 20 mM Tris/HCl pH 7.4, containing protease inhibitors. The renaturation should be performed over a period of 1.5 hours or more. After renaturation the proteins are eluted by the addition of 250 mM imidazole. Imidazole is removed by a final dialyzing step against PBS or 50 mM sodium acetate pH 6 buffer plus 200 mM NaCl. The purified protein is stored at 4°C or frozen at -80°C.

[950] In addition to the above expression vector, the present invention further includes an expression vector, called pHE4a (ATCC Accession Number 209645, deposited on February 25, 1998) which contains phage operator and promoter elements operatively linked to a polynucleotide of the present invention, called pHE4a. (ATCC Accession Number 209645, deposited on February 25, 1998.) This vector contains: 1) a neomycinphosphotransferase gene as a selection marker, 2) an *E. coli* origin of replication, 3) a T5 phage promoter sequence, 4) two lac operator sequences, 5) a Shine-Delgarno sequence, and 6) the lactose operon repressor gene (*lacIq*). The origin of replication (*oriC*) is derived from pUC19 (LTI, Gaithersburg, MD). The promoter and operator sequences are made synthetically.

[951] DNA can be inserted into the pHE4a by restricting the vector with NdeI and XbaI, BamHI, XhoI, or Asp718, running the restricted product on a gel, and isolating the larger fragment (the stuffer fragment should be about 310 base pairs). The DNA insert is generated according to the PCR protocol described in Example 1, using PCR primers having restriction sites for NdeI (5' primer) and XbaI, BamHI, XhoI, or Asp718 (3' primer). The PCR insert is gel purified and restricted with compatible enzymes. The insert and vector are ligated according to standard protocols.

[952] The engineered vector could easily be substituted in the above protocol to express protein in a bacterial system.

Example 6: Purification of a Polypeptide from an Inclusion Body

[953] The following alternative method can be used to purify a polypeptide expressed in *E. coli* when it is present in the form of inclusion bodies. Unless otherwise specified, all of the following steps are conducted at 4-10°C.

[954] Upon completion of the production phase of the *E. coli* fermentation, the cell culture is cooled to 4-10°C and the cells harvested by continuous centrifugation at 15,000 rpm (Heraeus Sepatech). On the basis of the expected yield of protein per unit weight of cell paste and the amount of purified protein required, an appropriate amount of cell paste, by weight, is suspended in a buffer solution containing 100 mM Tris, 50 mM EDTA, pH 7.4. The cells are dispersed to a homogeneous suspension using a high shear mixer.

[955] The cells are then lysed by passing the solution through a microfluidizer (Microfluidics, Corp. or APV Gaulin, Inc.) twice at 4000-6000 psi. The homogenate is then mixed with NaCl solution to a final concentration of 0.5 M NaCl, followed by centrifugation at 7000 xg for 15 min. The resultant pellet is washed again using 0.5M NaCl, 100 mM Tris, 50 mM EDTA, pH 7.4.

[956] The resulting washed inclusion bodies are solubilized with 1.5 M guanidine hydrochloride (GuHCl) for 2-4 hours. After 7000 xg centrifugation for 15 min., the pellet is discarded and the polypeptide containing supernatant is incubated at 4°C overnight to allow further GuHCl extraction.

[957] Following high speed centrifugation (30,000 xg) to remove insoluble particles, the GuHCl solubilized protein is refolded by quickly mixing the GuHCl extract with 20 volumes of buffer containing 50 mM sodium, pH 4.5, 150 mM NaCl, 2 mM EDTA by vigorous stirring. The refolded diluted protein solution is kept at 4°C without mixing for 12 hours prior to further purification steps.

[958] To clarify the refolded polypeptide solution, a previously prepared tangential filtration unit equipped with 0.16 µm membrane filter with appropriate surface area (e.g., Filtron), equilibrated with 40 mM sodium acetate, pH 6.0 is employed. The filtered sample is loaded onto a cation exchange resin (e.g., Poros HS-50, Perseptive Biosystems). The column is washed with 40 mM sodium acetate, pH 6.0 and eluted with 250 mM, 500 mM, 1000 mM, and 1500 mM NaCl in the same buffer, in a stepwise manner. The absorbance at 280 nm of the effluent is continuously monitored. Fractions are collected and further analyzed by SDS-PAGE.

[959] Fractions containing the polypeptide are then pooled and mixed with 4 volumes of water. The diluted sample is then loaded onto a previously prepared set of tandem columns of strong anion (Poros HQ-50, Perseptive Biosystems) and weak anion (Poros CM-20, Perseptive Biosystems) exchange resins. The columns are equilibrated with 40 mM sodium acetate, pH 6.0. Both columns are washed with 40 mM sodium acetate, pH

6.0, 200 mM NaCl. The CM-20 column is then eluted using a 10 column volume linear gradient ranging from 0.2 M NaCl, 50 mM sodium acetate, pH 6.0 to 1.0 M NaCl, 50 mM sodium acetate, pH 6.5. Fractions are collected under constant A₂₈₀ monitoring of the effluent. Fractions containing the polypeptide (determined, for instance, by 16% SDS-PAGE) are then pooled.

[960] The resultant polypeptide should exhibit greater than 95% purity after the above refolding and purification steps. No major contaminant bands should be observed from Commassie blue stained 16% SDS-PAGE gel when 5 µg of purified protein is loaded. The purified protein can also be tested for endotoxin/LPS contamination, and typically the LPS content is less than 0.1 ng/ml according to LAL assays.

Example 7: Cloning and Expression of a Polypeptide in a Baculovirus Expression System

[961] In this example, the plasmid shuttle vector pA2 is used to insert a polynucleotide into a baculovirus to express a polypeptide. This expression vector contains the strong polyhedrin promoter of the *Autographa californica* nuclear polyhedrosis virus (AcMNPV) followed by convenient restriction sites such as BamHI, Xba I and Asp718. The polyadenylation site of the simian virus 40 ("SV40") is used for efficient polyadenylation. For easy selection of recombinant virus, the plasmid contains the beta-galactosidase gene from *E. coli* under control of a weak *Drosophila* promoter in the same orientation, followed by the polyadenylation signal of the polyhedrin gene. The inserted genes are flanked on both sides by viral sequences for cell-mediated homologous recombination with wild-type viral DNA to generate a viable virus that express the cloned polynucleotide.

[962] Many other baculovirus vectors can be used in place of the vector above, such as pAc373, pVL941, and pAcIM1, as one skilled in the art would readily appreciate, as long as the construct provides appropriately located signals for transcription, translation, secretion and the like, including a signal peptide and an in-frame AUG as required. Such vectors are described, for instance, in Luckow et al., *Virology* 170:31-39 (1989).

[963] Specifically, the cDNA sequence contained in the deposited clone, including the AUG initiation codon, is amplified using the PCR protocol described in Example 1. If a naturally occurring signal sequence is used to produce the polypeptide of the present

invention, the pA2 vector does not need a second signal peptide. Alternatively, the vector can be modified (pA2 GP) to include a baculovirus leader sequence, using the standard methods described in Summers et al., "A Manual of Methods for Baculovirus Vectors and Insect Cell Culture Procedures," Texas Agricultural Experimental Station Bulletin No. 1555 (1987).

[964] The amplified fragment is isolated from a 1% agarose gel using a commercially available kit ("GeneClean," BIO 101 Inc., La Jolla, Ca.). The fragment then is digested with appropriate restriction enzymes and again purified on a 1% agarose gel.

[965] The plasmid is digested with the corresponding restriction enzymes and optionally, can be dephosphorylated using calf intestinal phosphatase, using routine procedures known in the art. The DNA is then isolated from a 1% agarose gel using a commercially available kit ("GeneClean" BIO 101 Inc., La Jolla, Ca.).

[966] The fragment and the dephosphorylated plasmid are ligated together with T4 DNA ligase. *E. coli* HB101 or other suitable *E. coli* hosts such as XL-1 Blue (Stratagene Cloning Systems, La Jolla, CA) cells are transformed with the ligation mixture and spread on culture plates. Bacteria containing the plasmid are identified by digesting DNA from individual colonies and analyzing the digestion product by gel electrophoresis. The sequence of the cloned fragment is confirmed by DNA sequencing.

[967] Five μg of a plasmid containing the polynucleotide is co-transfected with 1.0 μg of a commercially available linearized baculovirus DNA ("BaculoGold™ baculovirus DNA, Pharmingen, San Diego, CA), using the lipofection method described by Felgner et al., Proc. Natl. Acad. Sci. USA 84:7413-7417 (1987). One μg of BaculoGold™ virus DNA and 5 μg of the plasmid are mixed in a sterile well of a microtiter plate containing 50 μl of serum-free Grace's medium (Life Technologies Inc., Gaithersburg, MD). Afterwards, 10 μl Lipofectin plus 90 μl Grace's medium are added, mixed and incubated for 15 minutes at room temperature. Then the transfection mixture is added drop-wise to Sf9 insect cells (ATCC CRL 1711) seeded in a 35 mm tissue culture plate with 1 ml Grace's medium without serum. The plate is then incubated for 5 hours at 27° C. The transfection solution is then removed from the plate and 1 ml of Grace's insect medium supplemented with 10% fetal calf serum is added. Cultivation is then continued at 27° C for four days.

[968] After four days the supernatant is collected and a plaque assay is performed, as described by Summers and Smith, *supra*. An agarose gel with "Blue Gal" (Life Technologies Inc., Gaithersburg) is used to allow easy identification and isolation of gal-

expressing clones, which produce blue-stained plaques. (A detailed description of a "plaque assay" of this type can also be found in the user's guide for insect cell culture and baculovirology distributed by Life Technologies Inc., Gaithersburg, page 9-10.) After appropriate incubation, blue stained plaques are picked with the tip of a micropipettor (e.g., Eppendorf). The agar containing the recombinant viruses is then resuspended in a microcentrifuge tube containing 200 μ l of Grace's medium and the suspension containing the recombinant baculovirus is used to infect Sf9 cells seeded in 35 mm dishes. Four days later the supernatants of these culture dishes are harvested and then they are stored at 4° C.

[969] To verify the expression of the polypeptide, Sf9 cells are grown in Grace's medium supplemented with 10% heat-inactivated FBS. The cells are infected with the recombinant baculovirus containing the polynucleotide at a multiplicity of infection ("MOI") of about 2. If radiolabeled proteins are desired, 6 hours later the medium is removed and is replaced with SF900 II medium minus methionine and cysteine (available from Life Technologies Inc., Rockville, MD). After 42 hours, 5 μ Ci of 35 S-methionine and 5 μ Ci 35 S-cysteine (available from Amersham) are added. The cells are further incubated for 16 hours and then are harvested by centrifugation. The proteins in the supernatant as well as the intracellular proteins are analyzed by SDS-PAGE followed by autoradiography (if radiolabeled).

[970] Microsequencing of the amino acid sequence of the amino terminus of purified protein may be used to determine the amino terminal sequence of the produced protein.

Example 8: Expression of a Polypeptide in Mammalian Cells

[971] The polypeptide of the present invention can be expressed in a mammalian cell. A typical mammalian expression vector contains a promoter element, which mediates the initiation of transcription of mRNA, a protein coding sequence, and signals required for the termination of transcription and polyadenylation of the transcript. Additional elements include enhancers, Kozak sequences and intervening sequences flanked by donor and acceptor sites for RNA splicing. Highly efficient transcription is achieved with the early and late promoters from SV40, the long terminal repeats (LTRs) from Retroviruses, e.g., RSV, HTLV, HIV and the early promoter of the cytomegalovirus (CMV). However, cellular elements can also be used (e.g., the human actin promoter).

[972] Suitable expression vectors for use in practicing the present invention include, for example, vectors such as pSVL and pMSG (Pharmacia, Uppsala, Sweden), pRSVcat (ATCC 37152), pSV2dhfr (ATCC 37146), pBC12MI (ATCC 67109), pCMVSPORT 2.0, and pCMVSPORT 3.0. Mammalian host cells that could be used include, human Hela, 293, H9 and Jurkat cells, mouse NIH3T3 and C127 cells, Cos 1, Cos 7 and CV1, quail QC1-3 cells, mouse L cells and Chinese hamster ovary (CHO) cells.

[973] Alternatively, the polypeptide can be expressed in stable cell lines containing the polynucleotide integrated into a chromosome. The co-transfection with a selectable marker such as DHFR, gpt, neomycin, or hygromycin allows the identification and isolation of the transfected cells.

[974] The transfected gene can also be amplified to express large amounts of the encoded protein. The DHFR (dihydrofolate reductase) marker is useful in developing cell lines that carry several hundred or even several thousand copies of the gene of interest. (See, e.g., Alt, F. W., et al., J. Biol. Chem. 253:1357-1370 (1978); Hamlin, J. L. and Ma, C., Biochem. et Biophys. Acta, 1097:107-143 (1990); Page, M. J. and Sydenham, M. A., Biotechnology 9:64-68 (1991)). Another useful selection marker is the enzyme glutamine synthase (GS) (Murphy et al., Biochem J. 227:277-279 (1991); Bebbington et al., Bio/Technology 10:169-175 (1992)). Using these markers, the mammalian cells are grown in selective medium and the cells with the highest resistance are selected. These cell lines contain the amplified gene(s) integrated into a chromosome. Chinese hamster ovary (CHO) and NSO cells are often used for the production of proteins.

[975] Derivatives of the plasmid pSV2-dhfr (ATCC Accession No. 37146), the expression vectors pC4 (ATCC Accession No. 209646) and pC6 (ATCC Accession No.209647) contain the strong promoter (LTR) of the Rous Sarcoma Virus (Cullen et al., Molecular and Cellular Biology, 438-447 (March, 1985)) plus a fragment of the CMV-enhancer (Boshart et al., Cell 41:521-530 (1985)). Multiple cloning sites, e.g., with the restriction enzyme cleavage sites BamHI, XbaI and Asp718, facilitate the cloning of the gene of interest. The vectors also contain the 3' intron, the polyadenylation and termination signal of the rat preproinsulin gene, and the mouse DHFR gene under control of the SV40 early promoter.

[976] Specifically, the plasmid pC6, for example, is digested with appropriate restriction enzymes and then dephosphorylated using calf intestinal phosphates by procedures known in the art. The vector is then isolated from a 1% agarose gel.

[977] A polynucleotide of the present invention is amplified according to the protocol outlined in Example 1. If a naturally occurring signal sequence is used to produce the polypeptide of the present invention, the vector does not need a second signal peptide. Alternatively, if a naturally occurring signal sequence is not used, the vector can be modified to include a heterologous signal sequence. (See, e.g., International Publication No. WO 96/34891.)

[978] The amplified fragment is isolated from a 1% agarose gel using a commercially available kit ("GeneClean," BIO 101 Inc., La Jolla, Ca.). The fragment then is digested with appropriate restriction enzymes and again purified on a 1% agarose gel.

[979] The amplified fragment is then digested with the same restriction enzyme and purified on a 1% agarose gel. The isolated fragment and the dephosphorylated vector are then ligated with T4 DNA ligase. *E. coli* HB101 or XL-1 Blue cells are then transformed and bacteria are identified that contain the fragment inserted into plasmid pC6 using, for instance, restriction enzyme analysis.

[980] Chinese hamster ovary cells lacking an active DHFR gene is used for transfection. Five μ g of the expression plasmid pC6 or pC4 is cotransfected with 0.5 μ g of the plasmid pSVneo using lipofectin (Felgner et al., *supra*). The plasmid pSV2-neo contains a dominant selectable marker, the *neo* gene from Tn5 encoding an enzyme that confers resistance to a group of antibiotics including G418. The cells are seeded in alpha minus MEM supplemented with 1 mg/ml G418. After 2 days, the cells are trypsinized and seeded in hybridoma cloning plates (Greiner, Germany) in alpha minus MEM supplemented with 10, 25, or 50 ng/ml of methotrexate plus 1 mg/ml G418. After about 10-14 days single clones are trypsinized and then seeded in 6-well petri dishes or 10 ml flasks using different concentrations of methotrexate (50 nM, 100 nM, 200 nM, 400 nM, 800 nM). Clones growing at the highest concentrations of methotrexate are then transferred to new 6-well plates containing even higher concentrations of methotrexate (1 μ M, 2 μ M, 5 μ M, 10 mM, 20 mM). The same procedure is repeated until clones are obtained which grow at a concentration of 100 - 200 μ M. Expression of the desired gene product is analyzed, for instance, by SDS-PAGE and Western blot or by reversed phase HPLC analysis.

Example 9: Protein Fusions

[981] The polypeptides of the present invention are preferably fused to other proteins. These fusion proteins can be used for a variety of applications. For example, fusion of the present polypeptides to His-tag, HA-tag, protein A, IgG domains, and maltose binding protein facilitates purification. (See Example 5; see also EP A 394,827; Traunecker, et al., Nature 331:84-86 (1988)). Similarly, fusion to IgG-1, IgG-3, and albumin increases the half-life time *in vivo*. Nuclear localization signals fused to the polypeptides of the present invention can target the protein to a specific subcellular localization, while covalent heterodimer or homodimers can increase or decrease the activity of a fusion protein. Fusion proteins can also create chimeric molecules having more than one function. Finally, fusion proteins can increase solubility and/or stability of the fused protein compared to the non-fused protein. All of the types of fusion proteins described above can be made by modifying the following protocol, which outlines the fusion of a polypeptide to an IgG molecule, or the protocol described in Example 5.

[982] Briefly, the human Fc portion of the IgG molecule can be PCR amplified, using primers that span the 5' and 3' ends of the sequence described below. These primers also should have convenient restriction enzyme sites that will facilitate cloning into an expression vector, preferably a mammalian expression vector.

[983] For example, if pC4 (ATCC Accession No. 209646) is used, the human Fc portion can be ligated into the BamHI cloning site. Note that the 3' BamHI site should be destroyed. Next, the vector containing the human Fc portion is re-restricted with BamHI, linearizing the vector, and a polynucleotide of the present invention, isolated by the PCR protocol described in Example 1, is ligated into this BamHI site. Note that the polynucleotide is cloned without a stop codon, otherwise a fusion protein will not be produced.

[984] If the naturally occurring signal sequence is used to produce the polypeptide of the present invention, pC4 does not need a second signal peptide. Alternatively, if the naturally occurring signal sequence is not used, the vector can be modified to include a heterologous signal sequence. (See, e.g., International Publication No. WO 96/34891.)

Human IgG Fc region:

GGGATCCGGAGCCCAAATCTTCTGACAAACTCACACATGCCCACCGTGCCCA
 GCACCTGAATTCGAGGGTGCACCGTCAGTCTTCCTCTTCCCCCAAACCCAAG
 GACACCCTCATGATCTCCCGGACTCCTGAGGTCACATGCGTGGTGGTGGACGTA
 AGCCACGAAGACCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGGAGGT
 GCATAATGCCAAGACAAAGCCGCGGGAGGAGCAGTACAACAGCACGTACCGTG
 TGGTCAGCGTCCTCACCGTCCTGCACCAGGACTGGCTGAATGGCAAGGAGTAC
 AAGTGCAAGGTCTCCAACAAAGCCCTCCCAACCCCATCGAGAAAACCATCTC
 CAAAGCCAAAGGGCAGCCCCGAGAACCACAGGTGTACACCCTGCCCCCATCCC
 GGGATGAGCTGACCAAGAACCAGGTCAGCCTGACCTGCCTGGTCAAAGGCTTC
 TATCCAAGCGACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAACAA
 CTACAAGACCACGCCTCCCGTGCTGGACTCCGACGGCTCCTTCTTCCTCTACAG
 CAAGCTCACCGTGGACAAGAGCAGGTGGCAGCAGGGGAACGTCTTCTCATGCT
 CCGTGATGCATGAGGCTCTGCACAACCACTACACGCAGAAGAGCCTCTCCCTGT
 CTCCGGGTAAATGAGTGCGACGGCCGCGACTCTAGAGGAT (SEQ ID NO: 1)

Example 10: Production of an Antibody from a Polypeptide

a) Hybridoma Technology

[985] The antibodies of the present invention can be prepared by a variety of methods. (See, Current Protocols, Chapter 2.) As one example of such methods, cells expressing a polypeptide of the present invention are administered to an animal to induce the production of sera containing polyclonal antibodies. In a preferred method, a preparation of a polypeptide of the present invention is prepared and purified to render it substantially free of natural contaminants. Such a preparation is then introduced into an animal in order to produce polyclonal antisera of greater specific activity.

[986] Monoclonal antibodies specific for a polypeptide of the present invention are prepared using hybridoma technology (Kohler et al., Nature 256:495 (1975); Kohler et al., Eur. J. Immunol. 6:511 (1976); Kohler et al., Eur. J. Immunol. 6:292 (1976); Hammerling et al., in: Monoclonal Antibodies and T-Cell Hybridomas, Elsevier, N.Y., pp. 563-681 (1981)). In general, an animal (preferably a mouse) is immunized with a polypeptide of the present invention or, more preferably, with a secreted polypeptide-expressing cell. Such polypeptide-expressing cells are cultured in any suitable tissue culture medium, preferably in Earle's modified Eagle's medium supplemented with 10% fetal bovine serum

(inactivated at about 56°C), and supplemented with about 10 g/l of nonessential amino acids, about 1,000 U/ml of penicillin, and about 100 µg/ml of streptomycin.

[987] The splenocytes of such mice are extracted and fused with a suitable myeloma cell line. Any suitable myeloma cell line may be employed in accordance with the present invention; however, it is preferable to employ the parent myeloma cell line (SP2O), available from the ATCC. After fusion, the resulting hybridoma cells are selectively maintained in HAT medium, and then cloned by limiting dilution as described by Wands et al. (Gastroenterology 80:225-232 (1981)). The hybridoma cells obtained through such a selection are then assayed to identify clones which secrete antibodies capable of binding the polypeptide of the present invention .

[988] Alternatively, additional antibodies capable of binding to a polypeptide of the present invention can be produced in a two-step procedure using anti-idiotypic antibodies. Such a method makes use of the fact that antibodies are themselves antigens, and therefore, it is possible to obtain an antibody which binds to a second antibody. In accordance with this method, protein specific antibodies are used to immunize an animal, preferably a mouse. The splenocytes of such an animal are then used to produce hybridoma cells, and the hybridoma cells are screened to identify clones which produce an antibody whose ability to bind to the polypeptide-specific antibody can be blocked by said polypeptide. Such antibodies comprise anti-idiotypic antibodies to the polypeptide-specific antibody and are used to immunize an animal to induce formation of further polypeptide-specific antibodies.

[989] For *in vivo* use of antibodies in humans, an antibody is "humanized". Such antibodies can be produced using genetic constructs derived from hybridoma cells producing the monoclonal antibodies described above. Methods for producing chimeric and humanized antibodies are known in the art and are discussed herein. (See, for review, Morrison, Science 229:1202 (1985); Oi et al., BioTechniques 4:214 (1986); Cabilly et al., U.S. Patent No. 4,816,567; Taniguchi et al., EP 171496; Morrison et al., EP 173494; Neuberger et al., WO 8601533; Robinson et al., International Publication No. WO 8702671; Boulianne et al., Nature 312:643 (1984); Neuberger et al., Nature 314:268 (1985)).

b) Isolation Of Antibody Fragments Directed Against a Polypeptide of the Present Invention From A Library Of scFvs

[990] Naturally occurring V-genes isolated from human PBLs are constructed into a library of antibody fragments which contain reactivities against a polypeptide of the present invention to which the donor may or may not have been exposed (see e.g., U.S. Patent 5,885,793 incorporated herein by reference in its entirety).

[991] *Rescue of the Library.* A library of scFvs is constructed from the RNA of human PBLs as described in International Publication No. WO 92/01047. To rescue phage displaying antibody fragments, approximately 10^9 *E. coli* harboring the phagemid are used to inoculate 50 ml of 2xTY containing 1% glucose and 100 μ g/ml of ampicillin (2xTY-AMP-GLU) and grown to an O.D. of 0.8 with shaking. Five ml of this culture is used to inoculate 50 ml of 2xTY-AMP-GLU, 2×10^8 TU of delta gene 3 helper (M13 delta gene III, see International Publication No. WO 92/01047) are added and the culture incubated at 37°C for 45 minutes without shaking and then at 37°C for 45 minutes with shaking. The culture is centrifuged at 4000 r.p.m. for 10 min. and the pellet resuspended in 2 liters of 2xTY containing 100 μ g/ml ampicillin and 50 μ g/ml kanamycin and grown overnight. Phage are prepared as described in International Publication No. WO 92/01047.

[992] M13 delta gene III is prepared as follows: M13 delta gene III helper phage does not encode gene III protein, hence the phage(mid) displaying antibody fragments have a greater avidity of binding to antigen. Infectious M13 delta gene III particles are made by growing the helper phage in cells harboring a pUC19 derivative supplying the wild type gene III protein during phage morphogenesis. The culture is incubated for 1 hour at 37°C without shaking and then for a further hour at 37°C with shaking. Cells are spun down (IEC-Centra 8,400 r.p.m. for 10 min), resuspended in 300 ml 2xTY broth containing 100 μ g ampicillin/ml and 25 μ g kanamycin/ml (2xTY-AMP-KAN) and grown overnight, shaking at 37°C. Phage particles are purified and concentrated from the culture medium by two PEG-precipitations (Sambrook et al., 1990), resuspended in 2 ml PBS and passed through a 0.45 μ m filter (Minisart NML; Sartorius) to give a final concentration of approximately 10^{13} transducing units/ml (ampicillin-resistant clones).

[993] *Panning of the Library.* Immunotubes (Nunc) are coated overnight in PBS with 4 ml of either 100 μ g/ml or 10 μ g/ml of a polypeptide of the present invention. Tubes are blocked with 2% Marvel-PBS for 2 hours at 37°C and then washed 3 times in PBS. Approximately 10^{13} TU of phage is applied to the tube and incubated for 30 minutes at room temperature tumbling on an over and under turntable and then left to stand for another 1.5 hours. Tubes are washed 10 times with PBS 0.1% Tween-20 and 10 times with PBS.

Phage are eluted by adding 1 ml of 100 mM triethylamine and rotating 15 minutes on an under and over turntable after which the solution is immediately neutralized with 0.5 ml of 1.0M Tris-HCl, pH 7.4. Phage are then used to infect 10 ml of mid-log E. coli TG1 by incubating eluted phage with bacteria for 30 minutes at 37°C. The E. coli are then plated on TYE plates containing 1% glucose and 100 µg/ml ampicillin. The resulting bacterial library is then rescued with delta gene 3 helper phage as described above to prepare phage for a subsequent round of selection. This process is then repeated for a total of 4 rounds of affinity purification with tube-washing increased to 20 times with PBS, 0.1% Tween-20 and 20 times with PBS for rounds 3 and 4.

[994] *Characterization of Binders.* Eluted phage from the 3rd and 4th rounds of selection are used to infect E. coli HB 2151 and soluble scFv is produced (Marks, et al., 1991) from single colonies for assay. ELISAs are performed with microtitre plates coated with either 10 pg/ml of the polypeptide of the present invention in 50 mM bicarbonate pH 9.6. Clones positive in ELISA are further characterized by PCR fingerprinting (see, e.g., International Publication No. WO 92/01047) and then by sequencing. These ELISA positive clones may also be further characterized by techniques known in the art, such as, for example, epitope mapping, binding affinity, receptor signal transduction, ability to block or competitively inhibit antibody/antigen binding, and competitive agonistic or antagonistic activity.

Example 11: Method of Determining Alterations in a Gene Corresponding to a Polynucleotide

[995] RNA isolated from entire families or individual patients presenting with a phenotype of interest (such as a disease) is isolated. cDNA is then generated from these RNA samples using protocols known in the art. (See, Sambrook.) The cDNA is then used as a template for PCR, employing primers surrounding regions of interest in SEQ ID NO:X; and/or the nucleotide sequence of the cDNA contained in Clone ID NO:Z. Suggested PCR conditions consist of 35 cycles at 95 degrees C for 30 seconds; 60-120 seconds at 52-58 degrees C; and 60-120 seconds at 70 degrees C, using buffer solutions described in Sidransky et al., Science 252:706 (1991).

[996] PCR products are then sequenced using primers labeled at their 5' end with T4 polynucleotide kinase, employing SequiTherm Polymerase (Epicentre Technologies). The intron-exon boundaries of selected exons is also determined and genomic PCR products analyzed to confirm the results. PCR products harboring suspected mutations are then cloned and sequenced to validate the results of the direct sequencing.

[997] PCR products are cloned into T-tailed vectors as described in Holton et al., Nucleic Acids Research, 19:1156 (1991) and sequenced with T7 polymerase (United States Biochemical). Affected individuals are identified by mutations not present in unaffected individuals.

[998] Genomic rearrangements are also observed as a method of determining alterations in a gene corresponding to a polynucleotide. Genomic clones isolated according to Example 2 are nick-translated with digoxigenin deoxy-uridine 5'-triphosphate (Boehringer Mannheim), and FISH performed as described in Johnson et al., Methods Cell Biol. 35:73-99 (1991). Hybridization with the labeled probe is carried out using a vast excess of human cot-1 DNA for specific hybridization to the corresponding genomic locus.

[999] Chromosomes are counterstained with 4,6-diamino-2-phenylidole and propidium iodide, producing a combination of C- and R-bands. Aligned images for precise mapping are obtained using a triple-band filter set (Chroma Technology, Brattleboro, VT) in combination with a cooled charge-coupled device camera (Photometrics, Tucson, AZ) and variable excitation wavelength filters. (Johnson et al., Genet. Anal. Tech. Appl., 8:75 (1991)). Image collection, analysis and chromosomal fractional length measurements are performed using the ISee Graphical Program System. (Inovision Corporation, Durham, NC.) Chromosome alterations of the genomic region hybridized by the probe are identified as insertions, deletions, and translocations. These alterations are used as a diagnostic marker for an associated disease.

Example 12: Method of Detecting Abnormal Levels of a Polypeptide in a Biological Sample

[1000] A polypeptide of the present invention can be detected in a biological sample, and if an increased or decreased level of the polypeptide is detected, this polypeptide is a marker for a particular phenotype. Methods of detection are numerous, and thus, it is

understood that one skilled in the art can modify the following assay to fit their particular needs.

[1001] For example, antibody-sandwich ELISAs are used to detect polypeptides in a sample, preferably a biological sample. Wells of a microtiter plate are coated with specific antibodies, at a final concentration of 0.2 to 10 ug/ml. The antibodies are either monoclonal or polyclonal and are produced by the method described in Example 10. The wells are blocked so that non-specific binding of the polypeptide to the well is reduced.

[1002] The coated wells are then incubated for > 2 hours at RT with a sample containing the polypeptide. Preferably, serial dilutions of the sample should be used to validate results. The plates are then washed three times with deionized or distilled water to remove unbound polypeptide.

[1003] Next, 50 ul of specific antibody-alkaline phosphatase conjugate, at a concentration of 25-400 ng, is added and incubated for 2 hours at room temperature. The plates are again washed three times with deionized or distilled water to remove unbound conjugate.

[1004] Add 75 ul of 4-methylumbelliferyl phosphate (MUP) or p-nitrophenyl phosphate (NPP) substrate solution to each well and incubate 1 hour at room temperature. Measure the reaction by a microtiter plate reader. Prepare a standard curve, using serial dilutions of a control sample, and plot polypeptide concentration on the X-axis (log scale) and fluorescence or absorbance of the Y-axis (linear scale). Interpolate the concentration of the polypeptide in the sample using the standard curve.

Example 13: Formulation

[1005] The invention also provides methods of treatment and/or prevention of diseases or disorders (such as, for example, any one or more of the diseases or disorders disclosed herein) by administration to a subject of an effective amount of a Therapeutic. By therapeutic is meant polynucleotides or polypeptides of the invention (including fragments and variants), agonists or antagonists thereof, and/or antibodies thereto, in combination with a pharmaceutically acceptable carrier type (e.g., a sterile carrier).

[1006] The Therapeutic will be formulated and dosed in a fashion consistent with good medical practice, taking into account the clinical condition of the individual patient (especially the side effects of treatment with the Therapeutic alone), the site of delivery, the

method of administration, the scheduling of administration, and other factors known to practitioners. The "effective amount" for purposes herein is thus determined by such considerations.

[1007] As a general proposition, the total pharmaceutically effective amount of the Therapeutic administered parenterally per dose will be in the range of about 1ug/kg/day to 10 mg/kg/day of patient body weight, although, as noted above, this will be subject to therapeutic discretion. More preferably, this dose is at least 0.01 mg/kg/day, and most preferably for humans between about 0.01 and 1 mg/kg/day for the hormone. If given continuously, the Therapeutic is typically administered at a dose rate of about 1 ug/kg/hour to about 50 ug/kg/hour, either by 1-4 injections per day or by continuous subcutaneous infusions, for example, using a mini-pump. An intravenous bag solution may also be employed. The length of treatment needed to observe changes and the interval following treatment for responses to occur appears to vary depending on the desired effect.

[1008] Therapeutics can be administered orally, rectally, parenterally, intracisternally, intravaginally, intraperitoneally, topically (as by powders, ointments, gels, drops or transdermal patch), buccally, or as an oral or nasal spray. "Pharmaceutically acceptable carrier" refers to a non-toxic solid, semisolid or liquid filler, diluent, encapsulating material or formulation auxiliary of any. The term "parenteral" as used herein refers to modes of administration which include intravenous, intramuscular, intraperitoneal, intrasternal, subcutaneous and intraarticular injection and infusion.

[1009] Therapeutics of the invention are also suitably administered by sustained-release systems. Suitable examples of sustained-release Therapeutics are administered orally, rectally, parenterally, intracisternally, intravaginally, intraperitoneally, topically (as by powders, ointments, gels, drops or transdermal patch), buccally, or as an oral or nasal spray. "Pharmaceutically acceptable carrier" refers to a non-toxic solid, semisolid or liquid filler, diluent, encapsulating material or formulation auxiliary of any type. The term "parenteral" as used herein refers to modes of administration which include intravenous, intramuscular, intraperitoneal, intrasternal, subcutaneous and intraarticular injection and infusion.

[1010] Therapeutics of the invention are also suitably administered by sustained-release systems. Suitable examples of sustained-release Therapeutics include suitable polymeric materials (such as, for example, semi-permeable polymer matrices in the form of shaped articles, e.g., films, or microcapsules), suitable hydrophobic materials (for example as an

emulsion in an acceptable oil) or ion exchange resins, and sparingly soluble derivatives (such as, for example, a sparingly soluble salt).

[1011] Sustained-release matrices include polylactides (U.S. Pat. No. 3,773,919, EP 58,481), copolymers of L-glutamic acid and gamma-ethyl-L-glutamate (Sidman et al., *Biopolymers* 22:547-556 (1983)), poly (2- hydroxyethyl methacrylate) (Langer et al., *J. Biomed. Mater. Res.* 15:167-277 (1981), and Langer, *Chem. Tech.* 12:98-105 (1982)), ethylene vinyl acetate (Langer et al., *Id.*) or poly-D- (-)-3-hydroxybutyric acid (EP 133,988).

[1012] In a preferred embodiment, polypeptide, polynucleotide, and antibody compositions of the invention are formulated in a biodegradable, polymeric drug delivery system, for example as described in U.S. Patent Nos. 4,938,763; 5,278,201; 5,278,202; 5,324,519; 5,340,849; and 5,487,897 and in International Publication Numbers WO01/35929, WO00/24374, and WO00/06117 which are hereby incorporated by reference in their entirety. In specific preferred embodiments the polypeptide, polynucleotide, and antibody compositions of the invention are formulated using the ATRIGEL® Biodegradable System of Atrix Laboratories, Inc. (Fort Collins, Colorado).

[1013] Examples of biodegradable polymers which can be used in the formulation of polypeptide, polynucleotide, and antibody compositions, include but are not limited to, polylactides, polyglycolides, polycaprolactones, polyanhydrides, polyamides, polyurethanes, polyesteramides, polyorthoesters, polydioxanones, polyacetals, polyketals, polycarbonates, polyorthocarbonates, polyphosphazenes, polyhydroxybutyrates, polyhydroxyvalerates, polyalkylene oxalates, polyalkylene succinates, poly(malic acid), poly(amino acids), poly(methyl vinyl ether), poly(maleic anhydride), polyvinylpyrrolidone, polyethylene glycol, polyhydroxycellulose, chitin, chitosan, and copolymers, terpolymers, or combinations or mixtures of the above materials. The preferred polymers are those that have a lower degree of crystallization and are more hydrophobic. These polymers and copolymers are more soluble in the biocompatible solvents than the highly crystalline polymers such as polyglycolide and chitin which also have a high degree of hydrogen-bonding. Preferred materials with the desired solubility parameters are the polylactides, polycaprolactones, and copolymers of these with glycolide in which there are more amorphous regions to enhance solubility. In specific preferred embodiments, the biodegradable polymers which can be used in the formulation of polypeptide, polynucleotide, and antibody compositions are poly(lactide-co-glycolides). Polymer properties such as molecular weight, hydrophobicity, and lactide/glycolide ratio may be

modified to obtain the desired polypeptide, polynucleotide, or antibody release profile (See, e.g., Ravivarapu et al., Journal of Pharmaceutical Sciences 89:732-741 (2000), which is hereby incorporated by reference in its entirety).

[1014] It is also preferred that the solvent for the biodegradable polymer be non-toxic, water miscible, and otherwise biocompatible. Examples of such solvents include, but are not limited to, N-methyl-2-pyrrolidone, 2-pyrrolidone, C2 to C6 alkanols, C1 to C15 alcohols, dils, triols, and tetraols such as ethanol, glycerine propylene glycol, butanol; C3 to C15 alkyl ketones such as acetone, diethyl ketone and methyl ethyl ketone; C3 to C15 esters such as methyl acetate, ethyl acetate, ethyl lactate; alkyl ketones such as methyl ethyl ketone, C1 to C15 amides such as dimethylformamide, dimethylacetamide and caprolactam; C3 to C20 ethers such as tetrahydrofuran, or solketal; tweens, triacetin, propylene carbonate, decylmethylsulfoxide, dimethyl sulfoxide, oleic acid, 1-dodecylazacycloheptan-2-one, Other preferred solvents are benzyl alcohol, benzyl benzoate, dipropylene glycol, tributyrin, ethyl oleate, glycerin, glycofural, isopropyl myristate, isopropyl palmitate, oleic acid, polyethylene glycol, propylene carbonate, and triethyl citrate. The most preferred solvents are N-methyl-2-pyrrolidone, 2-pyrrolidone, dimethyl sulfoxide, triacetin, and propylene carbonate because of the solvating ability and their compatibility.

[1015] Additionally, formulations comprising polypeptide, polynucleotide, and antibody compositions and a biodegradable polymer may also include release-rate modification agents and/or pore-forming agents. Examples of release-rate modification agents include, but are not limited to, fatty acids, triglycerides, other like hydrophobic compounds, organic solvents, plasticizing compounds and hydrophilic compounds. Suitable release rate modification agents include, for example, esters of mono-, di-, and tricarboxylic acids, such as 2-ethoxyethyl acetate, methyl acetate, ethyl acetate, diethyl phthalate, dimethyl phthalate, dibutyl phthalate, dimethyl adipate, dimethyl succinate, dimethyl oxalate, dimethyl citrate, triethyl citrate, acetyl tributyl citrate, acetyl triethyl citrate, glycerol triacetate, di(n-butyl) sebecate, and the like; polyhydroxy alcohols, such as propylene glycol, polyethylene glycol, glycerin, sorbitol, and the like; fatty acids; triesters of glycerol, such as triglycerides, epoxidized soybean oil, and other epoxidized vegetable oils; sterols, such as cholesterol; alcohols, such as C.sub.6 -C.sub.12 alkanols, 2-ethoxyethanol. The release rate modification agent may be used singly or in combination with other such agents. Suitable combinations of release rate modification agents include, but are not limited to, glycerin/propylene glycol, sorbitol/glycerine, ethylene oxide/propylene oxide, butylene glycol/adipic acid, and the like. Preferred release rate modification agents include, but are

not limited to, dimethyl citrate, triethyl citrate, ethyl heptanoate, glycerin, and hexanediol. Suitable pore-forming agents that may be used in the polymer composition include, but are not limited to, sugars such as sucrose and dextrose, salts such as sodium chloride and sodium carbonate, polymers such as hydroxylpropylcellulose, carboxymethylcellulose, polyethylene glycol, and polyvinylpyrrolidone. Solid crystals that will provide a defined pore size, such as salt or sugar, are preferred.

[1016] In specific preferred embodiments the polypeptide, polynucleotide, and antibody compositions of the invention are formulated using the BEMA™ BioErodible Mucoadhesive System, MCA™ MucoCutaneous Absorption System, SMP™ Solvent MicroParticle System, or BCP™ BioCompatible Polymer System of Atrix Laboratories, Inc. (Fort Collins, Colorado).

[1017] Sustained-release Therapeutics also include liposomally entrapped Therapeutics of the invention (*see generally*, Langer, *Science* 249:1527-1533 (1990); Treat et al., in *Liposomes in the Therapy of Infectious Disease and Cancer*, Lopez-Berestein and Fidler (eds.), Liss, New York, pp. 317 -327 and 353-365 (1989)). Liposomes containing the Therapeutic are prepared by methods known per se: DE 3,218,121; Epstein et al., *Proc. Natl. Acad. Sci. (USA)* 82:3688-3692 (1985); Hwang et al., *Proc. Natl. Acad. Sci.(USA)* 77:4030-4034 (1980); EP 52,322; EP 36,676; EP 88,046; EP 143,949; EP 142,641; Japanese Pat. Appl. 83-118008; U.S. Pat. Nos. 4,485,045 and 4,544,545; and EP 102,324. Ordinarily, the liposomes are of the small (about 200-800 Angstroms) unilamellar type in which the lipid content is greater than about 30 mol. percent cholesterol, the selected proportion being adjusted for the optimal Therapeutic.

[1018] In yet an additional embodiment, the Therapeutics of the invention are delivered by way of a pump (*see* Langer, *supra*; Sefton, *CRC Crit. Ref. Biomed. Eng.* 14:201 (1987); Buchwald et al., *Surgery* 88:507 (1980); Saudek et al., *N. Engl. J. Med.* 321:574 (1989)).

[1019] Other controlled release systems are discussed in the review by Langer (*Science* 249:1527-1533 (1990)).

[1020] For parenteral administration, in one embodiment, the Therapeutic is formulated generally by mixing it at the desired degree of purity, in a unit dosage injectable form (solution, suspension, or emulsion), with a pharmaceutically acceptable carrier, i.e., one that is non-toxic to recipients at the dosages and concentrations employed and is compatible with other ingredients of the formulation. For example, the formulation preferably does not

include oxidizing agents and other compounds that are known to be deleterious to the Therapeutic.

[1021] Generally, the formulations are prepared by contacting the Therapeutic uniformly and intimately with liquid carriers or finely divided solid carriers or both. Then, if necessary, the product is shaped into the desired formulation. Preferably the carrier is a parenteral carrier, more preferably a solution that is isotonic with the blood of the recipient. Examples of such carrier vehicles include water, saline, Ringer's solution, and dextrose solution. Non-aqueous vehicles such as fixed oils and ethyl oleate are also useful herein, as well as liposomes.

[1022] The carrier suitably contains minor amounts of additives such as substances that enhance isotonicity and chemical stability. Such materials are non-toxic to recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate, succinate, acetic acid, and other organic acids or their salts; antioxidants such as ascorbic acid; low molecular weight (less than about ten residues) polypeptides, e.g., polyarginine or tripeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids, such as glycine, glutamic acid, aspartic acid, or arginine; monosaccharides, disaccharides, and other carbohydrates including cellulose or its derivatives, glucose, manose, or dextrans; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; counterions such as sodium; and/or nonionic surfactants such as polysorbates, poloxamers, or PEG.

[1023] The Therapeutic is typically formulated in such vehicles at a concentration of about 0.1 mg/ml to 100 mg/ml, preferably 1-10 mg/ml, at a pH of about 3 to 8. It will be understood that the use of certain of the foregoing excipients, carriers, or stabilizers will result in the formation of polypeptide salts.

[1024] Any pharmaceutical used for therapeutic administration can be sterile. Sterility is readily accomplished by filtration through sterile filtration membranes (e.g., 0.2 micron membranes). Therapeutics generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by a hypodermic injection needle.

[1025] Therapeutics ordinarily will be stored in unit or multi-dose containers, for example, sealed ampoules or vials, as an aqueous solution or as a lyophilized formulation for reconstitution. As an example of a lyophilized formulation, 10-ml vials are filled with 5 ml of sterile-filtered 1% (w/v) aqueous Therapeutic solution, and the resulting mixture is

lyophilized. The infusion solution is prepared by reconstituting the lyophilized Therapeutic using bacteriostatic Water-for-Injection.

[1026] The invention also provides a pharmaceutical pack or kit comprising one or more containers filled with one or more of the ingredients of the Therapeutics of the invention. Associated with such container(s) can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, which notice reflects approval by the agency of manufacture, use or sale for human administration. In addition, the Therapeutics may be employed in conjunction with other therapeutic compounds.

[1027] The Therapeutics of the invention may be administered alone or in combination with adjuvants. Adjuvants that may be administered with the Therapeutics of the invention include, but are not limited to, alum, alum plus deoxycholate (ImmunoAg), MTP-PE (Biocine Corp.), QS21 (Genentech, Inc.), BCG (e.g., THERACYS®), MPL and nonviable preparations of *Corynebacterium parvum*. In a specific embodiment, Therapeutics of the invention are administered in combination with alum. In another specific embodiment, Therapeutics of the invention are administered in combination with QS-21. Further adjuvants that may be administered with the Therapeutics of the invention include, but are not limited to, Monophosphoryl lipid immunomodulator, AdjuVax 100a, QS-21, QS-18, CRL1005, Aluminum salts, MF-59, and Virosomal adjuvant technology. Vaccines that may be administered with the Therapeutics of the invention include, but are not limited to, vaccines directed toward protection against MMR (measles, mumps, rubella), polio, varicella, tetanus/diphtheria, hepatitis A, hepatitis B, haemophilus influenzae B, whooping cough, pneumonia, influenza, Lyme's Disease, rotavirus, cholera, yellow fever, Japanese encephalitis, poliomyelitis, rabies, typhoid fever, and pertussis. Combinations may be administered either concomitantly, e.g., as an admixture, separately but simultaneously or concurrently; or sequentially. This includes presentations in which the combined agents are administered together as a therapeutic mixture, and also procedures in which the combined agents are administered separately but simultaneously, e.g., as through separate intravenous lines into the same individual. Administration "in combination" further includes the separate administration of one of the compounds or agents given first, followed by the second.

[1028] The Therapeutics of the invention may be administered alone or in combination with other therapeutic agents. Therapeutic agents that may be administered in combination

09950083-091201

compositions of the invention include, but are not limited to, aspirin, dipyridamole (e.g., PERSANTINE™), and ticlopidine (e.g., TICLID™).

[1032] In specific embodiments, the use of anti-coagulants, thrombolytic and/or antiplatelet drugs in combination with Therapeutics of the invention is contemplated for the prevention, diagnosis, and/or treatment of thrombosis, arterial thrombosis, venous thrombosis, thromboembolism, pulmonary embolism, atherosclerosis, myocardial infarction, transient ischemic attack, unstable angina. In specific embodiments, the use of anticoagulants, thrombolytic drugs and/or antiplatelet drugs in combination with Therapeutics of the invention is contemplated for the prevention of occlusion of saphenous grafts, for reducing the risk of periprocedural thrombosis as might accompany angioplasty procedures, for reducing the risk of stroke in patients with atrial fibrillation including nonrheumatic atrial fibrillation, for reducing the risk of embolism associated with mechanical heart valves and or mitral valves disease. Other uses for the therapeutics of the invention, alone or in combination with antiplatelet, anticoagulant, and/or thrombolytic drugs, include, but are not limited to, the prevention of occlusions in extracorporeal devices (e.g., intravascular canulas, vascular access shunts in hemodialysis patients, hemodialysis machines, and cardiopulmonary bypass machines).

[1033] In certain embodiments, Therapeutics of the invention are administered in combination with antiretroviral agents, nucleoside/nucleotide reverse transcriptase inhibitors (NRTIs), non-nucleoside reverse transcriptase inhibitors (NNRTIs), and/or protease inhibitors (PIs). NRTIs that may be administered in combination with the Therapeutics of the invention, include, but are not limited to, RETROVIR™ (zidovudine/AZT), VIDEX™ (didanosine/ddI), HIVID™ (zalcitabine/ddC), ZERIT™ (stavudine/d4T), EPIVIR™ (lamivudine/3TC), and COMBIVIR™ (zidovudine/lamivudine). NNRTIs that may be administered in combination with the Therapeutics of the invention, include, but are not limited to, VIRAMUNE™ (nevirapine), RESCRIPTOR™ (delavirdine), and SUSTIVA™ (efavirenz). Protease inhibitors that may be administered in combination with the Therapeutics of the invention, include, but are not limited to, CRIXIVAN™ (indinavir), NORVIR™ (ritonavir), INVIRASE™ (saquinavir), and VIRACEPT™ (nelfinavir). In a specific embodiment, antiretroviral agents, nucleoside reverse transcriptase inhibitors, non-nucleoside reverse transcriptase inhibitors, and/or protease

inhibitors may be used in any combination with Therapeutics of the invention to treat AIDS and/or to prevent or treat HIV infection.

[1034] Additional NRTIs include LODENOSINE™ (F-ddA; an acid-stable adenosine NRTI; Triangle/Abbott); COVIRACIL™ (emtricitabine/FTC; structurally related to lamivudine (3TC) but with 3- to 10-fold greater activity *in vitro*; Triangle/Abbott); dOTC (BCH-10652, also structurally related to lamivudine but retains activity against a substantial proportion of lamivudine-resistant isolates; Biochem Pharma); Adefovir (refused approval for anti-HIV therapy by FDA; Gilead Sciences); PREVEON® (Adefovir Dipivoxil, the active prodrug of adefovir; its active form is PMEA-pp); TENOFOVIR™ (bis-POC PMPA, a PMPA prodrug; Gilead); DAPD/DXG (active metabolite of DAPD; Triangle/Abbott); D-D4FC (related to 3TC, with activity against AZT/3TC-resistant virus); GW420867X (Glaxo Wellcome); ZIAGEN™ (abacavir/159U89; Glaxo Wellcome Inc.); CS-87 (3'-azido-2',3'-dideoxyuridine; WO 99/66936); and S-acyl-2-thioethyl (SATE)-bearing prodrug forms of β -L-FD4C and β -L-FddC (WO 98/17281).

[1035] Additional NNRTIs include COACTINON™ (Emivirine/MKC-442, potent NNRTI of the HEPT class; Triangle/Abbott); CAPRAVIRINE™ (AG-1549/S-1153, a next generation NNRTI with activity against viruses containing the K103N mutation; Agouron); PNU-142721 (has 20- to 50-fold greater activity than its predecessor delavirdine and is active against K103N mutants; Pharmacia & Upjohn); DPC-961 and DPC-963 (second-generation derivatives of efavirenz, designed to be active against viruses with the K103N mutation; DuPont); GW-420867X (has 25-fold greater activity than HBY097 and is active against K103N mutants; Glaxo Wellcome); CALANOLIDE A (naturally occurring agent from the latex tree; active against viruses containing either or both the Y181C and K103N mutations); and Propolis (WO 99/49830).

[1036] Additional protease inhibitors include LOPINAVIR™ (ABT378/r; Abbott Laboratories); BMS-232632 (an azapeptide; Bristol-Myers Squibb); TIPRANAVIR™ (PNU-140690, a non-peptic dihydropyrone; Pharmacia & Upjohn); PD-178390 (a nonpeptidic dihydropyrone; Parke-Davis); BMS 232632 (an azapeptide; Bristol-Myers Squibb); L-756,423 (an indinavir analog; Merck); DMP-450 (a cyclic urea compound; Avid & DuPont); AG-1776 (a peptidomimetic with *in vitro* activity against protease inhibitor-resistant viruses; Agouron); VX-175/GW-433908 (phosphate prodrug of amprenavir;

Vertex & Glaxo Wellcome); CGP61755 (Ciba); and AGENERASE™ (amprenavir; Glaxo Wellcome Inc.).

[1037] Additional antiretroviral agents include fusion inhibitors/gp41 binders. Fusion inhibitors/gp41 binders include T-20 (a peptide from residues 643-678 of the HIV gp41 transmembrane protein ectodomain which binds to gp41 in its resting state and prevents transformation to the fusogenic state; Trimeris) and T-1249 (a second-generation fusion inhibitor; Trimeris).

[1038] Additional antiretroviral agents include fusion inhibitors/chemokine receptor antagonists. Fusion inhibitors/chemokine receptor antagonists include CXCR4 antagonists such as AMD 3100 (a bicyclam), SDF-1 and its analogs, and ALX40-4C (a cationic peptide), T22 (an 18 amino acid peptide; Trimeris) and the T22 analogs T134 and T140; CCR5 antagonists such as RANTES (9-68), AOP-RANTES, NNY-RANTES, and TAK-779; and CCR5/CXCR4 antagonists such as NSC 651016 (a distamycin analog). Also included are CCR2B, CCR3, and CCR6 antagonists. Chemokine receptor agonists such as RANTES, SDF-1, MIP-1 α , MIP-1 β , etc., may also inhibit fusion.

[1039] Additional antiretroviral agents include integrase inhibitors. Integrase inhibitors include dicaffeoylquinic (DFQA) acids; L-chicoric acid (a dicaffeoyltartaric (DCTA) acid); quinalizarin (QLC) and related anthraquinones; ZINTEVIR™ (AR 177, an oligonucleotide that probably acts at cell surface rather than being a true integrase inhibitor; Arondex); and naphthols such as those disclosed in WO 98/50347.

[1040] Additional antiretroviral agents include hydroxyurea-like compounds such as BCX-34 (a purine nucleoside phosphorylase inhibitor; Biocryst); ribonucleotide reductase inhibitors such as DIDOX™ (Molecules for Health); inosine monophosphate dehydrogenase (IMPDH) inhibitors such as VX-497 (Vertex); and mycopholic acids such as CellCept (mycophenolate mofetil; Roche).

[1041] Additional antiretroviral agents include inhibitors of viral integrase, inhibitors of viral genome nuclear translocation such as arylene bis(methylketone) compounds; inhibitors of HIV entry such as AOP-RANTES, NNY-RANTES, RANTES-IgG fusion protein, soluble complexes of RANTES and glycosaminoglycans (GAG), and AMD-3100; nucleocapsid zinc finger inhibitors such as dithiane compounds; targets of HIV Tat and Rev; and pharmacoenhancers such as ABT-378.

[1042] Other antiretroviral therapies and adjunct therapies include cytokines and lymphokines such as MIP-1 α , MIP-1 β , SDF-1 α , IL-2, PROLEUKIN™ (aldesleukin/L2-

Therapeutics of the invention are used in any combination with TRIMETHOPRIM-SULFAMETHOXAZOLE™, DAPSONE™, PENTAMIDINE™, and/or ATOVAQUONE™ to prophylactically treat or prevent an opportunistic *Pneumocystis carinii* pneumonia infection. In another specific embodiment, Therapeutics of the invention are used in any combination with ISONIAZID™, RIFAMPIN™, PYRAZINAMIDE™, and/or ETHAMBUTOL™ to prophylactically treat or prevent an opportunistic *Mycobacterium avium* complex infection. In another specific embodiment, Therapeutics of the invention are used in any combination with RIFABUTIN™, CLARITHROMYCIN™, and/or AZITHROMYCIN™ to prophylactically treat or prevent an opportunistic *Mycobacterium tuberculosis* infection. In another specific embodiment, Therapeutics of the invention are used in any combination with GANCICLOVIR™, FOSCARNET™, and/or CIDOFOVIR™ to prophylactically treat or prevent an opportunistic cytomegalovirus infection. In another specific embodiment, Therapeutics of the invention are used in any combination with FLUCONAZOLE™, ITRACONAZOLE™, and/or KETOCONAZOLE™ to prophylactically treat or prevent an opportunistic fungal infection. In another specific embodiment, Therapeutics of the invention are used in any combination with ACYCLOVIR™ and/or FAMCICOLVIR™ to prophylactically treat or prevent an opportunistic herpes simplex virus type I and/or type II infection. In another specific embodiment, Therapeutics of the invention are used in any combination with PYRIMETHAMINE™ and/or LEUCOVORIN™ to prophylactically treat or prevent an opportunistic *Toxoplasma gondii* infection. In another specific embodiment, Therapeutics of the invention are used in any combination with LEUCOVORIN™ and/or NEUPOGEN™ to prophylactically treat or prevent an opportunistic bacterial infection.

[1045] In a further embodiment, the Therapeutics of the invention are administered in combination with an antibiotic agent. Antibiotic agents that may be administered with the Therapeutics of the invention include, but are not limited to, amoxicillin, beta-lactamases, aminoglycosides, beta-lactam (glycopeptide), beta-lactamases, Clindamycin, chloramphenicol, cephalosporins, ciprofloxacin, erythromycin, fluoroquinolones, macrolides, metronidazole, penicillins, quinolones, rapamycin, rifampin, streptomycin, sulfonamide, tetracyclines, trimethoprim, trimethoprim-sulfamethoxazole, and vancomycin.

[1046] In other embodiments, the Therapeutics of the invention are administered in combination with immunestimulants. Immunostimulants that may be administered in

combination with the Therapeutics of the invention include, but are not limited to, levamisole (e.g., ERGAMISOL™), isoprinosine (e.g. INOSIPLEX™), interferons (e.g. interferon alpha), and interleukins (e.g., IL-2).

[1047] In other embodiments, Therapeutics of the invention are administered in combination with immunosuppressive agents. Immunosuppressive agents that may be administered in combination with the Therapeutics of the invention include, but are not limited to, steroids, cyclosporine, cyclosporine analogs, cyclophosphamide methylprednisone, prednisone, azathioprine, FK-506, 15-deoxyspergualin, and other immunosuppressive agents that act by suppressing the function of responding T cells. Other immunosuppressive agents that may be administered in combination with the Therapeutics of the invention include, but are not limited to, prednisolone, methotrexate, thalidomide, methoxsalen, rapamycin, leflunomide, mizoribine (BREDININ™), brequinar, deoxyspergualin, and azaspirane (SKF 105685), ORTHOCLONE OKT® 3 (muromonab-CD3), SANDIMMUNE™, NEORAL™, SANGDYA™ (cyclosporine), PROGRAF® (FK506, tacrolimus), CELLCEPT® (mycophenolate mofetil, of which the active metabolite is mycophenolic acid), IMURAN™ (azathioprine), glucocorticosteroids, adrenocortical steroids such as DELTASONE™ (prednisone) and HYDELTRASOL™ (prednisolone), FOLEX™ and MEXATE™ (methotrxate), OXSORALEN-ULTRA™ (methoxsalen) and RAPAMUNE™ (sirolimus). In a specific embodiment, immunosuppressants may be used to prevent rejection of organ or bone marrow transplantation.

[1048] In an additional embodiment, Therapeutics of the invention are administered alone or in combination with one or more intravenous immune globulin preparations. Intravenous immune globulin preparations that may be administered with the Therapeutics of the invention include, but not limited to, GAMMAR™, IVEEGAM™, SANDOGLOBULIN™, GAMMAGARD S/D™, ATGAM™ (antithymocyte globulin), and GAMIMUNE™. In a specific embodiment, Therapeutics of the invention are administered in combination with intravenous immune globulin preparations in transplantation therapy (e.g., bone marrow transplant).

[1049] In certain embodiments, the Therapeutics of the invention are administered alone or in combination with an anti-inflammatory agent. Anti-inflammatory agents that may be administered with the Therapeutics of the invention include, but are not limited to, corticosteroids (e.g. betamethasone, budesonide, cortisone, dexamethasone, hydrocortisone, methylprednisolone, prednisolone, prednisone, and triamcinolone), nonsteroidal anti-

inflammatory drugs (e.g., diclofenac, diflunisal, etodolac, fenoprofen, floctafenine, flurbiprofen, ibuprofen, indomethacin, ketoprofen, meclofenamate, mefenamic acid, meloxicam, nabumetone, naproxen, oxaprozin, phenylbutazone, piroxicam, sulindac, tenoxicam, tiaprofenic acid, and tolmetin.), as well as antihistamines, aminoarylcarboxylic acid derivatives, arylacetic acid derivatives, arylbutyric acid derivatives, arylcarboxylic acids, arylpropionic acid derivatives, pyrazoles, pyrazolones, salicylic acid derivatives, thiazinecarboxamides, e-acetamidocaproic acid, S-adenosylmethionine, 3-amino-4-hydroxybutyric acid, amixetrine, bendazac, benzydamine, bucolome, difenpiramide, ditazol, emorfazone, guaiazulene, nabumetone, nimesulide, orgotein, oxaceprol, paranyline, perisoxal, pifoxime, proquazone, proxazole, and tenidap.

[1050] In an additional embodiment, the compositions of the invention are administered alone or in combination with an anti-angiogenic agent. Anti-angiogenic agents that may be administered with the compositions of the invention include, but are not limited to, Angiostatin (Entremed, Rockville, MD), Troponin-1 (Boston Life Sciences, Boston, MA), anti-Invasive Factor, retinoic acid and derivatives thereof, paclitaxel (Taxol), Suramin, Tissue Inhibitor of Metalloproteinase-1, Tissue Inhibitor of Metalloproteinase-2, VEGI, Plasminogen Activator Inhibitor-1, Plasminogen Activator Inhibitor-2, and various forms of the lighter "d group" transition metals.

[1051] Lighter "d group" transition metals include, for example, vanadium, molybdenum, tungsten, titanium, niobium, and tantalum species. Such transition metal species may form transition metal complexes. Suitable complexes of the above-mentioned transition metal species include oxo transition metal complexes.

[1052] Representative examples of vanadium complexes include oxo vanadium complexes such as vanadate and vanadyl complexes. Suitable vanadate complexes include metavanadate and orthovanadate complexes such as, for example, ammonium metavanadate, sodium metavanadate, and sodium orthovanadate. Suitable vanadyl complexes include, for example, vanadyl acetylacetonate and vanadyl sulfate including vanadyl sulfate hydrates such as vanadyl sulfate mono- and trihydrates.

[1053] Representative examples of tungsten and molybdenum complexes also include oxo complexes. Suitable oxo tungsten complexes include tungstate and tungsten oxide complexes. Suitable tungstate complexes include ammonium tungstate, calcium tungstate, sodium tungstate dihydrate, and tungstic acid. Suitable tungsten oxides include tungsten (IV) oxide and tungsten (VI) oxide. Suitable oxo molybdenum complexes include

09950083-091201

molybdate, molybdenum oxide, and molybdenyl complexes. Suitable molybdate complexes include ammonium molybdate and its hydrates, sodium molybdate and its hydrates, and potassium molybdate and its hydrates. Suitable molybdenum oxides include molybdenum (VI) oxide, molybdenum (VI) oxide, and molybdic acid. Suitable molybdenyl complexes include, for example, molybdenyl acetylacetonate. Other suitable tungsten and molybdenum complexes include hydroxo derivatives derived from, for example, glycerol, tartaric acid, and sugars.

[1054] A wide variety of other anti-angiogenic factors may also be utilized within the context of the present invention. Representative examples include, but are not limited to, platelet factor 4; protamine sulphate; sulphated chitin derivatives (prepared from queen crab shells), (Murata et al., *Cancer Res.* 51:22-26, (1991)); Sulphated Polysaccharide Peptidoglycan Complex (SP- PG) (the function of this compound may be enhanced by the presence of steroids such as estrogen, and tamoxifen citrate); Staurosporine; modulators of matrix metabolism, including for example, proline analogs, cishydroxyproline, d,L-3,4-dehydroproline, Thiaproline, alpha,alpha-dipyridyl, aminopropionitrile fumarate; 4-propyl-5-(4-pyridinyl)-2(3H)-oxazolone; Methotrexate; Mitoxantrone; Heparin; Interferons; 2 Macroglobulin-serum; ChIMP-3 (Pavloff et al., *J. Bio. Chem.* 267:17321-17326, (1992)); Chymostatin (Tomkinson et al., *Biochem J.* 286:475-480, (1992)); Cyclodextrin Tetradecasulfate; Eponemycin; Camptothecin; Fumagillin (Ingber et al., *Nature* 348:555-557, (1990)); Gold Sodium Thiomalate ("GST"; Matsubara and Ziff, *J. Clin. Invest.* 79:1440-1446, (1987)); anticollagenase-serum; alpha2-antiplasmin (Holmes et al., *J. Biol. Chem.* 262(4):1659-1664, (1987)); Bisantrene (National Cancer Institute); Lobenzarit disodium (N-(2)-carboxyphenyl-4- chloroanthronilic acid disodium or "CCA"; (Takeuchi et al., *Agents Actions* 36:312-316, (1992)); and metalloproteinase inhibitors such as BB94.

[1055] Additional anti-angiogenic factors that may also be utilized within the context of the present invention include Thalidomide, (Celgene, Warren, NJ); Angiostatic steroid; AGM-1470 (H. Brem and J. Folkman *J Pediatr. Surg.* 28:445-51 (1993)); an integrin alpha v beta 3 antagonist (C. Storgard et al., *J Clin. Invest.* 103:47-54 (1999)); carboxynaminolmidazole; Carboxyamidotriazole (CAI) (National Cancer Institute, Bethesda, MD); Conbretastatin A-4 (CA4P) (OXiGENE, Boston, MA); Squalamine (Magainin Pharmaceuticals, Plymouth Meeting, PA); TNP-470, (Tap Pharmaceuticals, Deerfield, IL); ZD-0101 AstraZeneca (London, UK); APRA (CT2584); Benefin, Byrostatin-1 (SC339555); CGP-41251 (PKC 412); CM101; Dexrazoxane (ICRF187);

[1058] In a particular embodiment, the use of compositions of the invention in combination with anti-angiogenic agents is contemplated for the treatment, prevention, and/or amelioration of arthritis. In a more particular embodiment, the use of compositions of the invention in combination with anti-angiogenic agents is contemplated for the treatment, prevention, and/or amelioration of rheumatoid arthritis.

[1059] In another embodiment, the polynucleotides encoding a polypeptide of the present invention are administered in combination with an angiogenic protein, or polynucleotides encoding an angiogenic protein. Examples of angiogenic proteins that may be administered with the compositions of the invention include, but are not limited to, acidic and basic fibroblast growth factors, VEGF-1, VEGF-2, VEGF-3, epidermal growth factor alpha and beta, platelet-derived endothelial cell growth factor, platelet-derived growth factor, tumor necrosis factor alpha, hepatocyte growth factor, insulin-like growth factor, colony stimulating factor, macrophage colony stimulating factor, granulocyte/macrophage colony stimulating factor, and nitric oxide synthase.

[1060] In additional embodiments, compositions of the invention are administered in combination with a chemotherapeutic agent. Chemotherapeutic agents that may be administered with the Therapeutics of the invention include, but are not limited to alkylating agents such as nitrogen mustards (for example, Mechlorethamine, cyclophosphamide, Cyclophosphamide Ifosfamide, Melphalan (L-sarcolysin), and Chlorambucil), ethylenimines and methylmelamines (for example, Hexamethylmelamine and Thiotepa), alkyl sulfonates (for example, Busulfan), nitrosoureas (for example, Carmustine (BCNU), Lomustine (CCNU), Semustine (methyl-CCNU), and Streptozocin (streptozotocin)), triazines (for example, Dacarbazine (DTIC; dimethyltriazenoimidazolecarboxamide)), folic acid analogs (for example, Methotrexate (amethopterin)), pyrimidine analogs (for example, Fluorouracil (5-fluorouracil; 5-FU), Floxuridine (fluorodeoxyuridine; FudR), and Cytarabine (cytosine arabinoside)), purine analogs and related inhibitors (for example, Mercaptopurine (6-mercaptopurine; 6-MP), Thioguanine (6-thioguanine; TG), and Pentostatin (2'-deoxycoformycin)), vinca alkaloids (for example, Vinblastine (VLB, vinblastine sulfate)) and Vincristine (vincristine sulfate)), epipodophyllotoxins (for example, Etoposide and Teniposide), antibiotics (for example, Dactinomycin (actinomycin D), Daunorubicin (daunomycin; rubidomycin), Doxorubicin, Bleomycin, Plicamycin (mithramycin), and Mitomycin (mitomycin C), enzymes (for example, L-Asparaginase), biological response modifiers (for example, Interferon-alpha

and interferon-alpha-2b), platinum coordination compounds (for example, Cisplatin (cis-DDP) and Carboplatin), anthracenedione (Mitoxantrone), substituted ureas (for example, Hydroxyurea), methylhydrazine derivatives (for example, Procarbazine (N-methylhydrazine; MIH), adrenocorticosteroids (for example, Prednisone), progestins (for example, Hydroxyprogesterone caproate, Medroxyprogesterone, Medroxyprogesterone acetate, and Megestrol acetate), estrogens (for example, Diethylstilbestrol (DES), Diethylstilbestrol diphosphate, Estradiol, and Ethinyl estradiol), antiestrogens (for example, Tamoxifen), androgens (Testosterone propionate, and Fluoxymesterone), antiandrogens (for example, Flutamide), gonadotropin-releasing hormone analogs (for example, Leuprolide), other hormones and hormone analogs (for example, methyltestosterone, estramustine, estramustine phosphate sodium, chlorotrianisene, and testolactone), and others (for example, dicarbazine, glutamic acid, and mitotane).

[1061] In one embodiment, the compositions of the invention are administered in combination with one or more of the following drugs: infliximab (also known as Remicade™ Centocor, Inc.), Trocade (Roche, RO-32-3555), Leflunomide (also known as Arava™ from Hoechst Marion Roussel), Kineret™ (an IL-1 Receptor antagonist also known as Anakinra from Amgen, Inc.)

[1062] In a specific embodiment, compositions of the invention are administered in combination with CHOP (cyclophosphamide, doxorubicin, vincristine, and prednisone) or combination of one or more of the components of CHOP. In one embodiment, the compositions of the invention are administered in combination with anti-CD20 antibodies, human monoclonal anti-CD20 antibodies. In another embodiment, the compositions of the invention are administered in combination with anti-CD20 antibodies and CHOP, or anti-CD20 antibodies and any combination of one or more of the components of CHOP, particularly cyclophosphamide and/or prednisone. In a specific embodiment, compositions of the invention are administered in combination with Rituximab. In a further embodiment, compositions of the invention are administered with Rituximab and CHOP, or Rituximab and any combination of one or more of the components of CHOP, particularly cyclophosphamide and/or prednisone. In a specific embodiment, compositions of the invention are administered in combination with tositumomab. In a further embodiment, compositions of the invention are administered with tositumomab and CHOP, or tositumomab and any combination of one or more of the components of CHOP, particularly cyclophosphamide and/or prednisone. The anti-CD20 antibodies may optionally be

associated with radioisotopes, toxins or cytotoxic prodrugs.

[1063] In another specific embodiment, the compositions of the invention are administered in combination Zevalin™. In a further embodiment, compositions of the invention are administered with Zevalin™ and CHOP, or Zevalin™ and any combination of one or more of the components of CHOP, particularly cyclophosphamide and/or prednisone. Zevalin™ may be associated with one or more radisotopes. Particularly preferred isotopes are ⁹⁰Y and ¹¹¹In.

[1064] In an additional embodiment, the Therapeutics of the invention are administered in combination with cytokines. Cytokines that may be administered with the Therapeutics of the invention include, but are not limited to, IL2, IL3, IL4, IL5, IL6, IL7, IL10, IL12, IL13, IL15, anti-CD40, CD40L, IFN-gamma and TNF-alpha. In another embodiment, Therapeutics of the invention may be administered with any interleukin, including, but not limited to, IL-1alpha, IL-1beta, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-12, IL-13, IL-14, IL-15, IL-16, IL-17, IL-18, IL-19, IL-20, and IL-21.

[1065] In one embodiment, the Therapeutics of the invention are administered in combination with members of the TNF family. TNF, TNF-related or TNF-like molecules that may be administered with the Therapeutics of the invention include, but are not limited to, soluble forms of TNF-alpha, lymphotoxin-alpha (LT-alpha, also known as TNF-beta), LT-beta (found in complex heterotrimer LT-alpha2-beta), OPGL, FasL, CD27L, CD30L, CD40L, 4-1BBL, DcR3, OX40L, TNF-gamma (International Publication No. WO 96/14328), AIM-I (International Publication No. WO 97/33899), endokine-alpha (International Publication No. WO 98/07880), OPG, and neutrokin-alpha (International Publication No. WO 98/18921, OX40, and nerve growth factor (NGF), and soluble forms of Fas, CD30, CD27, CD40 and 4-IBB, TR2 (International Publication No. WO 96/34095), DR3 (International Publication No. WO 97/33904), DR4 (International Publication No. WO 98/32856), TR5 (International Publication No. WO 98/30693), TRANK, TR9 (International Publication No. WO 98/56892), TR10 (International Publication No. WO 98/54202), 312C2 (International Publication No. WO 98/06842), and TR12, and soluble forms CD154, CD70, and CD153.

[1066] In an additional embodiment, the Therapeutics of the invention are administered in combination with angiogenic proteins. Angiogenic proteins that may be administered with the Therapeutics of the invention include, but are not limited to, Glioma Derived Growth Factor (GDGF), as disclosed in European Patent Number EP-399816; Platelet

Derived Growth Factor-A (PDGF-A), as disclosed in European Patent Number EP-682110; Platelet Derived Growth Factor-B (PDGF-B), as disclosed in European Patent Number EP-282317; Placental Growth Factor (PlGF), as disclosed in International Publication Number WO 92/06194; Placental Growth Factor-2 (PlGF-2), as disclosed in Hauser et al., Growth Factors, 4:259-268 (1993); Vascular Endothelial Growth Factor (VEGF), as disclosed in International Publication Number WO 90/13649; Vascular Endothelial Growth Factor-A (VEGF-A), as disclosed in European Patent Number EP-506477; Vascular Endothelial Growth Factor-2 (VEGF-2), as disclosed in International Publication Number WO 96/39515; Vascular Endothelial Growth Factor B (VEGF-3); Vascular Endothelial Growth Factor B-186 (VEGF-B186), as disclosed in International Publication Number WO 96/26736; Vascular Endothelial Growth Factor-D (VEGF-D), as disclosed in International Publication Number WO 98/02543; Vascular Endothelial Growth Factor-D (VEGF-D), as disclosed in International Publication Number WO 98/07832; and Vascular Endothelial Growth Factor-E (VEGF-E), as disclosed in German Patent Number DE19639601. The above mentioned references are herein incorporated by reference in their entireties.

[1067] In an additional embodiment, the Therapeutics of the invention are administered in combination with Fibroblast Growth Factors. Fibroblast Growth Factors that may be administered with the Therapeutics of the invention include, but are not limited to, FGF-1, FGF-2, FGF-3, FGF-4, FGF-5, FGF-6, FGF-7, FGF-8, FGF-9, FGF-10, FGF-11, FGF-12, FGF-13, FGF-14, and FGF-15.

[1068] In an additional embodiment, the Therapeutics of the invention are administered in combination with hematopoietic growth factors. Hematopoietic growth factors that may be administered with the Therapeutics of the invention include, but are not limited to, granulocyte macrophage colony stimulating factor (GM-CSF) (sargramostim, LEUKINE™, PROKINE™), granulocyte colony stimulating factor (G-CSF) (filgrastim, NEUPOGEN™), macrophage colony stimulating factor (M-CSF, CSF-1) erythropoietin (epoetin alfa, EPOGEN™, PROCRIT™), stem cell factor (SCF, c-kit ligand, steel factor), megakaryocyte colony stimulating factor, PIXY321 (a GMCSF/IL-3 fusion protein), interleukins, especially any one or more of IL-1 through IL-12, interferon-gamma, or thrombopoietin.

[1069] In certain embodiments, Therapeutics of the present invention are administered in combination with adrenergic blockers, such as, for example, acebutolol, atenolol, betaxolol, bisoprolol, carteolol, labetalol, metoprolol, nadolol, oxprenolol, penbutolol, pindolol, propranolol, sotalol, and timolol.

(liothyroine sodium), and THYROLAR™ (liotrix); antithyroid compounds such as 6-*n*-propylthiouracil (propylthiouracil), 1-methyl-2-mercaptoimidazole and TAPAZOLE™ (methimazole), NEO-MERCAZOLE™ (carbimazole); beta-adrenergic receptor antagonists such as propranolol and esmolol; Ca²⁺ channel blockers; dexamethasone and iodinated radiological contrast agents such as TELEPAQUE™ (iopanoic acid) and ORAGRAFIN™ (sodium ipodate).

[1073] Additional treatments for endocrine and/or hormone imbalance disorders include, but are not limited to, estrogens or conjugated estrogens such as ESTRACE™ (estradiol), ESTINYL™ (ethinyl estradiol), PREMARIN™, ESTRATAB™, ORTHO-EST™, OGEN™ and estropipate (estrone), ESTROVIS™ (quinestrol), ESTRADERM™ (estradiol), DELESTROGEN™ and VALERGEN™ (estradiol valerate), DEPO-ESTRADIOL CYPIONATE™ and ESTROJECT LA™ (estradiol cypionate); antiestrogens such as NOLVADEX™ (tamoxifen), SEROPHENE™ and CLOMID™ (clomiphene); progestins such as DURALUTIN™ (hydroxyprogesterone caproate), MPA™ and DEPO-PROVERA™ (medroxyprogesterone acetate), PROVERA™ and CYCRIN™ (MPA), MEGACE™ (megestrol acetate), NORLUTIN™ (norethindrone), and NORLUTATE™ and AYGESTIN™ (norethindrone acetate); progesterone implants such as NORPLANT SYSTEM™ (subdermal implants of norgestrel); antiprogestins such as RU 486™ (mifepristone); hormonal contraceptives such as ENOVID™ (norethynodrel plus mestranol), PROGESTASERT™ (intrauterine device that releases progesterone), LOESTRIN™, BREVICON™, MODICON™, GENORA™, NELONA™, NORINYL™, OVACON-35™ and OVACON-50™ (ethinyl estradiol/norethindrone), LEVLEN™, NORDETTE™, TRI-LEVLEN™ and TRIPHASIL-21™ (ethinyl estradiol/levonorgestrel) LO/OVRAL™ and OVRAL™ (ethinyl estradiol/norgestrel), DEMULEN™ (ethinyl estradiol/ethynodiol diacetate), NORINYL™, ORTHO-NOVUM™, NORETHIN™, GENORA™, and NELOVA™ (norethindrone/mestranol), DESOGEN™ and ORTHO-CEPT™ (ethinyl estradiol/desogestrel), ORTHO-CYCLEN™ and ORTHO-TRICYCLEN™ (ethinyl estradiol/norgestimate), MICRONOR™ and NOR-QD™ (norethindrone), and OVRETTE™ (norgestrel).

[1074] Additional treatments for endocrine and/or hormone imbalance disorders include, but are not limited to, testosterone esters such as methenolone acetate and testosterone

furoate), HALDRONE™ (paramethasone acetate), DELTA-CORTEF™ (prednisolone), ECONOPRED™ (prednisolone acetate), HYDELTRASOL™ (prednisolone sodium phosphate), HYDELTRA-T.B.A™ (prednisolone tebutate), DELTASONE™ (prednisone), ARISTOCORT™ and KENACORT™ (triamcinolone), KENALOG™ (triamcinolone acetonide), ARISTOCORT™ and KENACORT DIACETATE™ (triamcinolone diacetate), and ARISTOSPAN™ (triamcinolone hexacetonide); inhibitors of biosynthesis and action of adrenocortical steroids such as CYTADREN™ (aminoglutethimide), NIZORAL™ (ketoconazole), MODRASTANE™ (trilostane), and METOPIRONE™ (metyrapone); bovine, porcine or human insulin or mixtures thereof; insulin analogs; recombinant human insulin such as HUMULIN™ and NOVOLIN™; oral hypoglycemic agents such as ORAMIDE™ and ORINASE™ (tolbutamide), DIABINESE™ (chlorpropamide), TOLAMIDE™ and TOLINASE™ (tolazamide), DYMELOR™ (acetohexamide), glibenclamide, MICRONASE™, DIBETA™ and GLYNASE™ (glyburide), GLUCOTROL™ (glipizide), and DIAMICRON™ (gliclazide), GLUCOPHAGE™ (metformin), ciglitazone, pioglitazone, and alpha-glucosidase inhibitors; bovine or porcine glucagon; somatostatins such as SANDOSTATIN™ (octreotide); and diazoxides such as PROGLYCEM™ (diazoxide).

[1075] In one embodiment, the Therapeutics of the invention are administered in combination with treatments for uterine motility disorders. Treatments for uterine motility disorders include, but are not limited to, estrogen drugs such as conjugated estrogens (e.g., PREMARIN® and ESTRATAB®), estradiols (e.g., CLIMARA® and ALORA®), estropipate, and chlorotrianisene; progestin drugs (e.g., AMEN® (medroxyprogesterone), MICRONOR® (norethidrone acetate), PROMETRIUM® progesterone, and megestrol acetate); and estrogen/progesterone combination therapies such as, for example, conjugated estrogens/medroxyprogesterone (e.g., PREMPRO™ and PREMPHASE®) and norethindrone acetate/ethinyl estsradiol (e.g., FEMHRT™).

[1076] In an additional embodiment, the Therapeutics of the invention are administered in combination with drugs effective in treating iron deficiency and hypochromic anemias, including but not limited to, ferrous sulfate (iron sulfate, FEOSOL™), ferrous fumarate (e.g., FEOSTAT™), ferrous gluconate (e.g., FERGON™), polysaccharide-iron complex (e.g., NIFEREX™), iron dextran injection (e.g., INFED™), cupric sulfate, pyroxidine, riboflavin, Vitamin B₁₂, cyancobalamin injection (e.g., REDISOL™, RUBRAMIN PC™),

such an individual a composition comprising a therapeutically effective amount of an agonist of the invention (including polypeptides of the invention). Moreover, it will be appreciated that conditions caused by a decrease in the standard or normal expression level of a polypeptide of the present invention in an individual can be treated by administering the agonist or antagonist of the present invention. Thus, the invention also provides a method of treatment of an individual in need of an increased level of the polypeptide comprising administering to such an individual a Therapeutic comprising an amount of the agonist or antagonist to increase the activity level of the polypeptide in such an individual.

[1083] For example, a patient with decreased levels of a polypeptide receives a daily dose 0.1-100 ug/kg of the agonist or antagonist for six consecutive days. The exact details of the dosing scheme, based on administration and formulation, are provided in Example 13.

Example 15: Method of Treating Increased Levels of the Polypeptide

[1084] The present invention also relates to a method of treating an individual in need of a decreased level of a polypeptide of the invention in the body comprising administering to such an individual a composition comprising a therapeutically effective amount of an antagonist of the invention (including polypeptides and antibodies of the invention).

[1085] In one example, antisense technology is used to inhibit production of a polypeptide of the present invention. This technology is one example of a method of decreasing levels of a polypeptide, due to a variety of etiologies, such as cancer.

[1086] For example, a patient diagnosed with abnormally increased levels of a polypeptide is administered intravenously antisense polynucleotides at 0.5, 1.0, 1.5, 2.0 and 3.0 mg/kg day for 21 days. This treatment is repeated after a 7-day rest period if the treatment was well tolerated. The antisense polynucleotides of the present invention can be formulated using techniques and formulations described herein (e.g. see Example 13), or otherwise known in the art.

09950083 091201
102760 0000560

Example 16: Method of Treatment Using Gene Therapy-Ex Vivo

[1087] One method of gene therapy transplants fibroblasts, which are capable of expressing a polypeptide, onto a patient. Generally, fibroblasts are obtained from a subject by skin biopsy. The resulting tissue is placed in tissue-culture medium and separated into small pieces. Small chunks of the tissue are placed on a wet surface of a tissue culture flask, approximately ten pieces are placed in each flask. The flask is turned upside down, closed tight and left at room temperature over night. After 24 hours at room temperature, the flask is inverted and the chunks of tissue remain fixed to the bottom of the flask and fresh media (e.g., Ham's F12 media, with 10% FBS, penicillin and streptomycin) is added. The flasks are then incubated at 37 degree C for approximately one week.

[1088] At this time, fresh media is added and subsequently changed every several days. After an additional two weeks in culture, a monolayer of fibroblasts emerge. The monolayer is trypsinized and scaled into larger flasks.

[1089] pMV-7 (Kirschmeier, P.T. et al., DNA, 7:219-25 (1988)), flanked by the long terminal repeats of the Moloney murine sarcoma virus, is digested with EcoRI and HindIII and subsequently treated with calf intestinal phosphatase. The linear vector is fractionated on agarose gel and purified, using glass beads.

[1090] The cDNA encoding a polypeptide of the present invention can be amplified using PCR primers which correspond to the 5' and 3' end sequences respectively as set forth in Example 1 using primers and having appropriate restriction sites and initiation/stop codons, if necessary. Preferably, the 5' primer contains an EcoRI site and the 3' primer includes a HindIII site. Equal quantities of the Moloney murine sarcoma virus linear backbone and the amplified EcoRI and HindIII fragment are added together, in the presence of T4 DNA ligase. The resulting mixture is maintained under conditions appropriate for ligation of the two fragments. The ligation mixture is then used to transform bacteria HB101, which are then plated onto agar containing kanamycin for the purpose of confirming that the vector has the gene of interest properly inserted.

[1091] The amphotropic pA317 or GP+am12 packaging cells are grown in tissue culture to confluent density in Dulbecco's Modified Eagles Medium (DMEM) with 10% calf serum (CS), penicillin and streptomycin. The MSV vector containing the gene is then added to the media and the packaging cells transduced with the vector. The packaging cells now

09950083-091201

produce infectious viral particles containing the gene (the packaging cells are now referred to as producer cells).

[1092] Fresh media is added to the transduced producer cells, and subsequently, the media is harvested from a 10 cm plate of confluent producer cells. The spent media, containing the infectious viral particles, is filtered through a millipore filter to remove detached producer cells and this media is then used to infect fibroblast cells. Media is removed from a sub-confluent plate of fibroblasts and quickly replaced with the media from the producer cells. This media is removed and replaced with fresh media. If the titer of virus is high, then virtually all fibroblasts will be infected and no selection is required. If the titer is very low, then it is necessary to use a retroviral vector that has a selectable marker, such as neo or his. Once the fibroblasts have been efficiently infected, the fibroblasts are analyzed to determine whether protein is produced.

[1093] The engineered fibroblasts are then transplanted onto the host, either alone or after having been grown to confluence on cytodex 3 microcarrier beads.

***Example 17: Gene Therapy Using Endogenous Genes Corresponding To
Polynucleotides of the Invention***

[1094] Another method of gene therapy according to the present invention involves operably associating the endogenous polynucleotide sequence of the invention with a promoter via homologous recombination as described, for example, in U.S. Patent NO: 5,641,670, issued June 24, 1997; International Publication NO: WO 96/29411, published September 26, 1996; International Publication NO: WO 94/12650, published August 4, 1994; Koller et al., *Proc. Natl. Acad. Sci. USA*, 86:8932-8935 (1989); and Zijlstra et al., *Nature*, 342:435-438 (1989). This method involves the activation of a gene which is present in the target cells, but which is not expressed in the cells, or is expressed at a lower level than desired.

[1095] Polynucleotide constructs are made which contain a promoter and targeting sequences, which are homologous to the 5' non-coding sequence of endogenous polynucleotide sequence, flanking the promoter. The targeting sequence will be sufficiently near the 5' end of the polynucleotide sequence so the promoter will be operably linked to the endogenous sequence upon homologous recombination. The promoter and the

targeting sequences can be amplified using PCR. Preferably, the amplified promoter contains distinct restriction enzyme sites on the 5' and 3' ends. Preferably, the 3' end of the first targeting sequence contains the same restriction enzyme site as the 5' end of the amplified promoter and the 5' end of the second targeting sequence contains the same restriction site as the 3' end of the amplified promoter.

[1096] The amplified promoter and the amplified targeting sequences are digested with the appropriate restriction enzymes and subsequently treated with calf intestinal phosphatase. The digested promoter and digested targeting sequences are added together in the presence of T4 DNA ligase. The resulting mixture is maintained under conditions appropriate for ligation of the two fragments. The construct is size fractionated on an agarose gel, then purified by phenol extraction and ethanol precipitation.

[1097] In this Example, the polynucleotide constructs are administered as naked polynucleotides via electroporation. However, the polynucleotide constructs may also be administered with transfection-facilitating agents, such as liposomes, viral sequences, viral particles, precipitating agents, etc. Such methods of delivery are known in the art.

[1098] Once the cells are transfected, homologous recombination will take place which results in the promoter being operably linked to the endogenous polynucleotide sequence. This results in the expression of polynucleotide corresponding to the polynucleotide in the cell. Expression may be detected by immunological staining, or any other method known in the art.

[1099] Fibroblasts are obtained from a subject by skin biopsy. The resulting tissue is placed in DMEM + 10% fetal calf serum. Exponentially growing or early stationary phase fibroblasts are trypsinized and rinsed from the plastic surface with nutrient medium. An aliquot of the cell suspension is removed for counting, and the remaining cells are subjected to centrifugation. The supernatant is aspirated and the pellet is resuspended in 5 ml of electroporation buffer (20 mM HEPES pH 7.3, 137 mM NaCl, 5 mM KCl, 0.7 mM Na₂HPO₄, 6 mM dextrose). The cells are recentrifuged, the supernatant aspirated, and the cells resuspended in electroporation buffer containing 1 mg/ml acetylated bovine serum albumin. The final cell suspension contains approximately 3×10^6 cells/ml. Electroporation should be performed immediately following resuspension.

[1100] Plasmid DNA is prepared according to standard techniques. For example, to construct a plasmid for targeting to the locus corresponding to the polynucleotide of the invention, plasmid pUC18 (MBI Fermentas, Amherst, NY) is digested with HindIII. The

polynucleotide of the present invention may be operatively linked to (i.e., associated with) a promoter or any other genetic elements necessary for the expression of the polypeptide by the target tissue. Such gene therapy and delivery techniques and methods are known in the art, see, for example, WO90/11092, WO98/11779; U.S. Patent NO. 5693622, 5705151, 5580859; Tabata et al., *Cardiovasc. Res.* 35(3):470-479 (1997); Chao et al., *Pharmacol. Res.* 35(6):517-522 (1997); Wolff, *Neuromuscul. Disord.* 7(5):314-318 (1997); Schwartz et al., *Gene Ther.* 3(5):405-411 (1996); Tsurumi et al., *Circulation* 94(12):3281-3290 (1996) (incorporated herein by reference).

[1105] The polynucleotide constructs may be delivered by any method that delivers injectable materials to the cells of an animal, such as, injection into the interstitial space of tissues (heart, muscle, skin, lung, liver, intestine and the like). The polynucleotide constructs can be delivered in a pharmaceutically acceptable liquid or aqueous carrier.

[1106] The term "naked" polynucleotide, DNA or RNA, refers to sequences that are free from any delivery vehicle that acts to assist, promote, or facilitate entry into the cell, including viral sequences, viral particles, liposome formulations, lipofectin or precipitating agents and the like. However, the polynucleotides of the present invention may also be delivered in liposome formulations (such as those taught in Felgner P.L. et al. (1995) *Ann. NY Acad. Sci.* 772:126-139 and Abdallah B. et al. (1995) *Biol. Cell* 85(1):1-7) which can be prepared by methods well known to those skilled in the art.

[1107] The polynucleotide vector constructs used in the gene therapy method are preferably constructs that will not integrate into the host genome nor will they contain sequences that allow for replication. Any strong promoter known to those skilled in the art can be used for driving the expression of DNA. Unlike other gene therapy techniques, one major advantage of introducing naked nucleic acid sequences into target cells is the transitory nature of the polynucleotide synthesis in the cells. Studies have shown that non-replicating DNA sequences can be introduced into cells to provide production of the desired polypeptide for periods of up to six months.

[1108] The polynucleotide construct can be delivered to the interstitial space of tissues within an animal, including muscle, skin, brain, lung, liver, spleen, bone marrow, thymus, heart, lymph, blood, bone, cartilage, pancreas, kidney, gall bladder, stomach, intestine, testis, ovary, uterus, rectum, nervous system, eye, gland, and connective tissue. Interstitial space of the tissues comprises the intercellular fluid, mucopolysaccharide matrix among the reticular fibers of organ tissues, elastic fibers in the walls of vessels or chambers, collagen

fibers of fibrous tissues, or that same matrix within connective tissue ensheathing muscle cells or in the lacunae of bone. It is similarly the space occupied by the plasma of the circulation and the lymph fluid of the lymphatic channels. Delivery to the interstitial space of muscle tissue is preferred for the reasons discussed below. They may be conveniently delivered by injection into the tissues comprising these cells. They are preferably delivered to and expressed in persistent, non-dividing cells which are differentiated, although delivery and expression may be achieved in non-differentiated or less completely differentiated cells, such as, for example, stem cells of blood or skin fibroblasts. *In vivo* muscle cells are particularly competent in their ability to take up and express polynucleotides.

[1109] For the naked polynucleotide injection, an effective dosage amount of DNA or RNA will be in the range of from about 0.05 g/kg body weight to about 50 mg/kg body weight. Preferably the dosage will be from about 0.005 mg/kg to about 20 mg/kg and more preferably from about 0.05 mg/kg to about 5 mg/kg. Of course, as the artisan of ordinary skill will appreciate, this dosage will vary according to the tissue site of injection. The appropriate and effective dosage of nucleic acid sequence can readily be determined by those of ordinary skill in the art and may depend on the condition being treated and the route of administration. The preferred route of administration is by the parenteral route of injection into the interstitial space of tissues. However, other parenteral routes may also be used, such as, inhalation of an aerosol formulation particularly for delivery to lungs or bronchial tissues, throat or mucous membranes of the nose. In addition, naked polynucleotide constructs can be delivered to arteries during angioplasty by the catheter used in the procedure.

[1110] The dose response effects of injected polynucleotide in muscle *in vivo* is determined as follows. Suitable template DNA for production of mRNA coding for polypeptide of the present invention is prepared in accordance with a standard recombinant DNA methodology. The template DNA, which may be either circular or linear, is either used as naked DNA or complexed with liposomes. The quadriceps muscles of mice are then injected with various amounts of the template DNA.

[1111] Five to six week old female and male Balb/C mice are anesthetized by intraperitoneal injection with 0.3 ml of 2.5% Avertin. A 1.5 cm incision is made on the anterior thigh, and the quadriceps muscle is directly visualized. The template DNA is injected in 0.1 ml of carrier in a 1 cc syringe through a 27 gauge needle over one minute, approximately 0.5 cm from the distal insertion site of the muscle into the knee and about 0.2

cm deep. A suture is placed over the injection site for future localization, and the skin is closed with stainless steel clips.

[1112] After an appropriate incubation time (e.g., 7 days) muscle extracts are prepared by excising the entire quadriceps. Every fifth 15 um cross-section of the individual quadriceps muscles is histochemically stained for protein expression. A time course for protein expression may be done in a similar fashion except that quadriceps from different mice are harvested at different times. Persistence of DNA in muscle following injection may be determined by Southern blot analysis after preparing total cellular DNA and HIRT supernatants from injected and control mice. The results of the above experimentation in mice can be used to extrapolate proper dosages and other treatment parameters in humans and other animals using naked DNA.

Example 19: Transgenic Animals

[1113] The polypeptides of the invention can also be expressed in transgenic animals. Animals of any species, including, but not limited to, mice, rats, rabbits, hamsters, guinea pigs, pigs, micro-pigs, goats, sheep, cows and non-human primates, e.g., baboons, monkeys, and chimpanzees may be used to generate transgenic animals. In a specific embodiment, techniques described herein or otherwise known in the art, are used to express polypeptides of the invention in humans, as part of a gene therapy protocol.

[1114] Any technique known in the art may be used to introduce the transgene (i.e., polynucleotides of the invention) into animals to produce the founder lines of transgenic animals. Such techniques include, but are not limited to, pronuclear microinjection (Paterson et al., Appl. Microbiol. Biotechnol. 40:691-698 (1994); Carver et al., Biotechnology (NY) 11:1263-1270 (1993); Wright et al., Biotechnology (NY) 9:830-834 (1991); and Hoppe et al., U.S. Pat. No. 4,873,191 (1989)); retrovirus mediated gene transfer into germ lines (Van der Putten et al., Proc. Natl. Acad. Sci., USA 82:6148-6152 (1985)), blastocysts or embryos; gene targeting in embryonic stem cells (Thompson et al., Cell 56:313-321 (1989)); electroporation of cells or embryos (Lo, 1983, Mol Cell. Biol. 3:1803-1814 (1983)); introduction of the polynucleotides of the invention using a gene gun (see, e.g., Ulmer et al., Science 259:1745 (1993); introducing nucleic acid constructs into embryonic pluripotent stem cells and transferring the stem cells back into the blastocyst; and sperm-mediated gene transfer (Lavitrano et al., Cell 57:717-723 (1989); etc. For a

review of such techniques, see Gordon, "Transgenic Animals," Intl. Rev. Cytol. 115:171-229 (1989), which is incorporated by reference herein in its entirety.

[1115] Any technique known in the art may be used to produce transgenic clones containing polynucleotides of the invention, for example, nuclear transfer into enucleated oocytes of nuclei from cultured embryonic, fetal, or adult cells induced to quiescence (Campell et al., Nature 380:64-66 (1996); Wilmut et al., Nature 385:810-813 (1997)).

[1116] The present invention provides for transgenic animals that carry the transgene in all their cells, as well as animals which carry the transgene in some, but not all their cells, *i.e.*, mosaic animals or chimeric. The transgene may be integrated as a single transgene or as multiple copies such as in concatamers, *e.g.*, head-to-head tandems or head-to-tail tandems. The transgene may also be selectively introduced into and activated in a particular cell type by following, for example, the teaching of Lasko et al. (Lasko et al., Proc. Natl. Acad. Sci. USA 89:6232-6236 (1992)). The regulatory sequences required for such a cell-type specific activation will depend upon the particular cell type of interest, and will be apparent to those of skill in the art. When it is desired that the polynucleotide transgene be integrated into the chromosomal site of the endogenous gene, gene targeting is preferred. Briefly, when such a technique is to be utilized, vectors containing some nucleotide sequences homologous to the endogenous gene are designed for the purpose of integrating, via homologous recombination with chromosomal sequences, into and disrupting the function of the nucleotide sequence of the endogenous gene. The transgene may also be selectively introduced into a particular cell type, thus inactivating the endogenous gene in only that cell type, by following, for example, the teaching of Gu et al. (Gu et al., Science 265:103-106 (1994)). The regulatory sequences required for such a cell-type specific inactivation will depend upon the particular cell type of interest, and will be apparent to those of skill in the art.

[1117] Once transgenic animals have been generated, the expression of the recombinant gene may be assayed utilizing standard techniques. Initial screening may be accomplished by Southern blot analysis or PCR techniques to analyze animal tissues to verify that integration of the transgene has taken place. The level of mRNA expression of the transgene in the tissues of the transgenic animals may also be assessed using techniques which include, but are not limited to, Northern blot analysis of tissue samples obtained from the animal, *in situ* hybridization analysis, and reverse transcriptase-PCR (rt-PCR). Samples

09950087-091201

of transgenic gene-expressing tissue may also be evaluated immunocytochemically or immunohistochemically using antibodies specific for the transgene product.

[1118] Once the founder animals are produced, they may be bred, inbred, outbred, or crossbred to produce colonies of the particular animal. Examples of such breeding strategies include, but are not limited to: outbreeding of founder animals with more than one integration site in order to establish separate lines; inbreeding of separate lines in order to produce compound transgenics that express the transgene at higher levels because of the effects of additive expression of each transgene; crossing of heterozygous transgenic animals to produce animals homozygous for a given integration site in order to both augment expression and eliminate the need for screening of animals by DNA analysis; crossing of separate homozygous lines to produce compound heterozygous or homozygous lines; and breeding to place the transgene on a distinct background that is appropriate for an experimental model of interest.

[1119] Transgenic animals of the invention have uses which include, but are not limited to, animal model systems useful in elaborating the biological function of polypeptides of the present invention, studying conditions and/or disorders associated with aberrant expression, and in screening for compounds effective in ameliorating such conditions and/or disorders.

Example 20: Knock-Out Animals

[1120] Endogenous gene expression can also be reduced by inactivating or "knocking out" the gene and/or its promoter using targeted homologous recombination. (e.g., see Smithies et al., *Nature* 317:230-234 (1985); Thomas & Capecchi, *Cell* 51:503-512 (1987); Thompson et al., *Cell* 5:313-321 (1989); each of which is incorporated by reference herein in its entirety). For example, a mutant, non-functional polynucleotide of the invention (or a completely unrelated DNA sequence) flanked by DNA homologous to the endogenous polynucleotide sequence (either the coding regions or regulatory regions of the gene) can be used, with or without a selectable marker and/or a negative selectable marker, to transfect cells that express polypeptides of the invention *in vivo*. In another embodiment, techniques known in the art are used to generate knockouts in cells that contain, but do not express the gene of interest. Insertion of the DNA construct, via targeted homologous recombination, results in inactivation of the targeted gene. Such approaches are particularly suited in research and agricultural fields where modifications to embryonic stem cells can be used to

generate animal offspring with an inactive targeted gene (e.g., see Thomas & Capecchi 1987 and Thompson 1989, *supra*). However this approach can be routinely adapted for use in humans provided the recombinant DNA constructs are directly administered or targeted to the required site *in vivo* using appropriate viral vectors that will be apparent to those of skill in the art.

[1121] In further embodiments of the invention, cells that are genetically engineered to express the polypeptides of the invention, or alternatively, that are genetically engineered not to express the polypeptides of the invention (e.g., knockouts) are administered to a patient *in vivo*. Such cells may be obtained from the patient (i.e., animal, including human) or an MHC compatible donor and can include, but are not limited to fibroblasts, bone marrow cells, blood cells (e.g., lymphocytes), adipocytes, muscle cells, endothelial cells etc. The cells are genetically engineered *in vitro* using recombinant DNA techniques to introduce the coding sequence of polypeptides of the invention into the cells, or alternatively, to disrupt the coding sequence and/or endogenous regulatory sequence associated with the polypeptides of the invention, e.g., by transduction (using viral vectors, and preferably vectors that integrate the transgene into the cell genome) or transfection procedures, including, but not limited to, the use of plasmids, cosmids, YACs, naked DNA, electroporation, liposomes, etc. The coding sequence of the polypeptides of the invention can be placed under the control of a strong constitutive or inducible promoter or promoter/enhancer to achieve expression, and preferably secretion, of the polypeptides of the invention. The engineered cells which express and preferably secrete the polypeptides of the invention can be introduced into the patient systemically, e.g., in the circulation, or intraperitoneally.

[1122] Alternatively, the cells can be incorporated into a matrix and implanted in the body, e.g., genetically engineered fibroblasts can be implanted as part of a skin graft; genetically engineered endothelial cells can be implanted as part of a lymphatic or vascular graft. (See, for example, Anderson et al. U.S. Patent No. 5,399,349; and Mulligan & Wilson, U.S. Patent No. 5,460,959 each of which is incorporated by reference herein in its entirety).

[1123] When the cells to be administered are non-autologous or non-MHC compatible cells, they can be administered using well known techniques which prevent the development of a host immune response against the introduced cells. For example, the cells may be introduced in an encapsulated form which, while allowing for an exchange of

components with the immediate extracellular environment, does not allow the introduced cells to be recognized by the host immune system.

[1124] Transgenic and "knock-out" animals of the invention have uses which include, but are not limited to, animal model systems useful in elaborating the biological function of polypeptides of the present invention, studying conditions and/or disorders associated with aberrant expression, and in screening for compounds effective in ameliorating such conditions and/or disorders.

***Example 21: Assays Detecting Stimulation or Inhibition of B cell
Proliferation and Differentiation***

[1125] Generation of functional humoral immune responses requires both soluble and cognate signaling between B-lineage cells and their microenvironment. Signals may impart a positive stimulus that allows a B-lineage cell to continue its programmed development, or a negative stimulus that instructs the cell to arrest its current developmental pathway. To date, numerous stimulatory and inhibitory signals have been found to influence B cell responsiveness including IL-2, IL-4, IL-5, IL-6, IL-7, IL10, IL-13, IL-14 and IL-15. Interestingly, these signals are by themselves weak effectors but can, in combination with various co-stimulatory proteins, induce activation, proliferation, differentiation, homing, tolerance and death among B cell populations.

[1126] One of the best studied classes of B-cell co-stimulatory proteins is the TNF-superfamily. Within this family CD40, CD27, and CD30 along with their respective ligands CD154, CD70, and CD153 have been found to regulate a variety of immune responses. Assays which allow for the detection and/or observation of the proliferation and differentiation of these B-cell populations and their precursors are valuable tools in determining the effects various proteins may have on these B-cell populations in terms of proliferation and differentiation. Listed below are two assays designed to allow for the detection of the differentiation, proliferation, or inhibition of B-cell populations and their precursors.

[1127] In Vitro Assay- Agonists or antagonists of the invention can be assessed for its ability to induce activation, proliferation, differentiation or inhibition and/or death in B-cell populations and their precursors. The activity of the agonists or antagonists of the invention on purified human tonsillar B cells, measured qualitatively over the dose range from 0.1 to

[1132] The studies described in this example tested activity of agonists or antagonists of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides or polypeptides of the invention (e.g., gene therapy).

Example 22: T Cell Proliferation Assay

[1133] A CD3-induced proliferation assay is performed on PBMCs and is measured by the uptake of ³H-thymidine. The assay is performed as follows. Ninety-six well plates are coated with 100 µl/well of mAb to CD3 (HIT3a, Pharmingen) or isotype-matched control mAb (B33.1) overnight at 4 degrees C (1 µg/ml in .05M bicarbonate buffer, pH 9.5), then washed three times with PBS. PBMC are isolated by F/H gradient centrifugation from human peripheral blood and added to quadruplicate wells (5 x 10⁴/well) of mAb coated plates in RPMI containing 10% FCS and P/S in the presence of varying concentrations of agonists or antagonists of the invention (total volume 200 µl). Relevant protein buffer and medium alone are controls. After 48 hr. culture at 37 degrees C, plates are spun for 2 min. at 1000 rpm and 100 µl of supernatant is removed and stored -20 degrees C for measurement of IL-2 (or other cytokines) if effect on proliferation is observed. Wells are supplemented with 100 µl of medium containing 0.5 uCi of ³H-thymidine and cultured at 37 degrees C for 18-24 hr. Wells are harvested and incorporation of ³H-thymidine used as a measure of proliferation. Anti-CD3 alone is the positive control for proliferation. IL-2 (100 U/ml) is also used as a control which enhances proliferation. Control antibody which does not induce proliferation of T cells is used as the negative control for the effects of agonists or antagonists of the invention.

[1134] The studies described in this example tested activity of agonists or antagonists of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides or polypeptides of the invention (e.g., gene therapy).

0950083-091201
T02760-09005660

Example 23: Effect of Agonists or Antagonists of the Invention on the Expression of MHC Class II, Costimulatory and Adhesion Molecules and Cell Differentiation of Monocytes and Monocyte-Derived Human Dendritic Cells

[1135] Dendritic cells are generated by the expansion of proliferating precursors found in the peripheral blood: adherent PBMC or elutriated monocytic fractions are cultured for 7-10 days with GM-CSF (50 ng/ml) and IL-4 (20 ng/ml). These dendritic cells have the characteristic phenotype of immature cells (expression of CD1, CD80, CD86, CD40 and MHC class II antigens). Treatment with activating factors, such as TNF- α , causes a rapid change in surface phenotype (increased expression of MHC class I and II, costimulatory and adhesion molecules, downregulation of FC γ RII, upregulation of CD83). These changes correlate with increased antigen-presenting capacity and with functional maturation of the dendritic cells.

[1136] FACS analysis of surface antigens is performed as follows. Cells are treated 1-3 days with increasing concentrations of agonist or antagonist of the invention or LPS (positive control), washed with PBS containing 1% BSA and 0.02 mM sodium azide, and then incubated with 1:20 dilution of appropriate FITC- or PE-labeled monoclonal antibodies for 30 minutes at 4 degrees C. After an additional wash, the labeled cells are analyzed by flow cytometry on a FACScan (Becton Dickinson).

[1137] Effect on the production of cytokines. Cytokines generated by dendritic cells, in particular IL-12, are important in the initiation of T-cell dependent immune responses. IL-12 strongly influences the development of Th1 helper T-cell immune response, and induces cytotoxic T and NK cell function. An ELISA is used to measure the IL-12 release as follows. Dendritic cells (10^6 /ml) are treated with increasing concentrations of agonists or antagonists of the invention for 24 hours. LPS (100 ng/ml) is added to the cell culture as positive control. Supernatants from the cell cultures are then collected and analyzed for IL-12 content using commercial ELISA kit (e.g., R & D Systems (Minneapolis, MN)). The standard protocols provided with the kits are used.

[1138] Effect on the expression of MHC Class II, costimulatory and adhesion molecules. Three major families of cell surface antigens can be identified on monocytes: adhesion

molecules, molecules involved in antigen presentation, and Fc receptor. Modulation of the expression of MHC class II antigens and other costimulatory molecules, such as B7 and ICAM-1, may result in changes in the antigen presenting capacity of monocytes and ability to induce T cell activation. Increased expression of Fc receptors may correlate with improved monocyte cytotoxic activity, cytokine release and phagocytosis.

[1139] FACS analysis is used to examine the surface antigens as follows. Monocytes are treated 1-5 days with increasing concentrations of agonists or antagonists of the invention or LPS (positive control), washed with PBS containing 1% BSA and 0.02 mM sodium azide, and then incubated with 1:20 dilution of appropriate FITC- or PE-labeled monoclonal antibodies for 30 minutes at 4 degrees C. After an additional wash, the labeled cells are analyzed by flow cytometry on a FACScan (Becton Dickinson).

[1140] Monocyte activation and/or increased survival. Assays for molecules that activate (or alternatively, inactivate) monocytes and/or increase monocyte survival (or alternatively, decrease monocyte survival) are known in the art and may routinely be applied to determine whether a molecule of the invention functions as an inhibitor or activator of monocytes. Agonists or antagonists of the invention can be screened using the three assays described below. For each of these assays, Peripheral blood mononuclear cells (PBMC) are purified from single donor leukopacks (American Red Cross, Baltimore, MD) by centrifugation through a Histopaque gradient (Sigma). Monocytes are isolated from PBMC by counterflow centrifugal elutriation.

[1141] Monocyte Survival Assay. Human peripheral blood monocytes progressively lose viability when cultured in absence of serum or other stimuli. Their death results from internally regulated processes (apoptosis). Addition to the culture of activating factors, such as TNF-alpha dramatically improves cell survival and prevents DNA fragmentation. Propidium iodide (PI) staining is used to measure apoptosis as follows. Monocytes are cultured for 48 hours in polypropylene tubes in serum-free medium (positive control), in the presence of 100 ng/ml TNF-alpha (negative control), and in the presence of varying concentrations of the compound to be tested. Cells are suspended at a concentration of 2×10^6 /ml in PBS containing PI at a final concentration of 5 μ g/ml, and then incubated at room temperature for 5 minutes before FACScan analysis. PI uptake has been demonstrated to correlate with DNA fragmentation in this experimental paradigm.

[1142] Effect on cytokine release. An important function of monocytes/macrophages is their regulatory activity on other cellular populations of the immune system through the release of cytokines after stimulation. An ELISA to measure cytokine release is performed as follows. Human monocytes are incubated at a density of 5×10^5 cells/ml with increasing concentrations of agonists or antagonists of the invention and under the same conditions, but in the absence of agonists or antagonists. For IL-12 production, the cells are primed overnight with IFN (100 U/ml) in the presence of agonist or antagonist of the invention. LPS (10 ng/ml) is then added. Conditioned media are collected after 24h and kept frozen until use. Measurement of TNF-alpha, IL-10, MCP-1 and IL-8 is then performed using a commercially available ELISA kit (e.g., R & D Systems (Minneapolis, MN)) and applying the standard protocols provided with the kit.

[1143] Oxidative burst. Purified monocytes are plated in 96-w plate at 2×10^5 cell/well. Increasing concentrations of agonists or antagonists of the invention are added to the wells in a total volume of 0.2 ml culture medium (RPMI 1640 + 10% FCS, glutamine and antibiotics). After 3 days incubation, the plates are centrifuged and the medium is removed from the wells. To the macrophage monolayers, 0.2 ml per well of phenol red solution (140 mM NaCl, 10 mM potassium phosphate buffer pH 7.0, 5.5 mM dextrose, 0.56 mM phenol red and 19 U/ml of HRPO) is added, together with the stimulant (200 nM PMA). The plates are incubated at 37°C for 2 hours and the reaction is stopped by adding 20 μ l 1N NaOH per well. The absorbance is read at 610 nm. To calculate the amount of H_2O_2 produced by the macrophages, a standard curve of a H_2O_2 solution of known molarity is performed for each experiment.

[1144] The studies described in this example tested activity of agonists or antagonists of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides or polypeptides of the invention (e.g., gene therapy).

Example 24: Biological Effects of Agonists or Antagonists of the Invention

Astrocyte and Neuronal Assays

[1145] Agonists or antagonists of the invention, expressed in *Escherichia coli* and purified as described above, can be tested for activity in promoting the survival, neurite outgrowth, or phenotypic differentiation of cortical neuronal cells and for inducing the proliferation of glial fibrillary acidic protein immunopositive cells, astrocytes. The selection of cortical cells for the bioassay is based on the prevalent expression of FGF-1 and FGF-2 in cortical structures and on the previously reported enhancement of cortical neuronal survival resulting from FGF-2 treatment. A thymidine incorporation assay, for example, can be used to elucidate an agonist or antagonist of the invention's activity on these cells.

[1146] Moreover, previous reports describing the biological effects of FGF-2 (basic FGF) on cortical or hippocampal neurons *in vitro* have demonstrated increases in both neuron survival and neurite outgrowth (Walicke et al., "Fibroblast growth factor promotes survival of dissociated hippocampal neurons and enhances neurite extension." *Proc. Natl. Acad. Sci. USA* 83:3012-3016. (1986), assay herein incorporated by reference in its entirety). However, reports from experiments done on PC-12 cells suggest that these two responses are not necessarily synonymous and may depend on not only which FGF is being tested but also on which receptor(s) are expressed on the target cells. Using the primary cortical neuronal culture paradigm, the ability of an agonist or antagonist of the invention to induce neurite outgrowth can be compared to the response achieved with FGF-2 using, for example, a thymidine incorporation assay.

Fibroblast and endothelial cell assays

[1147] Human lung fibroblasts are obtained from Clonetics (San Diego, CA) and maintained in growth media from Clonetics. Dermal microvascular endothelial cells are obtained from Cell Applications (San Diego, CA). For proliferation assays, the human lung fibroblasts and dermal microvascular endothelial cells can be cultured at 5,000 cells/well in a 96-well plate for one day in growth medium. The cells are then incubated for one day in 0.1% BSA basal medium. After replacing the medium with fresh 0.1% BSA medium, the cells are incubated with the test

proteins for 3 days. Alamar Blue (Alamar Biosciences, Sacramento, CA) is added to each well to a final concentration of 10%. The cells are incubated for 4 hr. Cell viability is measured by reading in a CytoFluor fluorescence reader. For the PGE₂ assays, the human lung fibroblasts are cultured at 5,000 cells/well in a 96-well plate for one day. After a medium change to 0.1% BSA basal medium, the cells are incubated with FGF-2 or agonists or antagonists of the invention with or without IL-1 α for 24 hours. The supernatants are collected and assayed for PGE₂ by EIA kit (Cayman, Ann Arbor, MI). For the IL-6 assays, the human lung fibroblasts are cultured at 5,000 cells/well in a 96-well plate for one day. After a medium change to 0.1% BSA basal medium, the cells are incubated with FGF-2 or with or without agonists or antagonists of the invention IL-1 α for 24 hours. The supernatants are collected and assayed for IL-6 by ELISA kit (Endogen, Cambridge, MA).

[1148] Human lung fibroblasts are cultured with FGF-2 or agonists or antagonists of the invention for 3 days in basal medium before the addition of Alamar Blue to assess effects on growth of the fibroblasts. FGF-2 should show a stimulation at 10 - 2500 ng/ml which can be used to compare stimulation with agonists or antagonists of the invention.

Parkinson Models.

[1149] The loss of motor function in Parkinson's disease is attributed to a deficiency of striatal dopamine resulting from the degeneration of the nigrostriatal dopaminergic projection neurons. An animal model for Parkinson's that has been extensively characterized involves the systemic administration of 1-methyl-4 phenyl 1,2,3,6-tetrahydropyridine (MPTP). In the CNS, MPTP is taken-up by astrocytes and catabolized by monoamine oxidase B to 1-methyl-4-phenyl pyridine (MPP⁺) and released. Subsequently, MPP⁺ is actively accumulated in dopaminergic neurons by the high-affinity reuptake transporter for dopamine. MPP⁺ is then concentrated in mitochondria by the electrochemical gradient and selectively inhibits nicotinamide adenine disphosphate: ubiquinone oxidoreductionase (complex I), thereby interfering with electron transport and eventually generating oxygen radicals.

[1150] It has been demonstrated in tissue culture paradigms that FGF-2 (basic FGF) has trophic activity towards nigral dopaminergic neurons (Ferrari et al., Dev. Biol. 1989). Recently, Dr. Unsicker's group has demonstrated that administering FGF-2 in gel foam implants in the striatum results in the near complete protection of nigral dopaminergic neurons from the toxicity associated with MPTP exposure (Otto and Unsicker, J. Neuroscience, 1990).

[1151] Based on the data with FGF-2, agonists or antagonists of the invention can be evaluated to determine whether it has an action similar to that of FGF-2 in enhancing dopaminergic neuronal survival *in vitro* and it can also be tested *in vivo* for protection of dopaminergic neurons in the striatum from the damage associated with MPTP treatment. The potential effect of an agonist or antagonist of the invention is first examined *in vitro* in a dopaminergic neuronal cell culture paradigm. The cultures are prepared by dissecting the midbrain floor plate from gestation day 14 Wistar rat embryos. The tissue is dissociated with trypsin and seeded at a density of 200,000 cells/cm² on polyorthinine-laminin coated glass coverslips. The cells are maintained in Dulbecco's Modified Eagle's medium and F12 medium containing hormonal supplements (N1). The cultures are fixed with paraformaldehyde after 8 days *in vitro* and are processed for tyrosine hydroxylase, a specific marker for dopaminergic neurons, immunohistochemical staining. Dissociated cell cultures are prepared from embryonic rats. The culture medium is changed every third day and the factors are also added at that time.

[1152] Since the dopaminergic neurons are isolated from animals at gestation day 14, a developmental time which is past the stage when the dopaminergic precursor cells are proliferating, an increase in the number of tyrosine hydroxylase immunopositive neurons would represent an increase in the number of dopaminergic neurons surviving *in vitro*. Therefore, if an agonist or antagonist of the invention acts to prolong the survival of dopaminergic neurons, it would suggest that the agonist or antagonist may be involved in Parkinson's Disease.

[1153] The studies described in this example tested activity of agonists or antagonists of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides or polypeptides of the invention (e.g., gene therapy).

Example 25: The Effect of Agonists or Antagonists of the Invention on the Growth of Vascular Endothelial Cells

[1154] On day 1, human umbilical vein endothelial cells (HUVEC) are seeded at $2-5 \times 10^4$ cells/35 mm dish density in M199 medium containing 4% fetal bovine serum (FBS), 16 units/ml heparin, and 50 units/ml endothelial cell growth supplements (ECGS, Biotechnology, Inc.). On day 2, the medium is replaced with M199 containing 10% FBS, 8 units/ml heparin. An agonist or antagonist of the invention, and positive controls, such as VEGF and basic FGF (bFGF) are

added, at varying concentrations. On days 4 and 6, the medium is replaced. On day 8, cell number is determined with a Coulter Counter.

[1155] An increase in the number of HUVEC cells indicates that the compound of the invention may proliferate vascular endothelial cells, while a decrease in the number of HUVEC cells indicates that the compound of the invention inhibits vascular endothelial cells.

[1156] The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

Example 26: Rat Corneal Wound Healing Model

[1157] This animal model shows the effect of an agonist or antagonist of the invention on neovascularization. The experimental protocol includes:

- a) Making a 1-1.5 mm long incision from the center of cornea into the stromal layer.
- b) Inserting a spatula below the lip of the incision facing the outer corner of the eye.
- c) Making a pocket (its base is 1-1.5 mm from the edge of the eye).
- d) Positioning a pellet, containing 50ng- 5ug of an agonist or antagonist of the invention, within the pocket.
- e) Treatment with an agonist or antagonist of the invention can also be applied topically to the corneal wounds in a dosage range of 20mg - 500mg (daily treatment for five days).

[1158] The studies described in this example tested activity of agonists or antagonists of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides or polypeptides of the invention (e.g., gene therapy).

Example 27: Diabetic Mouse and Glucocorticoid-Impaired Wound Healing Models

Diabetic db+/db+ Mouse Model.

[1159] To demonstrate that an agonist or antagonist of the invention accelerates the healing process, the genetically diabetic mouse model of wound healing is used. The full thickness wound healing model in the db+/db+ mouse is a well characterized, clinically relevant and

0950083-091201

using a Keyes tissue punch. Immediately following wounding, the surrounding skin is gently stretched to eliminate wound expansion. The wounds are left open for the duration of the experiment. Application of the treatment is given topically for 5 consecutive days commencing on the day of wounding. Prior to treatment, wounds are gently cleansed with sterile saline and gauze sponges.

[1164] Wounds are visually examined and photographed at a fixed distance at the day of surgery and at two day intervals thereafter. Wound closure is determined by daily measurement on days 1-5 and on day 8. Wounds are measured horizontally and vertically using a calibrated Jameson caliper. Wounds are considered healed if granulation tissue is no longer visible and the wound is covered by a continuous epithelium.

[1165] An agonist or antagonist of the invention is administered using at a range different doses, from 4mg to 500mg per wound per day for 8 days in vehicle. Vehicle control groups received 50mL of vehicle solution.

[1166] Animals are euthanized on day 8 with an intraperitoneal injection of sodium pentobarbital (300mg/kg). The wounds and surrounding skin are then harvested for histology and immunohistochemistry. Tissue specimens are placed in 10% neutral buffered formalin in tissue cassettes between biopsy sponges for further processing.

[1167] Three groups of 10 animals each (5 diabetic and 5 non-diabetic controls) are evaluated: 1) Vehicle placebo control, 2) untreated group, and 3) treated group.

[1168] Wound closure is analyzed by measuring the area in the vertical and horizontal axis and obtaining the total square area of the wound. Contraction is then estimated by establishing the differences between the initial wound area (day 0) and that of post treatment (day 8). The wound area on day 1 is 64mm², the corresponding size of the dermal punch. Calculations are made using the following formula:

$$[\text{Open area on day 8}] - [\text{Open area on day 1}] / [\text{Open area on day 1}]$$

[1169] Specimens are fixed in 10% buffered formalin and paraffin embedded blocks are sectioned perpendicular to the wound surface (5mm) and cut using a Reichert-Jung microtome. Routine hematoxylin-eosin (H&E) staining is performed on cross-sections of bisected wounds. Histologic examination of the wounds are used to assess whether the healing process and the morphologic appearance of the repaired skin is altered by treatment with an agonist or antagonist of the invention. This assessment included verification of the presence of cell

accumulation, inflammatory cells, capillaries, fibroblasts, re-epithelialization and epidermal maturity (Greenhalgh, D.G. *et al.*, *Am. J. Pathol.* 136:1235 (1990)). A calibrated lens micrometer is used by a blinded observer.

[1170] Tissue sections are also stained immunohistochemically with a polyclonal rabbit anti-human keratin antibody using ABC Elite detection system. Human skin is used as a positive tissue control while non-immune IgG is used as a negative control. Keratinocyte growth is determined by evaluating the extent of reepithelialization of the wound using a calibrated lens micrometer.

[1171] Proliferating cell nuclear antigen/cyclin (PCNA) in skin specimens is demonstrated by using anti-PCNA antibody (1:50) with an ABC Elite detection system. Human colon cancer served as a positive tissue control and human brain tissue is used as a negative tissue control. Each specimen included a section with omission of the primary antibody and substitution with non-immune mouse IgG. Ranking of these sections is based on the extent of proliferation on a scale of 0-8, the lower side of the scale reflecting slight proliferation to the higher side reflecting intense proliferation.

[1172] Experimental data are analyzed using an unpaired t test. A p value of < 0.05 is considered significant.

Steroid Impaired Rat Model

[1173] The inhibition of wound healing by steroids has been well documented in various *in vitro* and *in vivo* systems (Wahl, Glucocorticoids and Wound healing. In: Anti-Inflammatory Steroid Action: Basic and Clinical Aspects. 280-302 (1989); Wahl *et al.*, *J. Immunol.* 115: 476-481 (1975); Werb *et al.*, *J. Exp. Med.* 147:1684-1694 (1978)). Glucocorticoids retard wound healing by inhibiting angiogenesis, decreasing vascular permeability (Ebert *et al.*, *An. Intern. Med.* 37:701-705 (1952)), fibroblast proliferation, and collagen synthesis (Beck *et al.*, *Growth Factors.* 5: 295-304 (1991); Haynes *et al.*, *J. Clin. Invest.* 61: 703-797 (1978)) and producing a transient reduction of circulating monocytes (Haynes *et al.*, *J. Clin. Invest.* 61: 703-797 (1978); Wahl, "Glucocorticoids and wound healing", In: Antiinflammatory Steroid Action: Basic and Clinical Aspects, Academic Press, New York, pp. 280-302 (1989)). The systemic administration of steroids to impaired wound healing is a well establish phenomenon in rats (Beck *et al.*, *Growth Factors.* 5: 295-304 (1991); Haynes *et al.*, *J. Clin. Invest.* 61: 703-797 (1978); Wahl, "Glucocorticoids and wound healing", In: Antiinflammatory Steroid Action:

Basic and Clinical Aspects, Academic Press, New York, pp. 280-302 (1989); Pierce *et al.*, *Proc. Natl. Acad. Sci. USA* 86: 2229-2233 (1989)).

[1174] To demonstrate that an agonist or antagonist of the invention can accelerate the healing process, the effects of multiple topical applications of the agonist or antagonist on full thickness excisional skin wounds in rats in which healing has been impaired by the systemic administration of methylprednisolone is assessed.

[1175] Young adult male Sprague Dawley rats weighing 250-300 g (Charles River Laboratories) are used in this example. The animals are purchased at 8 weeks of age and are 9 weeks old at the beginning of the study. The healing response of rats is impaired by the systemic administration of methylprednisolone (17mg/kg/rat intramuscularly) at the time of wounding. Animals are individually housed and received food and water *ad libitum*. All manipulations are performed using aseptic techniques. This study is conducted according to the rules and guidelines of Human Genome Sciences, Inc. Institutional Animal Care and Use Committee and the Guidelines for the Care and Use of Laboratory Animals.

[1176] The wounding protocol is followed according to section A, above. On the day of wounding, animals are anesthetized with an intramuscular injection of ketamine (50 mg/kg) and xylazine (5 mg/kg). The dorsal region of the animal is shaved and the skin washed with 70% ethanol and iodine solutions. The surgical area is dried with sterile gauze prior to wounding. An 8 mm full-thickness wound is created using a Keyes tissue punch. The wounds are left open for the duration of the experiment. Applications of the testing materials are given topically once a day for 7 consecutive days commencing on the day of wounding and subsequent to methylprednisolone administration. Prior to treatment, wounds are gently cleansed with sterile saline and gauze sponges.

[1177] Wounds are visually examined and photographed at a fixed distance at the day of wounding and at the end of treatment. Wound closure is determined by daily measurement on days 1-5 and on day 8. Wounds are measured horizontally and vertically using a calibrated Jameson caliper. Wounds are considered healed if granulation tissue is no longer visible and the wound is covered by a continuous epithelium.

[1178] The agonist or antagonist of the invention is administered using at a range different doses, from 4mg to 500mg per wound per day for 8 days in vehicle. Vehicle control groups received 50mL of vehicle solution.

[1179] Animals are euthanized on day 8 with an intraperitoneal injection of sodium pentobarbital (300mg/kg). The wounds and surrounding skin are then harvested for histology.

Tissue specimens are placed in 10% neutral buffered formalin in tissue cassettes between biopsy sponges for further processing.

[1180] Three groups of 10 animals each (5 with methylprednisolone and 5 without glucocorticoid) are evaluated: 1) Untreated group 2) Vehicle placebo control 3) treated groups.

[1181] Wound closure is analyzed by measuring the area in the vertical and horizontal axis and obtaining the total area of the wound. Closure is then estimated by establishing the differences between the initial wound area (day 0) and that of post treatment (day 8). The wound area on day 1 is 64mm², the corresponding size of the dermal punch. Calculations are made using the following formula:

$$[\text{Open area on day 8}] - [\text{Open area on day 1}] / [\text{Open area on day 1}]$$

[1182] Specimens are fixed in 10% buffered formalin and paraffin embedded blocks are sectioned perpendicular to the wound surface (5mm) and cut using an Olympus microtome. Routine hematoxylin-eosin (H&E) staining is performed on cross-sections of bisected wounds. Histologic examination of the wounds allows assessment of whether the healing process and the morphologic appearance of the repaired skin is improved by treatment with an agonist or antagonist of the invention. A calibrated lens micrometer is used by a blinded observer to determine the distance of the wound gap.

[1183] Experimental data are analyzed using an unpaired t test. A p value of < 0.05 is considered significant.

[1184] The studies described in this example tested activity of agonists or antagonists of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides or polypeptides of the invention (e.g., gene therapy).

Example 28: Lymphadema Animal Model

[1185] The purpose of this experimental approach is to create an appropriate and consistent lymphedema model for testing the therapeutic effects of an agonist or antagonist of the invention in lymphangiogenesis and re-establishment of the lymphatic circulatory system in the rat hind limb. Effectiveness is measured by swelling volume of the affected limb, quantification of the amount of lymphatic vasculature, total blood plasma protein, and histopathology. Acute

lymphedema is observed for 7-10 days. Perhaps more importantly, the chronic progress of the edema is followed for up to 3-4 weeks.

[1186] Prior to beginning surgery, blood sample is drawn for protein concentration analysis. Male rats weighing approximately ~350g are dosed with Pentobarbital. Subsequently, the right legs are shaved from knee to hip. The shaved area is swabbed with gauze soaked in 70% EtOH. Blood is drawn for serum total protein testing. Circumference and volumetric measurements are made prior to injecting dye into paws after marking 2 measurement levels (0.5 cm above heel, at mid-pt of dorsal paw). The intradermal dorsum of both right and left paws are injected with 0.05 ml of 1% Evan's Blue. Circumference and volumetric measurements are then made following injection of dye into paws.

[1187] Using the knee joint as a landmark, a mid-leg inguinal incision is made circumferentially allowing the femoral vessels to be located. Forceps and hemostats are used to dissect and separate the skin flaps. After locating the femoral vessels, the lymphatic vessel that runs along side and underneath the vessel(s) is located. The main lymphatic vessels in this area are then electrically coagulated or suture ligated.

[1188] Using a microscope, muscles in back of the leg (near the semitendinosus and adductors) are bluntly dissected. The popliteal lymph node is then located. The 2 proximal and 2 distal lymphatic vessels and distal blood supply of the popliteal node are then ligated by suturing. The popliteal lymph node, and any accompanying adipose tissue, is then removed by cutting connective tissues.

[1189] Care is taken to control any mild bleeding resulting from this procedure. After lymphatics are occluded, the skin flaps are sealed by using liquid skin (Vetbond) (AJ Buck). The separated skin edges are sealed to the underlying muscle tissue while leaving a gap of ~0.5 cm around the leg. Skin also may be anchored by suturing to underlying muscle when necessary.

[1190] To avoid infection, animals are housed individually with mesh (no bedding). Recovering animals are checked daily through the optimal edematous peak, which typically occurred by day 5-7. The plateau edematous peak are then observed. To evaluate the intensity of the lymphedema, the circumference and volumes of 2 designated places on each paw before operation and daily for 7 days are measured. The effect of plasma proteins on lymphedema is determined and whether protein analysis is a useful testing perimeter is also investigated. The weights of both control and edematous limbs are evaluated at 2 places. Analysis is performed in a blind manner.

[1191] Circumference Measurements: Under brief gas anesthetic to prevent limb movement, a cloth tape is used to measure limb circumference. Measurements are done at the ankle bone and dorsal paw by 2 different people and those 2 readings are averaged. Readings are taken from both control and edematous limbs.

[1192] Volumetric Measurements: On the day of surgery, animals are anesthetized with Pentobarbital and are tested prior to surgery. For daily volumetrics animals are under brief halothane anesthetic (rapid immobilization and quick recovery), and both legs are shaved and equally marked using waterproof marker on legs. Legs are first dipped in water, then dipped into instrument to each marked level then measured by Buxco edema software(Chen/Victor). Data is recorded by one person, while the other is dipping the limb to marked area.

[1193] Blood-plasma protein measurements: Blood is drawn, spun, and serum separated prior to surgery and then at conclusion for total protein and Ca^{2+} comparison.

[1194] Limb Weight Comparison: After drawing blood, the animal is prepared for tissue collection. The limbs are amputated using a quillitine, then both experimental and control legs are cut at the ligature and weighed. A second weighing is done as the tibio-cacaneal joint is disarticulated and the foot is weighed.

[1195] Histological Preparations: The transverse muscle located behind the knee (popliteal) area is dissected and arranged in a metal mold, filled with freezeGel, dipped into cold methylbutane, placed into labeled sample bags at - 80EC until sectioning. Upon sectioning, the muscle is observed under fluorescent microscopy for lymphatics..

[1196] The studies described in this example tested activity of agonists or antagonists of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides or polypeptides of the invention (e.g., gene therapy).

Example 29: Suppression of TNF alpha-induced adhesion molecule expression by an Agonist or Antagonist of the Invention

[1197] The recruitment of lymphocytes to areas of inflammation and angiogenesis involves specific receptor-ligand interactions between cell surface adhesion molecules (CAMs) on lymphocytes and the vascular endothelium. The adhesion process, in both normal and pathological settings, follows a multi-step cascade that involves intercellular adhesion molecule-1 (ICAM-1), vascular cell adhesion molecule-1 (VCAM-1), and endothelial leukocyte adhesion

molecule-1 (E-selectin) expression on endothelial cells (EC). The expression of these molecules and others on the vascular endothelium determines the efficiency with which leukocytes may adhere to the local vasculature and extravasate into the local tissue during the development of an inflammatory response. The local concentration of cytokines and growth factor participate in the modulation of the expression of these CAMs.

[1198] Tumor necrosis factor alpha (TNF- α), a potent proinflammatory cytokine, is a stimulator of all three CAMs on endothelial cells and may be involved in a wide variety of inflammatory responses, often resulting in a pathological outcome.

[1199] The potential of an agonist or antagonist of the invention to mediate a suppression of TNF- α induced CAM expression can be examined. A modified ELISA assay which uses ECs as a solid phase absorbent is employed to measure the amount of CAM expression on TNF- α treated ECs when co-stimulated with a member of the FGF family of proteins.

[1200] To perform the experiment, human umbilical vein endothelial cell (HUVEC) cultures are obtained from pooled cord harvests and maintained in growth medium (EGM-2; Clonetics, San Diego, CA) supplemented with 10% FCS and 1% penicillin/streptomycin in a 37 degree C humidified incubator containing 5% CO₂. HUVECs are seeded in 96-well plates at concentrations of 1×10^4 cells/well in EGM medium at 37 degree C for 18-24 hrs or until confluent. The monolayers are subsequently washed 3 times with a serum-free solution of RPMI-1640 supplemented with 100 U/ml penicillin and 100 mg/ml streptomycin, and treated with a given cytokine and/or growth factor(s) for 24 h at 37 degree C. Following incubation, the cells are then evaluated for CAM expression.

[1201] Human Umbilical Vein Endothelial cells (HUVECs) are grown in a standard 96 well plate to confluence. Growth medium is removed from the cells and replaced with 90 μ l of 199 Medium (10% FBS). Samples for testing and positive or negative controls are added to the plate in triplicate (in 10 μ l volumes). Plates are incubated at 37 degree C for either 5 h (selectin and integrin expression) or 24 h (integrin expression only). Plates are aspirated to remove medium and 100 μ l of 0.1% paraformaldehyde-PBS(with Ca⁺⁺ and Mg⁺⁺) is added to each well. Plates are held at 4°C for 30 min.

[1202] Fixative is then removed from the wells and wells are washed 1X with PBS(+Ca,Mg)+0.5% BSA and drained. Do not allow the wells to dry. Add 10 μ l of diluted primary antibody to the test and control wells. Anti-ICAM-1-Biotin, Anti-VCAM-1-Biotin and Anti-E-selectin-Biotin are used at a concentration of 10 μ g/ml (1:10 dilution of 0.1 mg/ml stock

antibody). Cells are incubated at 37°C for 30 min. in a humidified environment. Wells are washed X3 with PBS(+Ca,Mg)+0.5% BSA.

[1203] Then add 20 μ l of diluted ExtrAvidin-Alkaline Phosphotase (1:5,000 dilution) to each well and incubated at 37°C for 30 min. Wells are washed X3 with PBS(+Ca,Mg)+0.5% BSA. 1 tablet of p-Nitrophenol Phosphate pNPP is dissolved in 5 ml of glycine buffer (pH 10.4). 100 μ l of pNPP substrate in glycine buffer is added to each test well. Standard wells in triplicate are prepared from the working dilution of the ExtrAvidin-Alkaline Phosphotase in glycine buffer: 1:5,000 (10^0) > $10^{-0.5}$ > 10^{-1} > $10^{-1.5}$. 5 μ l of each dilution is added to triplicate wells and the resulting AP content in each well is 5.50 ng, 1.74 ng, 0.55 ng, 0.18 ng. 100 μ l of pNPP reagent must then be added to each of the standard wells. The plate must be incubated at 37°C for 4h. A volume of 50 μ l of 3M NaOH is added to all wells. The results are quantified on a plate reader at 405 nm. The background subtraction option is used on blank wells filled with glycine buffer only. The template is set up to indicate the concentration of AP-conjugate in each standard well [5.50 ng; 1.74 ng; 0.55 ng; 0.18 ng]. Results are indicated as amount of bound AP-conjugate in each sample.

[1204] The studies described in this example tested activity of agonists or antagonists of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides or polypeptides of the invention (e.g., gene therapy).

Example 30: Production Of Polypeptide of the Invention For High-Throughput Screening Assays

[1205] The following protocol produces a supernatant containing polypeptide of the present invention to be tested. This supernatant can then be used in the Screening Assays described in Examples 32-41.

[1206] First, dilute Poly-D-Lysine (644 587 Boehringer-Mannheim) stock solution (1mg/ml in PBS) 1:20 in PBS (w/o calcium or magnesium 17-516F Biowhittaker) for a working solution of 50ug/ml. Add 200 μ l of this solution to each well (24 well plates) and incubate at RT for 20 minutes. Be sure to distribute the solution over each well (note: a 12-channel pipetter may be used with tips on every other channel). Aspirate off the Poly-D-Lysine solution and rinse with 1ml PBS (Phosphate Buffered Saline). The PBS should

remain in the well until just prior to plating the cells and plates may be poly-lysine coated in advance for up to two weeks.

[1207] Plate 293T cells (do not carry cells past P+20) at 2×10^5 cells/well in .5ml DMEM(Dulbecco's Modified Eagle Medium)(with 4.5 G/L glucose and L-glutamine (12-604F Biowhittaker))/10% heat inactivated FBS(14-503F Biowhittaker)/1x Penstrep(17-602E Biowhittaker). Let the cells grow overnight.

[1208] The next day, mix together in a sterile solution basin: 300 ul Lipofectamine (18324-012 Gibco/BRL) and 5ml Optimem I (31985070 Gibco/BRL)/96-well plate. With a small volume multi-channel pipetter, aliquot approximately 2ug of an expression vector containing a polynucleotide insert, produced by the methods described in Examples 8-10, into an appropriately labeled 96-well round bottom plate. With a multi-channel pipetter, add 50ul of the Lipofectamine/Optimem I mixture to each well. Pipette up and down gently to mix. Incubate at RT 15-45 minutes. After about 20 minutes, use a multi-channel pipetter to add 150ul Optimem I to each well. As a control, one plate of vector DNA lacking an insert should be transfected with each set of transfections.

[1209] Preferably, the transfection should be performed by tag-teaming the following tasks. By tag-teaming, hands on time is cut in half, and the cells do not spend too much time on PBS. First, person A aspirates off the media from four 24-well plates of cells, and then person B rinses each well with .5-1ml PBS. Person A then aspirates off PBS rinse, and person B, using a 12-channel pipetter with tips on every other channel, adds the 200ul of DNA/Lipofectamine/Optimem I complex to the odd wells first, then to the even wells, to each row on the 24-well plates. Incubate at 37 degree C for 6 hours.

[1210] While cells are incubating, prepare appropriate media, either 1%BSA in DMEM with 1x penstrep, or HGS CHO-5 media (116.6 mg/L of CaCl_2 (anhyd); 0.00130 mg/L $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$; 0.050 mg/L of $\text{Fe}(\text{NO}_3)_3 \cdot 9\text{H}_2\text{O}$; 0.417 mg/L of $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$; 311.80 mg/L of Kcl; 28.64 mg/L of MgCl_2 ; 48.84 mg/L of MgSO_4 ; 6995.50 mg/L of NaCl; 2400.0 mg/L of NaHCO_3 ; 62.50 mg/L of $\text{NaH}_2\text{PO}_4 \cdot \text{H}_2\text{O}$; 71.02 mg/L of Na_2HPO_4 ; .4320 mg/L of $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$; .002 mg/L of Arachidonic Acid ; 1.022 mg/L of Cholesterol; .070 mg/L of DL-alpha-Tocopherol-Acetate; 0.0520 mg/L of Linoleic Acid; 0.010 mg/L of Linolenic Acid; 0.010 mg/L of Myristic Acid; 0.010 mg/L of Oleic Acid; 0.010 mg/L of Palmitric Acid; 0.010 mg/L of Palmitic Acid; 100 mg/L of Pluronic F-68; 0.010 mg/L of Stearic Acid; 2.20 mg/L of Tween 80; 4551 mg/L of D-Glucose; 130.85 mg/ml of L- Alanine;

147.50 mg/ml of L-Arginine-HCL; 7.50 mg/ml of L-Asparagine-H₂O; 6.65 mg/ml of L-Aspartic Acid; 29.56 mg/ml of L-Cystine-2HCL-H₂O; 31.29 mg/ml of L-Cystine-2HCL; 7.35 mg/ml of L-Glutamic Acid; 365.0 mg/ml of L-Glutamine; 18.75 mg/ml of Glycine; 52.48 mg/ml of L-Histidine-HCL-H₂O; 106.97 mg/ml of L-Isoleucine; 111.45 mg/ml of L-Leucine; 163.75 mg/ml of L-Lysine HCL; 32.34 mg/ml of L-Methionine; 68.48 mg/ml of L-Phenylalanine; 40.0 mg/ml of L-Proline; 26.25 mg/ml of L-Serine; 101.05 mg/ml of L-Threonine; 19.22 mg/ml of L-Tryptophan; 91.79 mg/ml of L-Tyrosine-2Na-2H₂O; and 99.65 mg/ml of L-Valine; 0.0035 mg/L of Biotin; 3.24 mg/L of D-Ca Pantothenate; 11.78 mg/L of Choline Chloride; 4.65 mg/L of Folic Acid; 15.60 mg/L of i-Inositol; 3.02 mg/L of Niacinamide; 3.00 mg/L of Pyridoxal HCL; 0.031 mg/L of Pyridoxine HCL; 0.319 mg/L of Riboflavin; 3.17 mg/L of Thiamine HCL; 0.365 mg/L of Thymidine; 0.680 mg/L of Vitamin B₁₂; 25 mM of HEPES Buffer; 2.39 mg/L of Na Hypoxanthine; 0.105 mg/L of Lipoic Acid; 0.081 mg/L of Sodium Putrescine-2HCL; 55.0 mg/L of Sodium Pyruvate; 0.0067 mg/L of Sodium Selenite; 20uM of Ethanolamine; 0.122 mg/L of Ferric Citrate; 41.70 mg/L of Methyl-B-Cyclodextrin complexed with Linoleic Acid; 33.33 mg/L of Methyl-B-Cyclodextrin complexed with Oleic Acid; 10 mg/L of Methyl-B-Cyclodextrin complexed with Retinal Acetate. Adjust osmolarity to 327 mOsm) with 2mm glutamine and 1x penstrep. (BSA (81-068-3 Bayer) 100gm dissolved in 1L DMEM for a 10% BSA stock solution). Filter the media and collect 50 ul for endotoxin assay in 15ml polystyrene conical.

[1211] The transfection reaction is terminated, preferably by tag-teaming, at the end of the incubation period. Person A aspirates off the transfection media, while person B adds 1.5ml appropriate media to each well. Incubate at 37 degree C for 45 or 72 hours depending on the media used: 1%BSA for 45 hours or CHO-5 for 72 hours.

[1212] On day four, using a 300ul multichannel pipetter, aliquot 600ul in one 1ml deep well plate and the remaining supernatant into a 2ml deep well. The supernatants from each well can then be used in the assays described in Examples 32-39.

[1213] It is specifically understood that when activity is obtained in any of the assays described below using a supernatant, the activity originates from either the polypeptide of the present invention directly (e.g., as a secreted protein) or by polypeptide of the present invention inducing expression of other proteins, which are then secreted into the

supernatant. Thus, the invention further provides a method of identifying the protein in the supernatant characterized by an activity in a particular assay.

Example 31: Construction of GAS Reporter Construct

[1214] One signal transduction pathway involved in the differentiation and proliferation of cells is called the Jaks-STATs pathway. Activated proteins in the Jaks-STATs pathway bind to gamma activation site "GAS" elements or interferon-sensitive responsive element ("ISRE"), located in the promoter of many genes. The binding of a protein to these elements alter the expression of the associated gene.

[1215] GAS and ISRE elements are recognized by a class of transcription factors called Signal Transducers and Activators of Transcription, or "STATs." There are six members of the STATs family. Stat1 and Stat3 are present in many cell types, as is Stat2 (as response to IFN-alpha is widespread). Stat4 is more restricted and is not in many cell types though it has been found in T helper class I, cells after treatment with IL-12. Stat5 was originally called mammary growth factor, but has been found at higher concentrations in other cells including myeloid cells. It can be activated in tissue culture cells by many cytokines.

[1216] The STATs are activated to translocate from the cytoplasm to the nucleus upon tyrosine phosphorylation by a set of kinases known as the Janus Kinase ("Jaks") family. Jaks represent a distinct family of soluble tyrosine kinases and include Tyk2, Jak1, Jak2, and Jak3. These kinases display significant sequence similarity and are generally catalytically inactive in resting cells.

[1217] The Jaks are activated by a wide range of receptors summarized in the Table below. (Adapted from review by Schidler and Darnell, Ann. Rev. Biochem. 64:621-51 (1995)). A cytokine receptor family, capable of activating Jaks, is divided into two groups: (a) Class 1 includes receptors for IL-2, IL-3, IL-4, IL-6, IL-7, IL-9, IL-11, IL-12, IL-15, Epo, PRL, GH, G-CSF, GM-CSF, LIF, CNTF, and thrombopoietin; and (b) Class 2 includes IFN-a, IFN-g, and IL-10. The Class 1 receptors share a conserved cysteine motif (a set of four conserved cysteines and one tryptophan) and a WSXWS motif (a membrane proximal region encoding Trp-Ser-Xaa-Trp-Ser (SEQ ID NO: 2)).

[1218] Thus, on binding of a ligand to a receptor, Jaks are activated, which in turn activate STATs, which then translocate and bind to GAS elements. This entire process is encompassed in the Jaks-STATs signal transduction pathway. Therefore, activation of the

Jaks-STATs pathway, reflected by the binding of the GAS or the ISRE element, can be used to indicate proteins involved in the proliferation and differentiation of cells. For example, growth factors and cytokines are known to activate the Jaks-STATs pathway (See Table below). Thus, by using GAS elements linked to reporter molecules, activators of the Jaks-STATs pathway can be identified.

0950033-091201
T02T60-5800560